

integration and more

alexander

November 27, 2025

trigonometric functions and the inverse

to obtain the inverse of a function, the function must be one-to-one, meaning that each input corresponds to exactly one unique output and no two different inputs share the same output value. however, the trigonometric functions such as sine, cosine, and tangent are not one-to-one over their entire domains because they are periodic and repeat their values infinitely many times. therefore, to define their inverses $\arcsin(x)$, $\arccos(x)$, and $\arctan(x)$ we must restrict the domain of each trigonometric function to an interval where it passes the horizontal line test. this ensures that each inverse function is well-defined and produces a single, unique output for every input within its range.

- $y = \sin(x)$, domain: $x \in (-\infty, \infty)$, range: $y \in [-1, 1]$, period: 2π
 $y = \arcsin(x)$, domain: $x \in [-1, 1]$, range: $y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
- $y = \cos(x)$, domain: $x \in (-\infty, \infty)$, range: $y \in [-1, 1]$, period: 2π
 $y = \arccos(x)$, domain: $x \in [-1, 1]$, range: $y \in [0, \pi]$
- $y = \tan(x) = \frac{\sin(x)}{\cos(x)}$, domain: $x \neq \frac{\pi}{2}, n \in \mathbb{Z}$, range: $y \in (-\infty, \infty)$, period: π
 $y = \arctan(x)$, domain: $x \in (-\infty, \infty)$, range: $y \in (-\frac{\pi}{2}, \frac{\pi}{2})$

FTC

- if f is a continuous function on an interval $[a, b]$, and $F(x) = \int_a^x f(t) dt$ is defined for $x \in [a, b]$, then $F'(x) = f(x)$
- if f is continuous function on $[a, b]$, and F is any antiderivative of f , meaning $F'(x) = f(x)$, then $\int_a^b f(x) dx = F(b) - F(a)$

integration power rule

- $\frac{d}{dx} [\frac{x^{n+1}}{n+1}] = x^n \Rightarrow \int x^n dx = \frac{x^{n+1}}{n+1} + C$

u-substitution

- $\int f(g(x))g'(x) dx \Rightarrow \int f(u) du$
- $\int_a^b f(g(x))g'(x) dx = \int_{u(a)}^{u(b)} f(u) du$

integration by parts

- $(uv)'(x) = u'(x)v(x) + u(x)v'(x) \Rightarrow \int u dv = uv - \int v du$

trigonometric substitution

1. $\sqrt{a^2 - b^2 x^2} = \sqrt{a^2(1 - \frac{b^2 x^2}{a^2})} = \sqrt{a^2(1 - (\frac{bx}{a})^2)} = \sqrt{a^2(1 - (\frac{b}{a}x)^2)} = a\sqrt{1 - (\frac{b}{a}x)^2}$
2. $\sin^2(\theta) = \cos^2(\theta) = 1 \Leftrightarrow \cos(\theta) = \sqrt{1 - (\sin(\theta))^2}$
3. $\frac{b}{a}x = \sin(\theta) \Leftrightarrow x = \frac{a}{b} \sin(\theta)$
4. $a \cos(\theta) = \sqrt{a^2 - b^2 x^2}$

when you do a trig substitution, you often let something like $x = \sin(\theta)$. that automatically restricts θ to the range of the inverse sine function, which is $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$. within this range, $\sin(\theta)$ can be positive or negative, but $\cos(\theta)$ is always nonnegative. now, when you use the pythagorean identity $\sin^2(\theta) + \cos^2(\theta) = 1$, and solve for $\cos(\theta)$, you get $\cos(\theta) = \pm\sqrt{1 - \sin^2(\theta)}$. mathematically, both the positive and negative square roots are valid - but you have to choose which sign is correct for the range of θ you are working in. because we restricted θ to $[-\frac{\pi}{2}, \frac{\pi}{2}]$ when we said $x = \sin(\theta)$, we know $\cos(\theta) \geq 0$ there. Therefore, we choose the positive square root: $\cos(\theta) = +\sqrt{1 - \sin^2(\theta)}$. if instead we had chosen a substitution involving $\cos(\theta)$, we would pick a domain where $\sin(\theta)$ has a definite sign and make the corresponding choice.

integration by partial fraction decomposition

- distinct linear factors $\frac{A}{x-a} + \frac{B}{x-b}$
- repeated linear factors $\frac{A_1}{x-a} + \frac{A_2}{(x-a)^2} + \dots + \frac{A_n}{(x-a)^n}$
- irreducible quadratic factors $\frac{Ax+B}{x^2+bx+c}$
- repeated quadratic factors $\frac{A_1x+B_1}{x^2+bx+c} + \dots + \frac{A_nx+B_n}{(x^2+bx+c)^n}$

a rational function is of the form $\frac{P(x)}{Q(x)}$ where $P(x)$ and $Q(x)$ are polynomials. with a proper fraction ($\deg(P) < \deg(Q)$) you can go straight to the decomposition, but for something improper ($\deg(P) > \deg(Q)$), you will need to use polynomial division ($\frac{P(x)}{Q(x)} = S(x) + \frac{R(x)}{Q(x)}$) where $S(x)$ is a polynomial (the quotient), and $\frac{R(x)}{Q(x)}$ is now a proper fraction (remainder over the denominator).

riemann sums

- given a definite integral like so: $\int_a^b f(x) dx$ you can approximate it by breaking $[a, b]$ into smaller subintervals with a width of $\Delta x = \frac{b-a}{n}$. you now have subintervals like so: $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$ where $x_i = a + i\Delta x$.

- $L_n = \sum_{i=0}^{n-1} f(x_i) \cdot \Delta x$
- $R_n = \sum_{i=1}^n f(x_i) \cdot \Delta x$
- $M_n = \sum_{i=1}^n f(m_i) \cdot \Delta x$ where $m_i = \frac{x_{i-1} + x_i}{2}$
- $T_n = \sum_{i=1}^n \frac{f(x_{i-1}) + f(x_i)}{2} \cdot \Delta x$

whats next?

1. probability
2. arrangement, permutation, combinations
3. pascals identity and the binomial theorem
4. logs and exponential functions
5. $A = P(1 + \frac{r}{n})^{nt} \Rightarrow A = Pe^{rt} \Rightarrow e = \lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n$
6. doubling time, half life,
7. $e^{i\pi} + 1 = 0$, hyperbolic functions
8. signed areas, area between curves, average value of a function, volumes of revolution
9. numeric integration techniques, improper integrals, arc lengths
10. taylor polynomials
11. first order linear equations
12. sequences, convergence, power series, taylor series
13. polar coords