

calculus is a branch of mathematics that deals with rates of change and the accumulation of quantities.

May 5, 2025

circle: $(x - a)^2 + (y - b)^2 = r^2$ where (a, b) is the center and the radius is r

midpoint between $P_1 = (x_1, y_1)$ and $P_2 = (x_2, y_2)$ is P_1P_2 divided by 2

distance: $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

laws of exponents:

- $x^m x^n = x^{m+n}$
- $\frac{x^m}{x^n} = x^{m-n}$
- $(x^m)^n = x^{mn}$
- $x^{-n} = \frac{1}{x^n}$
- $(xy)^n = x^n y^n$
- $(\frac{x}{y})^n = \frac{x^n}{y^n}$
- $x^{1/n} = \sqrt[n]{x}$
- $x^{\frac{m}{n}} = \sqrt[n]{x^m} = (\sqrt[n]{x})^m$

special factorizations

- $x^2 - y^2 = (x + y)(x - y)$
- $x^3 + y^3 = (x + y)(x^2 - xy + y^2)$
- $x^3 - y^3 = (x - y)(x^2 + xy + y^2)$

binomial theorem:

- $(x + y)^2 = x^2 + 2xy + y^2$
- $(x - y)^2 = x^2 - 2xy + y^2$
- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
- $(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$
- $(x + y)^n = x^n + nx^{n-1}y + \frac{n(n-1)}{2}x^{n-2}y^2 + \dots + \binom{n}{k}x^{n-k}y^k + \dots + nxy^{n-1} + y^n$
where $\binom{n}{k} = \frac{n(n-1)\dots(n-k+1)}{1 \cdot 2 \cdot 3 \cdot \dots \cdot k}$

quadratic formulae:

1. $ax^2 + bx + c = 0$
2. $x^2 + \frac{b}{a}x + \frac{c}{a} = 0$
3. $x^2 + \frac{b}{a}x = -\frac{c}{a}$
4. $x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$
5. $\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2}$
6. $x + \frac{b}{2a} = \pm \frac{\sqrt{b^2 - 4ac}}{2a}$
7. $x = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$
8. $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

polynomials $a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$ where n is non-neg and represents the degree

rational $p(x)/q(x)$

root $\sqrt[n]{g(x)}$

properties:

domain: set of all input values for which the function is defined


range: set of all possible output values of the function

continuity: most algebraic functions are continuous (no breaks or jumps), but rational functions have discontinuities at points

behavior: the functions behavior is influenced by the degree of the polynomial and the nature of the function

scaling:

- vertical scaling $y = kf(x)$: If $k \geq 1$, the graph is expanded vertically by the factor k . If $0 < k < 1$, the graph is compressed vertically. When the scale factor k is negative ($k < 0$), the graph is also reflected across the x-axis.
- horizontal scaling $y = f(kx)$: If $K \geq 1$, the graph is compressed in the horizontal direction. If $0 < k < 1$, the graph is expanded. If $k \leq 0$, then the graph is also reflected across the y-axis.



Angle (Degrees)	Angle (Radians)	$\cos(\theta)$	$\sin(\theta)$
0°	0	1	0
30°	$\frac{\pi}{6}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$
45°	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$
60°	$\frac{\pi}{3}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$
90°	$\frac{\pi}{2}$	0	1
120°	$\frac{2\pi}{3}$	$-\frac{1}{2}$	$\frac{\sqrt{3}}{2}$
135°	$\frac{3\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$
150°	$\frac{5\pi}{6}$	$-\frac{\sqrt{3}}{2}$	$\frac{1}{2}$
180°	π	-1	0
210°	$\frac{7\pi}{6}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$
225°	$\frac{5\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$
240°	$\frac{4\pi}{3}$	$-\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$
270°	$\frac{3\pi}{2}$	0	-1
300°	$\frac{5\pi}{3}$	$\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$
315°	$\frac{7\pi}{4}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$
330°	$\frac{11\pi}{6}$	$\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$
360°	2π	1	0

$$1 \text{ deg} = \frac{\pi}{180} \text{ rad}$$

trigonometric functions are mathematical functions that relate the angle of a triangle to the lengths of its sides... and can also be generalized to all real numbers using the unit circle.

To derive the rest of the fundamental trigonometric identities, you need a combination of a few key identities and principles. The most important starting point is the Pythagorean identity, but you'll also need the basic relationships between the trigonometric functions, such as the definitions of sine, cosine, tangent, secant, cosecant, and cotangent in terms of a right triangle or the unit circle.

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\sec \theta = \frac{1}{\cos \theta}, \quad \csc \theta = \frac{1}{\sin \theta}, \quad \cot \theta = \frac{1}{\tan \theta}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}, \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$$

$$\tan(2\theta) = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

$$\sin^2(2\theta) = \frac{1 - \cos^2(2\theta)}{2}$$

$$\cos^2(2\theta) = \frac{1 + \cos(2\theta)}{2}$$

$$\sin(90^\circ - \theta) = \cos \theta, \quad \cos(90^\circ - \theta) = \sin \theta$$

$$\tan(90^\circ - \theta) = \cot \theta, \quad \cot(90^\circ - \theta) = \tan \theta$$

$$\sec(90^\circ - \theta) = \csc \theta, \quad \csc(90^\circ - \theta) = \sec \theta$$

$$\sin(-\theta) = -\sin(\theta), \quad \cos(-\theta) = \cos(\theta)$$

$$\tan(-\theta) = -\tan(\theta), \quad \sec(-\theta) = \sec(\theta)$$

$$\csc(-\theta) = -\csc(\theta), \quad \cot(-\theta) = -\cot(\theta)$$

$$\sin \alpha \sin \beta = \frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2}[\cos(\alpha - \beta) + \cos(\alpha + \beta)]$$

$$\sin \alpha \cos \beta = \frac{1}{2}[\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

$$\text{law of sines: } \frac{\sin(A)}{a} = \frac{\sin(B)}{b} = \frac{\sin(C)}{c} \text{ (uppercase are the angles)}$$

$$\text{law of cosines: } a^2 = b^2 + c^2 - 2bc \cos(A)$$

power functions: $f(x) = x^n$...if even the function behaves symmetrical around the y-axis...if odd then the function has point symmetry (x^4 , x^3 , $x^{-n} = \frac{1}{x^n}$)

inverse trig functions: $\arcsin(x) = \sin^{-1}(x) = \theta$, $\arccos(x) = \cos^{-1}(x) = \theta$, $\arcsin(x) = \sin^{-1}(x) = \theta$

logs: $\log_a x = y \leftrightarrow a^y = x$, $\ln(x) = y \leftrightarrow e^y = x$

hyperbolic functions: $\sinh(x) = \frac{e^x - e^{-x}}{2}$, $\cosh(x) = \frac{e^x + e^{-x}}{2}$, $\tanh(x) = \frac{\sinh(x)}{\cosh(x)}$

differentiation rules:

1. $\frac{d}{dx}(c) = 0$
2. $\frac{d}{dx}x = 1$
3. $\frac{d}{dx}(x^n) = nx^{n-1}$ (power rule)
4. $\frac{d}{dx}[cf(x)] = cf'(x)$
5. $\frac{d}{dx}[f(x) + g(x)] = f'(x) + g'(x)$
6. $\frac{d}{dx}[f(x)g(x)] = f(x)g'(x) + g(x)f'(x)$ (product rule)
7. $\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$ (quotient rule)
8. $\frac{d}{dx}f(g(x)) = f'(g(x))g'(x)$ (chain rule)
9. $\frac{d}{dx}f(x)^n = nf(x)^{n-1}f'(x)$ (general power rule)
10. $\frac{d}{dx}\sin(x) = \cos(x)$
11. $\frac{d}{dx}\cos(x) = -\sin(x)$
12. $\frac{d}{dx}\tan(x) = \sec^2(x)$
13. $\frac{d}{dx}\csc(x) = -\csc(x)\cot(x)$
14. $\frac{d}{dx}\sec(x) = \sec(x)\tan(x)$
15. $\frac{d}{dx}\cot(x) = -\csc^2(x)$
16. $\frac{d}{dx}\sin^{-1}(x) = \frac{1}{\sqrt{1-x^2}}$
17. $\frac{d}{dx}\cos^{-1}(x) = -\frac{1}{\sqrt{1-x^2}}$

$$18. \frac{d}{dx} \tan^{-1}(x) = \frac{1}{1+x^2}$$

$$19. \frac{d}{dx} (e^x) = e^x$$

$$20. \frac{d}{dx} (a^x)(\ln a)a^x$$

$$21. \frac{d}{dx} \ln |x| = \frac{1}{x}$$

$$22. \frac{d}{dx} \log_a x = \frac{1}{(\ln a)x}$$

Essential Theorems:

- Intermediate Value Theorem
- Mean Value Theorem
- Extreme Values on a Closed Interval
- The Fundamental Theorem of Calculus, Part I
- Fundamental Theorem of Calculus, Part II

review problems

verifying trigonometric identities
solving trigonometric equations

direct substitution
indeterminate forms and algebraic simplification
trigonometric limits
piecewise functions
infinite limits and vertical asymptotes

squeeze theorem
epsilon-delta

basic polynomial differentiation
implicit differentiation
higher-order derivatives
applications of derivatives (tangent line equations, critical points, motion problems, related rates)

basic antiderivatives
u-sub (reverse chain rule)
trig integrals

partial fractions
applications (area under curves, average value of a function, volume by disks/washers/shells, accumulation problems)