

0.1 Properties of Complex Numbers (★)

Complex numbers are just as complex as the name suggests, but they are so so elegant when used properly - from circles to trigonometric expressions, complex numbers link them all.




Figure 1: Comic from <https://xkcd.wtf/2028/>

0.1 Properties of Complex Numbers (★)

Complex numbers can be written in a few ways, and each of these ways have their own benefits that make information more apparent.

Proposition 0.1 (Notations for Complex Numbers)

Let z be a complex number and let i be the imaginary number such that $i^2 = -1$. We write $z = a + bi$ for real a and b .

1. The *real part* of z is denoted as $Re(z)$ or $\Re(z)$. For instance, $\Re(z) = a$.
2. The *imaginary part* of z is denoted as $Im(z)$ or $\Im(z)$. For instance, $\Im(z) = b$.
3. The *modulus* of z is denoted as $|z|$. For instance, $|z| = \sqrt{a^2 + b^2}$.
4. The *argument* of z is denoted as $\arg z$. For instance, $\arg z = \tan^{-1}\left(\frac{b}{a}\right)$.
5. The *polar form* of z can be written as

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta) = r \operatorname{cis} \theta.$$

Then, $|z| = r$ and $\arg z = \theta$.

6. The *conjugate* of z is denoted as \bar{z} or z^* . For instance,

$$\bar{z} = a - bi = re^{-i\theta} = r(\cos \theta - i \sin \theta).$$

Remark 0.2. Note that conjugation is distributive. For instance,

1. $\overline{z \pm w} = \bar{z} \pm \bar{w}$,
2. $\overline{z \cdot w} = \bar{z} \cdot \bar{w}$,
3. $\overline{\frac{z}{w}} = \frac{\bar{z}}{\bar{w}}$.

With the different representations for complex numbers in mind, we shall briefly introduce some properties that relate to each form. As again, let

$$z = a + bi = re^{i\theta} = r(\cos \theta + i \sin \theta).$$

Proposition 0.3 (Absolute Square)

$$z \cdot \bar{z} = |z|^2.$$

Proof.

$$\begin{aligned} z \cdot \bar{z} &= (a + bi)(a - bi) = a^2 - (bi)^2 \\ &= a^2 + b^2 = (\sqrt{a^2 + b^2})^2 \\ &= |z|^2. \end{aligned}$$

□

Proposition 0.4 (Rotation)

Multiplying a complex number by $e^{i\theta}$ rotates the vector by θ radians counterclockwise about the origin.

0.2 Exercise

Problem 0.5. Prove that $\arg(zw) = \arg(z) + \arg(w)$ and $\arg\left(\frac{z}{w}\right) = \arg(z) - \arg(w)$.

Problem 0.6. De Moivre's theorem states that

$$(\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta).$$

Prove de Moivre's theorem by induction.

0.3 Roots of Unity (★★)

Let n be a positive integer. An n th root of unity is a complex number z that satisfies the equation

$$z^n = 1.$$

Then, the n th roots of unity can be described by

$$z = \exp\left(\frac{2k\pi}{n}\right) = \cos \frac{2k\pi}{n} + i \sin \frac{2k\pi}{n}, \quad k = 0, 1, \dots, n-1.$$

Proposition 0.7 (Conjugates)

If z satisfies $z^n = 1$, meaning z is an n th root of unity, then so does \bar{z} .

Proof. Recall that $z \cdot \bar{z} = |z|^2 = 1$, so $\bar{z} = \frac{1}{z} = z$.

□

To start off the section, we showcase a trick on evaluating (suspiciously convenient) trigonometric sums through the lenses of the **roots of unity**.

0.3 Roots of Unity (★★)

Example 0.8 (2021 H2 Further Math P1/9)

Let $\omega = \cos \frac{2\pi}{11} + \sin \frac{2\pi}{11}$. Show that

$$\sum_{j=1}^{10} \omega^j = -1.$$

Simple enough, ω is an 11th roots of unity, and so it satisfies the polynomial $z^{11} - 1 = 0$. Moreover, the required sum is a geometric progression, so

$$\sum_{j=1}^{10} \omega^j = \frac{\omega^{11} - 1}{\omega - 1} - 1 = -1.$$

Example 0.9 (cont.)

The complex numbers α and β are such that

$$\alpha = w + w^3 + w^4 + w^5 + w^9 \text{ and } \beta = w^{-1} + w^{-3} + w^{-4} + w^{-5} + w^{-9}.$$

Find in its simplest form, the quadratic equation whose roots are α and β .

Most jarringly, the exponents for α don't come in order: it jumps from 1 to 3, then 5 to 9 and β mirrors this behaviour, except with negative exponents. Fortunately, we see that $w^9 = e^{\frac{18\pi i}{11}} = e^{\frac{-4\pi i}{11}} = w^{-2}$ and that $\Re(w^{-2}) = \Re(w^2)$ (why doesn't the problem use w^2 and w^{-2} instead?). We also make a mental note that $\alpha = \beta^*$. Maybe this will come in handy later.

For now, we have

$$\begin{aligned} \alpha + \beta &= (w + w^{-1}) + (w^3 + w^{-3}) + (w^4 + w^{-4}) + (w^5 + w^{-5}) + (w^9 + w^{-9}) \\ &= 2\Re(w + w^3 + w^4 + w^5 + w^9) = 2\Re(w + w^2 + w^3 + w^4 + w^5) \\ &= 2 \left(\cos \frac{2\pi}{11} + \cos \frac{4\pi}{11} + \cos \frac{6\pi}{11} + \cos \frac{8\pi}{11} + \cos \frac{10\pi}{11} \right) \end{aligned}$$

This is usually a tough sum to evaluate, but the values here just line up way too nicely for us to ignore... With that, here comes the *trick*:

$$S = \cos \frac{2\pi}{11} + \cos \frac{4\pi}{11} + \cos \frac{6\pi}{11} + \cos \frac{8\pi}{11} + \cos \frac{10\pi}{11}.$$

We consider

$$\begin{aligned} \frac{S \cdot 2 \sin \frac{2\pi}{11}}{2 \sin \frac{2\pi}{11}} &= \frac{2 \sin \frac{2\pi}{11} \cos \frac{2\pi}{11} + 2 \sin \frac{2\pi}{11} \cos \frac{4\pi}{11} + \cdots + 2 \sin \frac{2\pi}{11} \cos \frac{10\pi}{11}}{2 \sin \frac{2\pi}{11}} \\ &= \frac{\sin \frac{10\pi}{11} + \sin \frac{12\pi}{11} - \sin \frac{2\pi}{11}}{2 \sin \frac{2\pi}{11}} \quad \text{using the product-to-sum formula} \\ &= -\frac{1}{2} \end{aligned}$$

Remark 0.10. This doesn't work if we have one fewer term though. For instance if we exclude $\cos \frac{10\pi}{11}$, then the sum doesn't come out nice anymore. Hmm...

0.3 Roots of Unity (★★)

Hence, $\alpha + \beta = -1$. Furthermore, we have $w^{-i} = w^{-11+i}$ and $w^i = w^{11-i}$, so that

$$\begin{aligned}\alpha\beta &= (w + w^3 + w^4 + w^5 + w^9)(w^{-1} + w^{-3} + w^{-4} + w^{-5} + w^{-9}) \\ &= 5 + w^{-8} + w^{-6} + w^{-5} + 2w^{-4} + w^{-3} + 2w^{-2} + 2w^{-1} + 2w + 2w^2 + w^3 + 2w^4 + w^5 + w^6 + w^8 \\ &= 5 + (w^{-10} + w^{-9} + \dots + w^{-1}) + (w + w^2 + \dots + w^{10}) \\ &= 5 + \sum_{j=1}^{10} \omega^j + \left(\sum_{j=1}^{10} \omega^j \right)^* = 3\end{aligned}$$

where the final sum holds because

$$\left(\sum_{j=1}^{10} \omega^j \right) \left(\sum_{j=1}^{10} \omega^j \right)^* = \left| \left(\sum_{j=1}^{10} \omega^j \right) \right|^2 = 1$$

Hence, a possible quadratic equation is $x^2 - x + 3 = 0$.

Example 0.11 (cont..)

Prove that $|\alpha| = \sqrt{3}$

Aha! Here's where the property that $\alpha = \beta^*$ comes in handy.

$$|\alpha|^2 = \alpha \cdot \alpha^* = \alpha\beta = 3$$

and the result follows.

Example 0.12 (cont...)

Prove that

$$\sin \frac{2\pi}{11} + \sin \frac{6\pi}{11} + \sin \frac{8\pi}{11} + \sin \frac{10\pi}{11} + \sin \frac{18\pi}{11} = \frac{\sqrt{11}}{2}.$$

Observe that this is simply the imaginary part of α . In addition,

$$\begin{aligned}\Im(\alpha) &= \sin \frac{2\pi}{11} + \sin \frac{6\pi}{11} + \sin \frac{8\pi}{11} + \sin \frac{10\pi}{11} + \sin \frac{18\pi}{11} \\ &= \sin \frac{2\pi}{11} + \sin \frac{6\pi}{11} + \left(\sin \frac{8\pi}{11} - \sin \frac{7\pi}{11} \right) + \sin \frac{10\pi}{11} \\ &> 0\end{aligned}$$

since each term is positive. Thus,

$$\begin{aligned}\Im(\alpha) &= \frac{1}{2} \cdot 2\Im(\alpha) = \frac{1}{2} (\alpha - \alpha^*) \\ &= \frac{1}{2} (\alpha - \beta) \\ &= \frac{1}{2} \Im \left(\sqrt{(\alpha + \beta)^2 - 4\alpha\beta} \right) \\ &= \frac{\sqrt{11}}{2}\end{aligned}$$

by taking the positive square root because $\alpha - \beta = 2\Im(\alpha) > 0$.

Tangent: Gaussian Periods So why didn't the question use w^{-2} and w^2 ? The set $(1, 3, 4, 5, 9)$ feels too specific anyways. As it turns out, this is the set of **quadratic residues** modulo 11 (prove this!), and the sums α and β are the so-called quadratic Gauss sums:

Definition 0.13 (Quadratic Gauss Sum).

$$g_a = \sum_{t=0}^{p-1} \left(\frac{t}{p} \right) \zeta^{at}$$

where $\left(\frac{t}{p} \right)$ is the Legendre's symbol

Definition 0.14 (Legendre's symbol). $\left(\frac{t}{p} \right) = \begin{cases} 1 & \text{if } t \text{ is a quadratic residue,} \\ -1 & \text{if } t \text{ is a quadratic non-residue,} \\ 0 & \text{if } p|t. \end{cases}$

In what follows, we showcase a technique of *summing in two ways* that reduces this problem to a sum of roots of unity.

Proposition 0.15 (Expressing g_a in terms of g_1)

$$g_a = \left(\frac{a}{p} \right) g_1.$$

Proof. Firstly if $p|a$, then $g_a = 0$ (why?).

Henceforth, suppose p does not divide a . Then,

$$\left(\frac{a}{p} \right) g_a = \sum_{t=0}^{p-1} \left(\frac{at}{p} \right) \zeta^{at} = \sum_{x=0}^{p-1} \left(\frac{x}{p} \right) \zeta^x = g_1.$$

The first equality follows because $a \nmid p \implies at \nmid p$, so that as t runs over the set of residues (mod p), then so does at . Since at runs over the set of residues, then we may simply let the sum run over all x in the set of residues (mod p).

Now observe by definition that $\left(\frac{a}{p} \right)^2 = 1$, and we are done. \square

What follows is the *coup de grace*:

Proposition 0.16 (Summing in two ways)

$$g_1^2 = (-1)^{\frac{p-1}{2}} p.$$

For convenience, write $g = g_1$. According to Ireland-Rosen, the main idea is to consider the sum $\sum_{a=0}^{p-1} g_a g_{-a}$ in two ways. Again, we consider only the case when $p \nmid a$.

On one hand, we have

$$g_a g_{-a} = \left(\frac{a}{p} \right) \left(\frac{-a}{p} \right) g^2 = \left(\frac{-1}{p} \right) g^2,$$

whence it follows that

$$\sum_{a=1}^{p-1} g_a g_{-a} = \left(\frac{-1}{p} \right) (p-1) g^2.$$

On the other hand, we have

$$g_a g_{-a} = \sum_{x=0}^{p-1} \sum_{y=0}^{p-1} \left(\frac{x}{p}\right) \left(\frac{y}{p}\right) \zeta^{a(x-y)},$$

and summing both sides over a ,

$$\begin{aligned} \sum_{a=0}^{p-1} g_a g_{-a} &= \sum_{a=0}^{p-1} \sum_{x=0}^{p-1} \sum_{y=0}^{p-1} \left(\frac{x}{p}\right) \left(\frac{y}{p}\right) \zeta^{a(x-y)} \\ &= \sum_{x=0}^{p-1} \sum_{y=0}^{p-1} \left(\frac{x}{p}\right) \left(\frac{y}{p}\right) \sum_{a=0}^{p-1} \zeta^{a(x-y)} \\ &= \sum_{x=0}^{p-1} \sum_{y=0}^{p-1} \left(\frac{x}{p}\right) \left(\frac{y}{p}\right) s(x, y) p \\ &= p(p-1) \end{aligned}$$

where we have $p^{-1} \sum_{t=0}^{p-1} \zeta^{at} = s(x, y) = \begin{cases} 1 & \text{if } p \mid t \\ 0 & \text{otherwise.} \end{cases}$. Putting both parts together, the result follows.