

Trigonometric identities are *triggier* than you may realise! However, they are also useful in many ways, from parametrisations to geometry, and even bridging over to complex numbers.




Figure 1: Comic from <https://xkcd.wtf/2028/>

### Proposition 0.1 (Trigonometric Identities)

Let  $A$  and  $B$  denote angles.

1. *Pythagorean Identity*:  $\sin^2 A + \cos^2 A = 1$

(a)  $\tan^2 A + 1 = \sec^2 A$ ,

(b)  $\cot^2 A + 1 = \csc^2 A$ .

2. *Addition Formulae*

(a)  $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$ ,

(b)  $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$ ,

(c)  $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$ .

3. *Double Angle Formulae*

(a)  $\sin 2A = 2 \sin A \cos A$ ,

(b)  $\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$ ,

(c)  $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$ .

4. *Sum-to-Product Formulae*

(a)  $\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$ ,

(b)  $\sin A - \sin B = 2 \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$ ,

(c)  $\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$ ,

(d)  $\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$ .

---

**Proposition 0.2** (Trigonometric R Method (or R-formula))

Let  $a, b, \theta$  be reals. The R-formula expresses the *weighted sum*  $a \sin \theta + b \cos \theta$  as a single sin or cos term.

Letting  $R = \sqrt{a^2 + b^2}$  and  $\tan \alpha = \frac{b}{a}$ ,

1. *Sine form:*  $a \sin \theta \pm b \cos \theta = R \sin(\theta \pm \alpha)$ .
2. *Cosine form:*  $a \cos \theta \pm b \sin \theta = R \cos(\theta \mp \alpha)$ .

# 1 Applying Identities (★)

For fundamentals, it is important that we are able to simplify complicated expressions into simpler forms.

## Example 1.1

Simplify  $\cos 49^\circ + \cos 71^\circ$  as a single trigonometric term.

*Proof.* Observe the symmetry about 60:  $49 = 60 - 11$  and  $71 = 60 + 11$ , so

$$\begin{aligned}\cos 49^\circ + \cos 71^\circ &= \cos(60 - 11)^\circ + \cos(60 + 11)^\circ \\ &= 2 \cos 60^\circ \cos 11^\circ \\ &= \boxed{\cos 11^\circ}.\end{aligned}$$

□

## Proposition 1.2 (Toolkit)

There usually isn't much to do with trigonometry. Most useful properties relate the sum or difference of angles, and so the usual techniques go by:

1. squaring and adding the given relations,
2. squaring and subtracting the given relations,
3. simply adding or subtracting the given relations,
4. multiplying the given relations,
5. using sum-to-product formulae, or
6. expressing the required expression in terms of the given angles.

## Example 1.3 (2020 SMO(S) P9)

If  $8 \cos x - 8 \sin x = 3$ , find the value of  $\tan x + \frac{1}{\tan x}$ .

*Proof.* Firstly,  $(8 \cos x - 8 \sin x)^2 = 64 - 64 \sin 2x = 9 \implies \sin 2x = \frac{55}{64}$ .

Moreover,

$$\tan x + \frac{1}{\tan x} = \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{2}{\sin 2x} = \boxed{\frac{128}{55}}.$$

□

## Example 1.4 (2021 SMO(O) P1)

It is given that  $\frac{\pi}{2} < \beta < \alpha < \frac{3\pi}{4}$ ,  $\cos(\alpha - \beta) = \frac{12}{13}$  and  $\sin(\alpha + \beta) = -\frac{3}{5}$ . Find  $\lfloor |2021 \sin 2\alpha| \rfloor$ .

*Proof.* From the angle conditions, we have  $0 < \alpha - \beta < \frac{\pi}{4}$  and  $\pi < \alpha + \beta < \frac{3\pi}{2}$ . We now pull a big sneaky! Note that

$$\begin{aligned}\sin 2\alpha &= \sin((\alpha - \beta) + (\alpha + \beta)) \\ &= \sin(\alpha - \beta) \cos(\alpha + \beta) + \cos(\alpha - \beta) \sin(\alpha + \beta) \\ &= \frac{5}{13} \cdot \left(-\frac{4}{5}\right) + \frac{12}{13} \cdot \left(-\frac{3}{5}\right) \\ &= -\frac{4}{13} - \frac{36}{65} \\ &= -\frac{56}{65}.\end{aligned}$$

Thus,

$$\lfloor 2021 \sin 2\alpha \rfloor = \left\lfloor \frac{2021 \cdot 56}{65} \right\rfloor = \left\lfloor \frac{113176}{65} \right\rfloor = \boxed{1741}.$$

□

**Example 1.5** (2021 SMO(O) P12)

Given that  $\sin \alpha + \sin \beta = \frac{1}{10}$  and  $\cos \alpha + \cos \beta = \frac{1}{9}$ . Find the value of  $\lfloor \tan^2(\alpha + \beta) \rfloor$ .

*Proof.* By the sum-to-product formulae,

$$\frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta} = \tan \left( \frac{A+B}{2} \right) = \frac{9}{10}.$$

Thus,

$$\begin{aligned}\lfloor \tan^2(\alpha + \beta) \rfloor &= \left\lfloor \left( \frac{2 \tan \left( \frac{A+B}{2} \right)}{1 - \tan^2 \left( \frac{A+B}{2} \right)} \right)^2 \right\rfloor \\ &= \left\lfloor \left( \frac{2 \cdot \frac{9}{10}}{1 - \frac{81}{100}} \right)^2 \right\rfloor \\ &= \left\lfloor \left( \frac{180}{19} \right)^2 \right\rfloor = \left\lfloor \frac{32400}{361} \right\rfloor \\ &= \boxed{89}.\end{aligned}$$

□

**Example 1.6** (2021 SMO(S) P7)

If  $\cos A - \cos B = \frac{1}{2}$  and  $\sin A - \sin B = -\frac{1}{4}$ , find the value of  $100 \sin(A + B)$ .

*Proof.* With some foresight, we notice that squaring and then summing or subtracting the relations wouldn't be too useful at this point.

We turn to the sum-to-product formulae:

$$\cos A - \cos B = \frac{1}{2} \implies \sin \frac{A+B}{2} \sin \frac{A-B}{2} = -\frac{1}{4}, \quad (1)$$

and

$$\sin A - \sin B = -\frac{1}{4} \implies \cos \frac{A+B}{2} \sin \frac{A-B}{2} = -\frac{1}{8}. \quad (2)$$

Now, squaring seems like a good idea:

$$(1)^2 + (2)^2 = \sin^2 \frac{A-B}{2} = \frac{5}{64}.$$

Moreover,

$$(1) \cdot (2) = \frac{1}{2} \sin A + B \sin \frac{A-B}{2} = \frac{1}{32}.$$

Thus,  $100 \sin(A+B) = 100 \cdot \frac{4}{5} = 80$ . □

**Example 1.7** (2018 SMO(S) P17)

Let

$$L = \sum_{k=7}^{16} \left( 1 + \tan(15k^\circ + 15^\circ) \tan 15k^\circ \right).$$

Find  $\lfloor L \rfloor$ .

*Proof.* We see that the expression is dominated by tangents, and that the summand resembles the denominator of the tangent addition formula!

For simplicity, let  $A = 15k^\circ + 15^\circ$  and  $B = 15k^\circ$ , then  $L = \sum_{k=7}^{16} (1 + \tan A \tan B)$ .

To simplify this sum, consider

$$\begin{aligned} L \cdot \tan(A-B) &= L \cdot \tan 15^\circ \\ &= \sum_{k=7}^{16} \tan(A-B) (1 + \tan A \tan B) \\ &= \sum_{k=7}^{16} (\tan A - \tan B) = \sum_{k=7}^{16} \left( \tan(15k^\circ + 15^\circ) - \tan 15k^\circ \right) \\ &= \left( \tan 120^\circ - \tan 105^\circ \right) + \left( \tan 135^\circ - \tan 120^\circ \right) + \cdots + \left( \tan 255^\circ - \tan 240^\circ \right) \\ &= \tan 255^\circ - \tan 105^\circ \\ &= 2 \tan 75^\circ \end{aligned}$$

To this end, we first compute  $\tan 15^\circ$ :

$$\tan 30^\circ = \frac{\sqrt{3}}{3} = \frac{2 \tan 15^\circ}{1 - \tan^2 15^\circ} \implies \tan 15^\circ = 2 - \sqrt{3}.$$

Then,

$$\begin{aligned} L &= \frac{2 \tan 75^\circ}{\tan 15^\circ} = \frac{2 \tan(30+45)^\circ}{2 - \sqrt{3}} \\ &= \frac{4 + 2\sqrt{3}}{2 - \sqrt{3}} \cdot \frac{2 + \sqrt{3}}{2 + \sqrt{3}} \\ &= 14 + 8\sqrt{3}. \end{aligned}$$

Finally, since  $1.7 < \sqrt{3} < 1.75$ ,

$$14 + 8 \cdot 1.7 = 27.6 < L = 14 + 8\sqrt{3} < 14 + 8 \cdot 1.75 = 28,$$

thus  $\lfloor L \rfloor = \boxed{27}$ . □

**Example 1.8** (2021 SMO(S) P15)

Find the minimum value of

$$\frac{8}{\sin 2\theta} + 12 \tan \theta,$$

where  $0 < \theta < \frac{\pi}{2}$ .

*Proof.*

$$\begin{aligned} \frac{8}{\sin 2\theta} + 12 \tan \theta &= \frac{4 + 12 \sin^2 \theta}{\sin \theta \cos \theta} \\ &= \frac{16 \sin^2 \theta + 4 \cos^2 \theta}{\sin \theta \cos \theta} \\ &\geq \frac{2\sqrt{64 \sin^2 \theta \cos^2 \theta}}{\sin \theta \cos \theta} \quad \text{by AM-GM} \\ &= \boxed{16}. \end{aligned}$$

Equality occurs when  $4 \cos^2 \theta = 16 \sin^2 \theta \implies \tan \theta = \frac{1}{2}$ , for which  $\theta$  lies in the given domain. □

## 1.1 Sums and Products (★)

As a disclaimer, some parts of this section may tie into complex numbers, so you may wish to omit these parts until you have had some form of treatment on the connection between complex numbers and trigonometry.

That said, most of this section should be self-contained, and we do our best to not tie into complex numbers, although sometimes they may be used as motivation. This is just a testament to the interconnectedness between these two beautiful ideas!

**Example 1.9** (2014 SMO(S) P33)

Find the value of

$$2 \tan 1^\circ (\sin 2^\circ + \sin 4^\circ + \cdots + \sin 178^\circ).$$

With some foresight, we first note the useful identity that

$$\cos A - 1 + \cos A + 1 = 2 \cos A \cos 1 \tag{3}$$

*Proof.* Let  $S = \sin 2^\circ + \sin 4^\circ + \cdots + \sin 178^\circ$ , then

$$\begin{aligned} S &= 2(\sin 2^\circ + \sin 4^\circ + \cdots + \sin 178^\circ + \sin 180^\circ) \\ &= 2(\sin 2^\circ + \sin 4^\circ + \cdots + \sin 88^\circ) + 1 \\ &= 2(\cos 2^\circ + \cdots) + (\cos 0^\circ + \cos 90^\circ) \\ &= (\cos 0^\circ + \cos 2^\circ) + (\cos 2^\circ + \cos 4^\circ) + \cdots + (\cos 88^\circ + \cos 90^\circ) \\ &= 2 \cos 1^\circ (\cos 1^\circ + \cos 3^\circ + \cdots + \cos 89^\circ) \quad \text{using (3)} \end{aligned}$$

Now, we use a trick to simplify the terms:

$$\begin{aligned} S &= \cot 1^\circ (2 \sin 1^\circ \cos 1^\circ + 2 \sin 3^\circ \cos 1^\circ + \cdots + 2 \sin 89^\circ \cos 1^\circ) \\ &= \cot 1^\circ ((\sin 2^\circ - \sin 0^\circ) + (\sin 4^\circ - \sin 2^\circ) + \cdots + (\sin 90^\circ - \sin 88^\circ)) \\ &= \cot 1^\circ (\sin 90^\circ - \sin 0^\circ) = \cot 1^\circ. \end{aligned}$$

Hence, the required value is  $\boxed{2}$ . □

**Example 1.10** (2013 AIME I/14)

For  $\pi \leq \theta \leq 2\pi$ , let

$$P = \frac{1}{2} \cos \theta - \frac{1}{4} \sin 2\theta - \frac{1}{8} \cos 3\theta + \frac{1}{16} \sin 4\theta + \frac{1}{32} \cos 5\theta - \frac{1}{64} \sin 6\theta - \frac{1}{128} \cos 7\theta + \cdots$$

and

$$Q = 1 - \frac{1}{2} \sin \theta - \frac{1}{4} \cos 2\theta + \frac{1}{8} \sin 3\theta + \frac{1}{16} \cos 4\theta - \frac{1}{32} \sin 5\theta - \frac{1}{64} \cos 6\theta + \frac{1}{128} \sin 7\theta + \cdots$$

so that  $\frac{P}{Q} = \frac{2\sqrt{2}}{7}$ . Find the value of  $\sin \theta$ .

Surely  $P$  and  $Q$  are connected by some obscure relation - this is almost by design! For the confused reader (I sure was!) who can't spot the pattern just yet, the terms in  $P$  and  $Q$  have signs that oscillate in the pattern of  $(+, -, -, +)$ , while the trigonometric terms oscillate between  $\sin$  and  $\cos$ .

The terms in  $P$  and  $Q$  are quite unruly, so let us attempt to line them up. A naive attempt would be to first "unify" the  $\cos \theta$  term from  $P$  and the  $\sin \theta$  term from  $Q$  by considering  $P \cdot \sin \theta$  and  $Q \cdot \cos \theta$ .

Now, for each term in  $P$  and  $Q$  with the same coefficient, note that we have the identities

$$\begin{cases} \sin n\theta \cos \theta + \cos n\theta \sin \theta = \sin(n+1)\theta \\ \cos n\theta \sin \theta - \sin n\theta \cos \theta = \cos(n+1)\theta. \end{cases}$$

Somehow, we now have the miraculous identity:

$$P \cdot \sin \theta - Q \cdot \cos \theta = -2P.$$

Dividing across by  $Q$  yields

$$\frac{P}{Q} = \frac{\cos \theta}{\sin \theta + 2} = \frac{2\sqrt{2}}{7},$$

and squaring and simplifying produces the quadratic

$$57 \sin^2 \theta + 32 \sin \theta - 17 = 0 \implies (19 \sin \theta + 17)(3 \sin \theta - 1) = 0.$$

Since  $\pi \leq \theta \leq 2\pi$ ,  $\sin \theta < 0$  so  $\sin \theta = \boxed{-\frac{17}{19}}$  only.

## 2 Some Geometry

As it turns out, many concepts of a triangle is unified through its *area*.

We start by asserting (without proof) the sine and cosine law. Let  $A, B, C$  be angles in a triangle and let  $a, b, c$  be the lengths opposite their respective angles.

### Proposition 2.1 (Sine Law)

Given a triangle  $ABC$ ,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

### Proposition 2.2 (Cosine Law)

Given a triangle  $ABC$ ,

$$c^2 = a^2 + b^2 - 2ab \cos C$$

. Upon a rearrangement, we also have the useful form

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}.$$

### 2.1 Lengths (★)

Trigonometry is very closely related to the (unit) circle. We have seen the sine rule, but as it turns out, the sine rule can also tie in neatly to the circumcircle. This is the so-called *extended* sine rule:

### Proposition 2.3 (Extended sine rule)

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$$

It suffices to show that  $\frac{a}{\sin A} = 2R$ , and certainly,  $2R$  is the diameter of the circumcircle. It would thus make sense to tie in its diameter.




Figure 2: Two setups of the Sine Rule. <sup>1</sup>



2

*Proof.* Let  $BX$  be a diameter of the circumcircle as above. Then,  $BXC$  is a right triangle with right angle at  $C$ . Thus,  $BC = a$ ,  $BX = 2R$ . Moreover, either  $\angle BXC = \angle BAC$  or  $\angle BXC = \pi - \angle BAC$ .

In any case,  $\sin A = \sin \angle BXC = \frac{a}{2R}$ , and we are done.  $\square$

We have seen setups involving a circle and a triangle, but what about a circle and *two* triangles?

A canonical result is **Ptolemy's theorem**.

**Theorem 2.4** (Ptolemy's Theorem)

Let  $ABCD$  be a cyclic quadrilateral. Then,

$$AB \cdot CD + AD \cdot BC = AC \cdot BD.$$

There are certainly more elegant proofs involving similar triangles, complex numbers, *inversion*, etc. But here, we showcase a trigonometric approach.

We should first anticipate what our mode of attack should look like. While this is a result involving lengths, it is obstructive to work purely in lengths since we don't have a way to tie in to the cyclic nature of the setup.

On the other hand, angles should work nicely because angles uniquely define our quadrilateral.

**Remark 2.5.** This may be counterintuitive, since this is a length result! However this should be expected: we don't really care about each individual length, but rather *how* each length relates to each other.




Figure 3: Looks kind of neat to be honest.

<sup>2</sup>From Euclidean Geometry in Mathematical Olympiads by Evan Chen, pp.52.

With some foreshadowing, we consider the area of our quadrilateral  $ABCD$ , as well as the area of another useful quadrilateral. Ideally, we want our second quadrilateral to have an easily computable area.

*Proof.* Let  $E$  be the point such that  $BE$  is parallel to  $AC$ .

Then,

$$\begin{aligned} [ABCD] &= [ADE] + [DCE] \\ &= \frac{1}{2} (AD \cdot AE \sin \angle DAE + CD \cdot CE \sin \angle DCE) \\ &= \frac{1}{2} (AD \cdot AE + CD \cdot CE) \sin \angle DAE \end{aligned}$$

where the third equality follows since  $\sin \angle DAE = \sin \angle DCE$  and the last equality follows because  $AEBC$  is an isosceles trapezium.

On the other hand, if we let  $X$  be the intersection of the diagonals  $AC$  and  $BD$ , then

$$\begin{aligned} [ABCD] &= \frac{1}{2} \cdot AC \cdot BD \sin \angle BXC \\ &= \frac{1}{2} \cdot AC \cdot BD \sin \angle XBE = \frac{1}{2} \cdot AC \cdot BD \sin \angle DBE \\ &= \frac{1}{2} \cdot AC \cdot BD \sin \angle DCE = \frac{1}{2} \cdot AC \cdot BD \sin \angle DAE. \end{aligned}$$

Thus,

$$\begin{aligned} \frac{1}{2} (AD \cdot AE + CD \cdot CE) \sin \angle DAE &= \frac{1}{2} \cdot AC \cdot BD \sin \angle DAE \\ \implies AB \cdot CD + AD \cdot BC &= AC \cdot BD. \end{aligned}$$

□

An immediate-ish consequence of Ptolemy's theorem is Stewart's theorem, which allows us to find the length of a *cevian*.

**Theorem 2.6** (Stewart's Theorem)

Let  $ABC$  be a triangle and let  $D$  be a point on  $BC$ . Define also  $m = DB, n = DC, d = AD$ . Then


$$man + ad^2 = b^2m + c^2n.$$

This is also written in the more comical form

$$man + dad = bmb + cnc,$$

which fits into a nifty mnemonic "a *man* and his *dad* put a *bomb* in the *sink*."

We present two proofs - the first is a consequence of Ptolemy's theorem.

Figure 4:  $AD$  is a cevian of  $ABC$ .

*Proof.* Let the ray  $AD$  meet the circumcircle of  $ABC$  again at  $P$ . Then,  $BDP \sim ADC$  and  $CDP \sim ADB$ , whence

$$\frac{BP}{b} = \frac{m}{d} \quad \text{and} \quad \frac{CP}{c} = \frac{n}{d}.$$

Moreover, by *power of a point*,

$$DP = \frac{mn}{d}.$$

Finally, by Ptolemy's theorem,

$$\begin{aligned} AP \cdot BC &= AC \cdot BP + AB \cdot CP \\ a \left( d + \frac{mn}{d} \right) &= b \cdot \frac{bm}{d} + c \cdot \frac{cn}{d} \\ ad^2 + man &= b^2m + c^2n. \end{aligned}$$

□

Our second proof goes by way of applying the cosine rule to point  $D$  in two ways.

*Proof.*

$$\frac{n^2 + d^2 - b^2}{2dn} = \cos \angle ADC = -\cos \angle ADB = -\frac{d^2 + m^2 - c^2}{2dm}.$$

This gives

$$n(d^2 + m^2 - c^2) + m(n^2 + d^2 - b^2) = 0,$$

upon which expanding and simplifying gives the result. □

## 2.2 Area Results (★)

With these results, it turns out that we can formulate the area of a triangle in many different ways. For convenience, we denote the area of a triangle  $ABC$  as  $[ABC]$ .

**Theorem 2.7** (Area of a Triangle)

$$\begin{aligned}
[ABC] &= \frac{1}{2}ab \sin C = \frac{1}{2}bc \sin A = \frac{1}{2}ca \sin B \\
&= \sqrt{s(s-a)(s-b)(s-c)} \\
&= rs.
\end{aligned}$$

Here,  $r$  is the **inradius** and  $s = \frac{1}{2}(a+b+c)$  is the **semiperimeter** of  $ABC$ .

The relation  $[ABC] = \sqrt{s(s-a)(s-b)(s-c)}$  is the so-called **Heron's formula**, and allows us to relate the side lengths of a triangle to its semiperimeter and inradius.

To prove this, we consider the **contact triangle** of  $ABC$ . The contact triangle is the triangle formed from the three points of tangency of the incircle.




Figure 5:  $DEF$  is the contact triangle of  $ABC$ .<sup>3</sup>

<sup>4</sup> We state, again without proof, the key trigonometric identity of use here:

**Lemma 2.8**

If  $A, B, C$  are angles in a triangle and are each acute, then

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

The proof of this is sneakily left as a problem in the exercises section.

*Proof.* Now, we have

$$\tan \frac{\pi}{2} - \frac{A}{2} = \tan \angle AIF = \frac{s-a}{r},$$

and similarly,

$$\tan \frac{\pi}{2} - \frac{B}{2} = \tan \angle BID = \frac{s-b}{r}$$

$$\tan \frac{\pi}{2} - \frac{C}{2} = \tan \angle CIE = \frac{s-c}{r}.$$

<sup>4</sup>From Euclidean Geometry in Mathematical Olympiads by Evan Chen, pp.78.

Since we have  $(\frac{\pi}{2} - \frac{A}{2}) + \frac{\pi}{2} - \frac{A}{2} + \frac{\pi}{2} - \frac{A}{2} = \pi$ , our lemma gives, helpfully,

$$\begin{aligned}\frac{s-a}{r} + \frac{s-b}{r} + \frac{s-c}{r} &= \frac{s-a}{r} \cdot \frac{s-b}{r} \cdot \frac{s-c}{r} \\ &= \frac{3s - (a+b+c)}{r} \\ &= \frac{s}{r}.\end{aligned}$$

Finally,  $(sr)^2 = s(s-a)(s-b)(s-c)$ , as needed. □

### 3 Parametrising (★★)

Consider the unit circle  $x^2 + y^2 = 1$ . Then, we may describe the set of points along the circumference of the circle as

$$(x, y) = (\cos \theta, \sin \theta), \quad -\pi \leq \theta \leq \pi.$$

We say that this is a *parametrisation* of the set of points on the circle.

And certainly, the circle  $x^2 + y^2 = r^2$  with radius  $r$  is simply a scaled unit circle, so it's points can be described as

$$(x, y) = (r \cos \theta, r \sin \theta), \quad -\pi \leq \theta \leq \pi.$$

This idea is extremely useful to re-parametrise conditions and to maximise or minimise the so-called objective functions.

**Example 3.1** (2018 SMO(S) P11)

Let  $x$  and  $y$  be real numbers. Find the maximum value of  $2x^2 - 3xy - 2y^2$  subject to the condition that

$$25x^2 - 20xy + 40y^2 = 36.$$

The "cross-term"  $20xy$  suggests that we should complete the square.

*Proof.* Write

$$25x^2 - 20xy + 40y^2 = (25x^2 - 20xy + 4y^2) + 36y^2 = (5x - 2y)^2 + 36y^2 = 36.$$

Now, let  $36a^2 = (5x - 2y)^2$ , then  $36a^2 + 36y^2 = 36 \implies a^2 + y^2 = 1$ , and so we may parametrise

$$(a, y) = (\cos \theta, \sin \theta), \quad -\pi \leq \theta \leq \pi \implies (x, y) = \left( \frac{6 \cos \theta + 2 \sin \theta}{5}, \sin \theta \right).$$

Thus,

$$\begin{aligned} 2x^2 - 3xy - 2y^2 &= \frac{2(6 \cos \theta + 2 \sin \theta)^2 - 15(6 \cos \theta + 2 \sin \theta) \sin \theta - 50 \sin^2 \theta}{25} \\ &= \frac{2(36 \cos^2 \theta + 12 \sin 2\theta + 4 \sin^2 \theta) - (45 \sin 2\theta + 30 \sin^2 \theta) - 50 \sin^2 \theta}{25} \\ &= \frac{72 \cos^2 \theta - 72 \sin^2 \theta - 21 \sin 2\theta}{25} \\ &= \frac{72 \cos 2\theta - 21 \sin 2\theta}{25} \\ &= \frac{\sqrt{72^2 + 21^2} \cos(2\theta + \alpha)}{25}, \quad \text{where } \tan \alpha = \frac{21}{72} \\ &\geq \frac{\sqrt{72^2 + 21^2}}{25} = \boxed{3}. \end{aligned}$$

The minimum occurs when  $2\theta = -\tan^{-1} \frac{21}{72}$ , which lies in the domain of  $\theta$ . □

## 4 Exercises

### 4.1 Warmup

**Problem 4.1.** Let  $ABC$  be a triangle. Prove that

1.  $[ABC] = \frac{abc}{4R}$ , **Hint:** 4
2.  $[ABC] = rs$ . **Hints:** 29 12

**Problem 4.2.** If  $A, B, C$  are angles in a triangle and are each acute, prove that

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

**Hint:** 8

**Problem 4.3** (Tangent Rule). Let  $A, B, C$  be angles in a triangle and let  $a, b, c$  denote the side length of the triangle opposite each respective angle. Prove that

$$\frac{a-b}{a+b} = \frac{\tan \frac{A+B}{2}}{\tan \frac{A-B}{2}}.$$

**Hints:** 23 15

**Problem 4.4** (Catastrophic Cancellation). Let  $A, B, C$  be angles in a triangle and let  $a, b, c$  denote the side length of the triangle opposite each respective angle.

Using the tangent rule, prove that

$$\tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2}.$$

1. In the time before digital calculators, the tangent rule is preferred over the cosine rule when used to find a missing length when given two side lengths  $a, b$  and their enclosed angle  $C$ . Why is this so?
2. For small values of  $C$ , is the tangent rule still preferred?

**Problem 4.5** (Cotangent rule). Let  $A, B, C$  be angles in a triangle and let  $a, b, c$  denote the side length of the triangle opposite each respective angle. Prove that

$$\frac{\cot \frac{A}{2}}{s-a} = \frac{\cot \frac{B}{2}}{s-b} = \frac{\cot \frac{C}{2}}{s-c} = \frac{1}{r}.$$

**Hints:** 3 30

**Problem 4.6** (Mollweide's Formulas). Prove that

$$1. \quad \frac{a+b}{c} = \frac{\cos \frac{A-B}{2}}{\sin \frac{C}{2}}.$$

**Hints:** 28 31

$$2. \quad \frac{a-b}{c} = \frac{\sin \frac{A-B}{2}}{\cos \frac{C}{2}}.$$

**Hints:** 18 21

## 4.2 Problems

**Problem 4.7** (2019 SMO(O) P18). In triangle  $ABC$ ,  $BC = 10$ ,  $CA = 9$ ,  $\cos C = \frac{5}{14}$ . The angle bisector of  $C$  meets the circumcircle again at  $N$ , the altitude from  $A$  meets the circumcircle again at  $L$  and  $NL$  intersects  $BC$  at  $M$ . Find the length of  $MC$ . **Hints:** 19 2 13

**Problem 4.8** (2021 Taiwan APMO Preliminary I). In triangle  $ABC$ ,  $\angle A = 23^\circ$  and  $\angle B = 46^\circ$ . Let  $\Gamma$  be a circle with center  $C$  with radius  $AC$  and let the angle bisector of  $B$  intersect  $\Gamma$  at  $M$  and at  $N$ . Find  $\angle MAN$ . **Hints:** 25 27 26

**Problem 4.9** (2021 SMO(O) P8). Find the minimum value of  $(x+7)^2 + (x+4)^2$  subject to the constraint  $(x-5)^2 + (y-7)^2 = 4$ . **Hint:** 17

**Problem 4.10** (2021 SMO(O) P18). Let  $ABC$  be a triangle with  $AB = 10$  and

$$\frac{\cos A}{\cos B} = \frac{b}{a} = \frac{4}{3}.$$

Let  $P$  be a point on the incircle of  $ABC$ . Find the maximum value of

$$PA^2 + PB^2 + PC^2.$$

**Hints:** 14 9 16 24 6

**Problem 4.11** (2022 AIME I/15). Let  $x, y, z$  be positive reals satisfying the following system of equations:

$$\begin{aligned}\sqrt{2x - xy} + \sqrt{2y - xy} &= 1 \\ \sqrt{2y - yz} + \sqrt{2z - yz} &= \sqrt{2} \\ \sqrt{2z - zx} + \sqrt{2x - zx} &= \sqrt{3}.\end{aligned}$$

Find the value of  $[(1-x)(1-y)(1-z)]^2$ . **Hints:** 1 5

**Problem 4.12** (2019 SMO(O) P17). Two circles are tangent to each other internally at point  $T$ . Let chord  $AB$  of the larger circle be tangent to the smaller circle at  $P$ . If  $AB = \frac{59}{\sqrt{3}}$ ,  $TA = 32$  and  $TB = 27$ , find the length of  $TP$ . **Hints:** 22 20

**Problem 4.13** (2019 SMO(O) P15). In triangle  $ABC$ ,  $AB = 209$ ,  $BC = 171$ ,  $CA = 190$  and  $M$  is the midpoint of  $BC$ . Point  $L$  lies on the extension of  $BA$  and point  $N$  lies on the extension of  $MA$  such that points  $C, L, N$  are collinear and  $AL = NL$ . Find the length of  $AN$ . **Hints:** 11 10 7



## 5 Hints

1. What is the domain of  $x, y, z$ ?
2. Use the angle bisector theorem on  $BLD$ .
3. Find an expression for  $\cot \frac{A}{2}$ .
4. Sine rule.
5. Let  $x = 2 \sin^2 \theta$ . Why is this so?
6. Find  $PA^2 + PB^2 + PC^2$  in terms of sines and cosines and hence maximise this.
7. Find  $NK$ .
8.  $C = \pi - (A + B)$ .
9. Prove that  $ABC$  is a right-angled triangle.
10. Use Apollonius' theorem (google it if you have to!).
11. Construct a point  $K$  such that  $ABKC$  is a parallelogram.
12. Divide  $[ABC]$  into the area of three right triangles.
13. Find  $CD$ .
14. Let  $D$  be the foot of perpendicular from  $C$  onto the angle bisector of  $B$  and let  $C'$  be the foot of perpendicular from  $C$  onto  $BC$ .
15. Use the sum-to-product identities.
16. You may find it convenient to impose the setup on the cartesian plane. Let  $C = (0, 0)$  and define  $A, B$  in a similar way.
17. Parametrise  $x$  and  $y$  in terms of sines and cosines.
18.  $\cos \frac{C}{2} = \sin \frac{A}{2} + \sin \frac{B}{2}$ .
19. Draw  $LB$ .
20. Stewart's Theorem.
21. Convert everything to cotangents.
22. Prove that  $TP$  bisects  $\angle ATB$ . This is called the death star lemma or the shooting lemma (useful result, so google a proof if you have to!).
23. Prove that  $\frac{a-b}{a+b} = \frac{\sin A - \sin B}{\sin A + \sin B}$ .
24. Find the inradius of the incircle and hence parametrise  $P$  in terms of sines and cosines.
25. Let  $D$  be the foot of perpendicular from  $C$  onto the angle bisector of  $B$  and let  $C'$  be the foot of perpendicular from  $C$  onto  $BC$ .
26. Use the law of sines to find  $\frac{BC}{AC}$ .
27. Use the extended sine law to find  $MN$ . You may also note that  $MN = 2MC'$ .
28. Start from  $\frac{\cos \frac{A}{2} - \frac{B}{2}}{\cos \frac{A}{2} + \cos \frac{B}{2}}$ .

29. Draw a good diagram.
30. Consider the triple cotangent identity.
31. Convert everything to cotangents. You may also need to use the second problem.