


5

Eigenvalues and Eigenvectors

5.5

COMPLEX EIGENVALUES



COMPLEX EIGENVALUES

- The matrix eigenvalue-eigenvector theory already developed for \mathbb{R}^n applies equally well to \mathbb{C}^n .
- So a complex scalar λ satisfied $\det(A - \lambda I) = 0$ if and only if there is a nonzero vector x in \mathbb{C}^n such that $Ax = \lambda x$.
- We call λ a **(complex) eigenvalue** and x a **(complex) eigenvector** corresponding to λ .

COMPLEX EIGENVALUES

- **Example 1** If $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then the linear transformation $x \mapsto Ax$ on \mathbb{R}^2 rotates the plane counterclockwise through a quarter-turn.
- The action of A is periodic, since after four quarter-turns, a vector is back where it started.
- Obviously, no nonzero vector is mapped into a multiple of itself, so A has no eigenvectors in \mathbb{R}^2 and hence no real eigenvalues.
- In fact, the characteristic equation of A is

$$\lambda^2 + 1 = 0$$

COMPLEX EIGENVALUES

- The only roots are complex: $\lambda = i$ and $\lambda = -i$. However, if we permit A to act on \mathbb{C}^2 , then

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ -i \end{bmatrix} = \begin{bmatrix} i \\ 1 \end{bmatrix} = i \begin{bmatrix} 1 \\ -i \end{bmatrix}$$
$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ i \end{bmatrix} = \begin{bmatrix} -i \\ 1 \end{bmatrix} = -i \begin{bmatrix} 1 \\ i \end{bmatrix}$$

- Thus i and $-i$ are eigenvalues, with $\begin{bmatrix} 1 \\ -i \end{bmatrix}$ and $\begin{bmatrix} 1 \\ i \end{bmatrix}$ as corresponding eigenvectors.

REAL AND IMAGINARY PARTS OF VECTORS

- The complex conjugate of a complex vector x in \mathbb{C}^n is the vector \bar{x} in \mathbb{C}^n whose entries are the complex conjugates of the entries in x .
- The **real** and **imaginary parts** of a complex vector x are the vectors $\operatorname{Re} x$ and $\operatorname{Im} x$ in \mathbb{R}^n formed from the real and imaginary parts of the entries of x .

REAL AND IMAGINARY PARTS OF VECTORS

■ **Example 4** If $x = \begin{bmatrix} 3 - i \\ i \\ 2 + 5i \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix} + i \begin{bmatrix} -1 \\ 1 \\ 5 \end{bmatrix}$, then

$$\operatorname{Re} x = \begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix}, \quad \operatorname{Im} x = \begin{bmatrix} -1 \\ 1 \\ 5 \end{bmatrix}, \quad \text{and} \quad \bar{x} = \begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix} - i \begin{bmatrix} -1 \\ 1 \\ 5 \end{bmatrix} = \begin{bmatrix} 3 + i \\ -i \\ 2 - 5i \end{bmatrix}$$

EIGENVALUES AND EIGENVECTORS OF A REAL MATRIX THAT ACTS ON

- **Theorem 9:** Let A be a real 2×2 matrix with a complex eigenvalue $\lambda = a - bi$ ($b \neq 0$) and an associated eigenvector v in \mathbb{C}^2 . Then

$$A = PCP^{-1}, \text{ where } P = [\operatorname{Re} v \quad \operatorname{Im} v] \text{ and } C = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$