

HOMOTOPY LIE THEORY

DANIEL ROBERT-NICOUD AND BRUNO VALLETTE

ABSTRACT. To be typed at the very end.

CONTENTS

Introduction	2
1. Recollections [Bruno]	2
2. Cosimplicial object [Bruno]	4
3. Relation with Maurer–Cartan spaces [Daniel]	8
4. Higher Lawrence–Sullivan algebra [Daniel] [Bruno]	8
5. How to fill horns in Maurer–Cartan spaces [Bruno]	9
6. Homotopy theory [Bruno]	11
Appendix A. Formal equations [Bruno]	13
References	14

-
- Convention homologique
 - \mathfrak{L}_∞ -algebras
 - $\mathfrak{L}_\infty\text{-alg}$: the category of complete \mathfrak{L}_∞ -algebras
 - α Maurer–Cartan element
 - R : representation functor
 - C_\bullet : objet simplicial
 - C^\bullet : objet cosimplicial
 - \mathbf{sSet} : category of simplicial sets
 - font sans serif pour les categories
 - \mathbb{K} field (char 0)
 - $C^\bullet := C(\Delta^\bullet)$ cellular chain complex (cosimplicial object)
 - $C_\bullet := C(\Delta^\bullet)^\vee$ (co)cellular chain complex (simplicial object)
 - operad \mathbf{Com}
 - \mathbf{CC}_∞ -algebras for $\Omega\mathbf{BCom}$ -algebras.
 - A^\vee : linear dual
 - $\mathfrak{mc}_\bullet := \Omega_\pi C^\bullet$: the *Maurer–Cartan cosimplicial \mathfrak{L}_∞ -algebra*
 - \mathfrak{Lie} : operad
 - \mathbf{MC}_\bullet : Deligne–Hinich ∞ -groupoid.
 - γ_\bullet : Getzler ∞ -groupoid
 - \mathbf{RT} set of rooted trees
 - $\overline{\mathbf{RT}}$: reduced rooted trees (without |)
 - \mathbb{S} symmetric group
 - \mathcal{T} free operad
 - $\widehat{\mathfrak{L}_\infty(V)}$ free complete \mathfrak{L}_∞ -algebra
 - $\tau, |\tau|$: a rooted tree and its number of vertices
 - d différentielle de la \mathfrak{L}_∞ -algebra

Date: March 10, 2018.

The first author was supported by grants from Région Ile-de-France, the second author was supported by the Institut Universitaire de France, and both authors were supported by the grant ANR-14-CE25-0008-01 project SAT.

- $\text{PT}^{[n]}$: set of planar rooted trees with internal vertices having at least one input and such that the total number of vertices and leaves is equal to n .
- Λ_k^n horns [toujours pas fan des deux petites barres en bas ... Pas trouve mieux](#)
- Δ^n ensemble simpliciaux "principaux"
- \mathbb{N} : integers
- \mathbb{R} : real numbers
- Δ simplex category

INTRODUCTION

To be typed at the very end.

Layout.

Conventions.

convention homologique : que des complexes de chaines (differentielles de degrees -1); les complexes de "cochaines" sont concentres en degrees de $-n$ a 0

corps de base

(degree-wise, arity-wise) linear dual A^\vee .

We work over a field \mathbb{K} of characteristic 0 .

1. RECOLLECTIONS [\[BRUNO\]](#)

The classical Lie theory is made of an equivalence between a category of Lie algebras and a category of groups. The higher analogue is actually a Quillen equivalence between a category of homotopy Lie algebras and a category of ∞ -groupoid, notions that we recall and make precise in this section. [a reprendre a la fin.](#)

1.1. Complete shifted homotopy Lie algebras. In this section, we recall the main features of complete shifted homotopy Lie algebras.

Definition 1.1 (Complete dg module). *A complete differential graded (dg) module (A, d, F) is a chain complex, i.e. with homological degree convention $|d| = -1$, equipped with a filtration*

$$A = F_0 A = F_1 A \supset F_2 A \supset \cdots \supset F_k A \supset F_{k+1} A \supset \cdots$$

made up of dg sub-modules, such that the associated topology is complete.

Notice that in the present paper, we restrict ourselves to the case $F_0 A = F_1 A$. A morphism of complete dg module is a chain map preserves the respective filtrations. Any dg module V is viewed as a complete dg module endowed with the discrete thus complete filtration $V = F_0 V = F_1 V \supset F_2 V = F_3 V = \cdots = 0$.

Definition 1.2 (Complete shifted \mathcal{L}_∞ -algebra). *A complete shifted \mathcal{L}_∞ -algebra structure, on a complete differential graded module (A, d, F) is a collection $(\ell_2, \ell_3, \dots) \in \prod_{n \geq 2} \text{hom}(A^{\odot n}, A)$ of symmetric maps which respect the filtration, of degree -1 , and satisfying*

$$\partial(\ell_n) + \sum_{\substack{p+q=n+1 \\ 2 \leq p, q \leq n}} \sum_{\sigma \in \text{Sh}_{p,q}^{-1}} (\ell_{p+1} \circ_1 \ell_q)^\sigma = 0,$$

where $\text{Sh}_{p,q}^{-1}$ denotes the set of the inverses of (p, q) -shuffles.

These are actually algebras over the operad $\mathfrak{s}\mathcal{L}_\infty := \mathcal{S} \otimes \mathcal{L}_\infty \cong \Omega \text{Com}^\vee$, which is the suspension of the operad encoding \mathcal{L}_∞ -algebra. So we will call them *complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras*. In the present paper, we consider the category of algebras over this operad, denoted $\mathfrak{s}\mathcal{L}_\infty\text{-alg}$, that is with strict morphisms of $\mathfrak{s}\mathcal{L}_\infty$ -algebras, which are made up of one map of degree 0 commuting strictly with the structure operations. This operadic interpretation over the symmetric monoidal category of complete dg modules with the completed tensor product $\widehat{\otimes}$ [\[DSV18\]](#) allows us to apply to complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras all the classical operadic constructions and results [\[LV12\]](#), like the following one.

Proposition 1.3. *The category of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras is locally small, complete, and cocomplete.*

Proof. All the arguments of [DSV18, Lemma 4] hold true for complete dg modules whose filtration satisfies $F_0 = F_1$. Therefore this latter category is locally small, complete, and cocomplete and thus is the category of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras as a category of algebras over an operad, by [DSV18, Theorem 1]. \square

Remark 1.4. *By definition of the suspension operad $\mathcal{S} := \text{End}_{\mathbb{K}, \mathcal{S}}$, the category of (complete) $\mathfrak{s}\mathcal{L}_\infty$ -algebras is isomorphic to the category of \mathcal{L}_∞ -algebras, the two isomorphism functors are given by the (de)suspension of the underlying chain complexes: $\mathfrak{g} \mapsto s^{\pm 1}\mathfrak{g}$. In the present context, $\mathfrak{s}\mathcal{L}_\infty$ -algebras appear more naturally than their analogues and they have the great advantage of carrying much less signs (if not none).*

Definition 1.5 (Maurer–Cartan element). *A Maurer–Cartan element is a degree 0 element $\alpha \in \mathfrak{g}_0$ of a complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras satisfying the Maurer–Cartan equation:*

$$d(\alpha) + \sum_{n \geq 2} \frac{1}{n!} \ell_n(\alpha, \dots, \alpha) = 0 ;$$


the associated set is denoted by $\text{MC}(\mathfrak{g})$.

For any $n \geq 2$, we consider the set RT_n of rooted trees with n leaves such that each vertex has at least two inputs. The set $\text{RT}_1 := \{|\}$ is made up of the trivial rooted tree. We denote the number of vertices of a rooted tree $\tau \in \text{RT}$ by $|\tau|$. The set of *reduced* rooted trees, that is without the trivial tree, is denoted by $\overline{\text{RT}}$.

Proposition 1.6. *Let V be a dg module with basis $\{v_i\}_{i \in I}$. The free complete $\mathfrak{s}\mathcal{L}_\infty$ -algebra generated by V is isomorphic to the following product*

$$\widehat{\mathfrak{s}\mathcal{L}_\infty(V)} := \mathfrak{s}\mathcal{L}_\infty \widehat{\circ} V \cong \prod_{n \geq 1} \mathfrak{s}\mathcal{L}_\infty(n) \otimes_{\mathbb{S}_n} V^{\otimes n} ,$$

with a basis given series indexed by n of linear combinations of rooted trees τ , of degree $-|\tau|$, with n leaves labelled by elements v_i .



The differential is the sum of the internal differential on V with the splitting of all corollas into two. The operations ℓ_n amounts to graft n rooted trees to n leaves of a corolla.

Proof. The set RT of rooted trees provides us with a basis of the operad $\mathfrak{s}\mathcal{L}_\infty \cong \Omega\text{Com}^\vee \cong \mathcal{T}(s^{-1}\overline{\text{Com}})$. The form of the complete free algebra on a discrete dg module follows from [DSV18, Section 2.2]. The differential and the structure operations are produced by the general theory of operads, see [LV12, Section 6.5.2]. \square

1.2. Algebraic ∞ -groupoid.

Definition 1.7 (Simplex category Δ). *The simplex category, denoted by Δ , has, for objects, the totally ordered sets $[n] = \{0 < \dots < n\}$, for $n \in \mathbb{N}$, and, for morphisms, the order-preserving maps.*

Definition 1.8 (Simplicial set). *The category of simplicial sets, denoted by sSet , is the category of presheaves $\text{sSet} := \text{Fun}(\Delta^{\text{op}}, \text{Set})$ over the simplex category.*

Presheaves over the opposite category are called *cosimplicial sets*. To avoid confusion, when it is necessary, we denote simplicial sets by X_\bullet and respectively cosimplicial sets by X^\bullet .

Example 1.9. For $n \in \mathbb{N}$, the standard n -simplex is the simplicial set $\Delta^n := \text{Hom}_\Delta(-, [n])$ represented by $[n]$. This combinatorial object encodes the cellular decomposition of the geometric n -simplex, which is the convex hull of the unit vectors in \mathbb{R}^{n+1} :

$$|\Delta^n| := \{(t_0, \dots, t_n) \in [0, 1]^{n+1} \mid t_0 + \dots + t_n = 1\} .$$

Definition 1.10 (Horn). The k^{th} horn Λ_k^n of dimension n , for $n \geq 2$ and $0 \leq k \leq n$, is ...

We call an n -horn in a simplicial set X_\bullet a morphism of simplicial sets $\Lambda_k^n \rightarrow X_\bullet$ and a *horn filler* a morphism of simplicial sets $\Delta^n \rightarrow X_\bullet$ which lifts the horn:

$$\begin{array}{ccc} \Lambda_k^n & \longrightarrow & X_\bullet \\ \downarrow & \nearrow & \\ \Delta^n & & \end{array} .$$

Definition 1.11 (Kan complex). A Kan complex is simplicial set X_\bullet for which every horn $\Lambda_k^n \rightarrow X_\bullet$, for $n \geq 2$ and $0 \leq k \leq n$, can be filled.

Proposition 1.12 (see [GJ09, Lemma 3.3]). The singular simplicial set $\text{Hom}_{\text{Top}}(|\Delta^\bullet|, X)$ associated to any topological space X is a Kan complex.

Moreover, the singular simplicial set functor is faithful (but not full) and Kan complexes are known to share the same homotopy theory than compactly generated Hausdorff spaces notably after the seminal works of D. Quillen [Qui67]. That is why, many people refer to Kan complexes simply as to “spaces”.

The notion of a Kan complex is a property, one can get make it into a structure by the following definition due to T. Nikolaus in [Nik11], called algebraic Kan complex in *loc. cit.*.

Definition 1.13 (Algebraic ∞ -groupoid). An algebraic ∞ -groupoid is a simplicial set X_\bullet with a given filler for which every horn $\Lambda_k^n \rightarrow X_\bullet$, for $n \geq 2$ and $0 \leq k \leq n$.

Proposition 1.14. The nerve of a groupoid but here we have no choice: there exists one and only one horn filler each time.

A reprendre ! modeles pour les ∞ -groupoid, hypothese homotopique, notion de complexe de Kan algebrique de Nikolaus ("mieux que Getzer"). Terminologie : "Algebraic ∞ -groupoid" (ou "algebraic Kan complex") car structure algebrique, pas propriete?

We refer the reader to [GJ09] for more details on simplicial sets and their homotopy properties.

2. COSIMPLICIAL OBJECT [BRUNO]

2.1. Dold–Kan correspondence. Let us start with the following observation, which goes back to D.M. Kan.

Lemma 2.1 ([Kan58]). Let \mathcal{C} be a locally small cocomplete category. The data of a pair of adjoint functors whose left adjoint has domain in simplicial sets

$$L : \text{sSet} \xrightarrow{\perp} \mathcal{C} : R$$

is equivalent to the data of a cosimplicial objects $C^\bullet : \Delta \rightarrow \mathcal{C}$ in \mathcal{C} , under the restriction

$$C^\bullet = LY : \Delta \xrightarrow{Y} \text{sSet} \xrightarrow{L} \mathcal{C}$$

of the left adjoint functor to the Yoneda embedding Y .

Proof. Any cosimplicial object $C^\bullet : \Delta \rightarrow \mathcal{C}$ induces a functor

$$\begin{array}{ccc} R & : & \mathcal{C} \rightarrow \text{sSet} \\ c & \mapsto & \text{Hom}_{\mathcal{C}}(C^\bullet, c), \end{array}$$

which admits a left adjoint given by the following left Kan extension:

$$\begin{array}{ccc} \Delta & \xrightarrow{C^\bullet} & \mathcal{C} \\ & \searrow Y & \nearrow L_{\text{an}_Y} C^\bullet \\ & \text{sSet} & \end{array} .$$

Finally, notice that left adjoint functors preserve colimits and that there is a unique colimit preserving functor $L : \text{sSet} \rightarrow \mathcal{C}$ with prescribed restriction to Δ , by the density theorem: any simplicial set X_\bullet is isomorphic to the colimit

$$X_\bullet \cong \text{colim}_{E(X_\bullet)} Y \circ \Pi .$$

over its category of elements, where $\Pi : E(X_\bullet) \rightarrow \Delta$ is the canonical projection which sends $x \in X_n$ onto $[n]$. \square

This fact shows that if one wants to introduce a functor from a category \mathcal{C} to simplicial sets, a good way is to consider a cosimplicial objects in \mathcal{C} .

Example 2.2. Let $\mathcal{C} = \text{dgMod}$ be the category of dg modules. We consider the cosimplicial dg module

$$C^\bullet := C(\Delta^\bullet)$$

made up of the cellular chain complexes of the geometric n -simplices $|\Delta^n|$ or equivalently the normalised chain complex of the standard n -simplices Δ^n . The associated adjunction

$$L_{\text{DK}} : \text{sSet} \xrightleftharpoons[\perp]{} \text{dgMod} : R_{\text{DK}}$$

is nothing but the one involved in the Dold–Kan correspondence: equivalence of categories between abelian simplicial abelian groups and non-negatively graded chain complexes of abelian groups.

mettre aussi l'exemple du foncteur singulier et de la realisation geometrique ? Coupe le flot mais on s'en sert (un peu) dans la partie categorie de modeles

2.2. Maurer–Cartan cosimplicial complete \mathcal{L}_∞ -algebra. In the present paper, we work with the category $\mathcal{C} = \mathcal{L}_\infty\text{-alg}$ of complete \mathcal{L}_∞ -algebras. To define in this category a suitable cosimplicial object, we consider the following elements involved in Dupont's simplicial version of the de Rham Theorem.

Proposition 2.3 ([Dup76]). *There exists a simplicial contraction*

$$h_\bullet \hookrightarrow \Omega_\bullet \xrightleftharpoons[i_\bullet]{p_\bullet} C_\bullet$$

between the simplicial dg commutative algebras of piecewise polynomial differential forms on the geometric n -simplices

$$\Omega_n := \mathbb{K}[t_0, \dots, t_n, dt_0, \dots, dt_n] / (t_0 + \dots + t_n - 1, dt_0 + \dots + dt_n)$$

and the simplicial dg module made up of the linear dual of the cellular chain complexes of the geometric n -simplices

$$C_\bullet := C(\Delta^\bullet)^\vee .$$

Remark 2.4. Since we work with the homological degree convention, the elements dt_i have all degree -1 and the dg module $C(\Delta^n)^\vee$ is concentrated in degree range $-n$ to 0 with differential of degree -1 .

The former chain complex carries an algebra structure over the operad Com , which admits a canonical cofibrant replacement by means of the bar-cobar construction $\Omega\text{BCom} \xrightarrow{\sim} \text{Com}$. By pulling back, each Ω_n carries an ΩBCom -algebra structure that we can transfer onto $C(\Delta^n)^\vee$ under the homotopy transfer theorem, see [LV12, Section 10.3] for instance. Since the Dupont contraction is simplicial, the various maps can be extended to ∞ -morphisms in order to form a simplicial ΩBCom -algebra structure on $C(\Delta^\bullet)^\vee$. Since the $C(\Delta^n)^\vee$ are degree-wise finite dimensional and since ΩBCom is degree-wise and arity-wise finite dimensional, the cellular chain complexes $C^\bullet = C(\Delta^\bullet)$ admit a cosimplicial $\text{B}\Omega\text{Com}^\vee \cong (\Omega\text{BCom})^\vee$ -coalgebra structure. The canonical twisting morphism

$\pi: \mathbf{B}\Omega\mathbf{Com}^\vee \rightarrow \Omega\mathbf{Com}^\vee \cong \mathcal{SL}_\infty$ induces a bar-cobar adjunction between complete \mathcal{SL}_∞ -algebras and $\mathbf{B}\Omega\mathbf{Com}^\vee$ -coalgebras, see [LV12, Section 11.2]:

$$\Omega_\pi : \mathbf{B}\Omega\mathbf{Com}^\vee\text{-}\mathbf{coalg} \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} \mathbf{d}\mathcal{L}_\infty\text{-}\mathbf{alg} : \mathbf{B}_\pi.$$

Proposition 2.5. *All together, the $\Omega_\pi \mathbf{C}^\bullet$ form a cosimplicial complete $\mathfrak{s}\mathcal{L}_\infty$ -algebra.*

Proof. It remains to notice that the cobar functor Ω_π sends ∞ -morphisms of $\mathbf{B}\Omega\mathbf{Com}^\vee$ -coalgebras to strict morphisms of $\mathfrak{s}\mathcal{L}_\infty$ -algebras, see [RNW17, Section 3]. \square

Definition 2.6 (The Maurer–Cartan cosimplicial $\mathfrak{s}\mathcal{L}_\infty$ -algebra). *The Maurer–Cartan cosimplicial $\mathfrak{s}\mathcal{L}_\infty$ -algebra is defined by*

$$\mathrm{mc}_\bullet := \Omega_\pi \mathbb{C}^\bullet.$$

Let us now try to make this seminal object more explicit. The proof of the following fact will give more details.

Proposition 2.7. *The 0-simplex \mathfrak{mc}_0 is the free complete \mathfrak{L}_∞ -algebra on one Maurer–Cartan element:*

$$\mathrm{mc}_0 \cong (\widehat{\mathfrak{L}_\infty(a_0)}, \mathrm{d}(a_0) = - \sum_{m \geq 2} \frac{1}{m!} \ell_m(a_0, \dots, a_0)) .$$

Proof. The chain complex $C(\Delta^n)^\vee$ can be seen as a sub-chain complex of Ω_n spanned by the basis made up of the differential forms ω_I , for $I = \{i_0, \dots, i_k\} \subset [n]$, with degree $|\omega_I| = -|I| = -k - 1$:

$$\omega_{i_0 \dots i_k} := k! \sum_{j=0}^k (-1)^j t_{i_j} dt_{i_0} \cdots \widehat{dt_{i_j}} \cdots dt_{i_k} .$$

For $n = 0$, the dg commutative algebra $\Omega_0 = \mathbb{K}\omega_0 = \mathbb{K}1$ is trivial and the chain complex $C(\Delta^n)^\vee = \mathbb{K}\omega_0$ is one-dimensional.

Since the dg operad ΩBCom is quasi-free on $s^{-1}\text{BCom} = s^{-1}\mathcal{T}^c(s\overline{\text{Com}})$, the transferred ΩBCom -algebra structure on $C(\Delta^n)^\vee$ amounts to a collection of operations $\{\mu_\tau\}_{\tau \in \text{RT}}$ indexed by rooted trees. The operation $\mu_1 = \text{id}$ is the identity. Given a rooted tree $\tau \in \text{RT}_m$, the associated operation $\mu_\tau : (C(\Delta^n)^\vee)^{\otimes m} \rightarrow C(\Delta^n)^\vee$ has degree $|\tau| - 1$ and is explicitly given by the formula

$$\mu_\tau(\omega_{I_1}, \dots, \omega_{I_6}) =$$

where the corollas means the (iterated) commutative product in Ω_n

For $n = 0$, the contracting homotopy $h_0 = 0$ vanishes and only the operations indexed by corollas c_m do not collapse: $\mu_{c_m}(\omega_0, \dots, \omega_0) = \omega_0$.

Let us denote by $a_I := \omega_I^\vee$ the dual basis on $C(\Delta^n)$. The linear dual of the operad algebra structure corresponding to $s^{-1}\mathrm{BCom}(C(\Delta^n)^\vee) \rightarrow C(\Delta^n)^\vee$ produces the cooperad algebra structure corresponding to $C(\Delta^n) \rightarrow s\Omega\mathrm{Com}^\vee(C(\Delta^n))$. Notice that, since this linear dual includes the isomorphism between invariants and coinvariants under the actions of the symmetric groups \mathbb{S}_m , it carries a coefficient $\frac{1}{m!}$. So, if the element $\lambda\omega_J$, with $\lambda \in \mathbb{K}$, appears in a product $\mu_\tau(\omega_{I_1}, \dots, \omega_{I_m})$, then, dually, the image under the coproduct map of the element a_J includes the term $\frac{1}{\lambda m!} \sigma\tau(a_{I_1}, \dots, a_{I_m})$.


For $n = 0$, the image under the coproduct map of a_0 is equal to

$$a_0 \mapsto \sum_{m \geq 1} \frac{1}{m!} sc_m(a_0, \dots, a_0) .$$

The cobar construction $\Omega_\pi C(\Delta^n)$ admits the free complete \mathfrak{sL}_∞ -algebra on $C(\Delta^n)$

$$\widehat{\mathfrak{L}_\infty}(C(\Delta^n)) \cong \widehat{\mathfrak{L}_\infty}(a_I, I \subset [n])$$

as underlying module; its differential is produced by the difference between the internal differential of the chain complex $C(\Delta^n)$ and the desuspension of the above coalgebra structure, without the primitive term, that is:

$$d(a_J) = \sum_{l=0}^k (-1)^l a_{j_0 \dots \widehat{j_l} \dots j_k} - \sum_{\tau \in \overline{\text{RT}}} \frac{1}{\lambda.6!} \text{ (diagram) }, \text{ for } J = \{j_0, \dots, j_k\}.$$


For $n = 0$, this

$$d(a_0) = - \sum_{m \geq 2} \frac{1}{m!} \ell_m(a_0, \dots, a_0),$$

which concludes the proof. \square

2.3. Non-abelian Dold–Kan adjunction. The Maurer–Cartan cosimplicial \mathfrak{sL}_∞ -algebra mc_\bullet induces canonically the following pair of adjoint functors.

Definition 2.8 (Simplicial representation). *The simplicial representation of complete \mathfrak{sL}_∞ -algebras is produced by the functor*

$$\begin{aligned} \mathbf{R} &: \mathfrak{sL}_\infty\text{-alg} \longrightarrow \mathbf{sSet} \\ \mathfrak{g} &\longmapsto \text{Hom}_{\mathfrak{sL}_\infty\text{-alg}}(\text{mc}_\bullet, \mathfrak{g}). \end{aligned}$$

As a direct corollary of Proposition 2.7, we have

$$\mathbf{R}(\mathfrak{g})_0 = \text{Hom}_{\mathfrak{sL}_\infty\text{-alg}}(\text{mc}_0, \mathfrak{g}) \cong \text{MC}(\mathfrak{g}).$$

Definition 2.9 (Realisation functor). *The \mathfrak{sL}_∞ -realisation functor is defined by*

$$\mathfrak{L}_\infty := \text{Lan}_{\mathbf{Y}} \text{mc}_\bullet : \mathbf{sSet} \rightarrow \mathfrak{sL}_\infty\text{-alg}.$$

Theorem 2.10. *The simplicial representation functor is right adjoint to the \mathfrak{sL}_∞ -realisation functor:*

$$\mathfrak{L}_\infty : \mathbf{sSet} \xrightleftharpoons[\perp]{} \mathfrak{sL}_\infty\text{-alg} : \mathbf{R}.$$

Proof. This is a direct corollary of Lemma 2.1 and Proposition 2.5. \square

The sub-category of (complete) abelian \mathfrak{sL}_∞ -algebras, where all the operations ℓ_n , i.e. for $n \geq 2$, are trivial, is nothing but the category of (complete) dg modules.

Proposition 2.11. *The two simplicial sets which are obtained from a complete dg module under the Dold–Kan representation functor and the \mathfrak{sL}_∞ -representation functor are isomorphic.*

Proof. Let \mathfrak{g} be a complete dg module, viewed as a complete abelian \mathfrak{sL}_∞ -algebra. Its image under the \mathfrak{sL}_∞ -representation functor is equal to

$$\mathbf{R}(\mathfrak{g}) = \text{Hom}_{\mathfrak{sL}_\infty\text{-alg}}(\text{mc}_\bullet, \mathfrak{g}) = \text{Hom}_{\mathfrak{sL}_\infty\text{-alg}}(\widehat{\mathfrak{sL}_\infty}(\mathbf{C}^\bullet), \mathfrak{g}) \cong \text{Hom}_{\text{dgMod}}(\mathbf{C}^\bullet, \mathfrak{g}) = \mathbf{R}_{\text{DK}}(\mathfrak{g}).$$

\square

The sub-category of (complete) \mathfrak{sL}_∞ -algebras where all the operations ℓ_n are trivial, for $n \geq 3$, is nothing but the category of (complete) shifted dg Lie algebras. This latter category is controlled by the operad $\mathfrak{sLie} := \mathcal{S} \otimes \text{Lie}$. In the same way as above, one can consider the cofibrant Koszul resolution $\Omega\text{Com}^i \xrightarrow{\sim} \text{Com}$, and then endow $C(\Delta^\bullet)^\vee$ with a simplicial ΩCom^i -algebra structures by the homotopy transfer theorem. Its linear dual $C(\Delta^\bullet)$ thus admits a cosimplicial $\text{B}\mathfrak{sLie} \cong (\Omega\text{Com}^i)^\vee$ -coalgebra structure. Finally, the bar-cobar adjunction associated to the canonical twisting morphism $\bar{\pi} : \text{B}\mathfrak{sLie} \rightarrow \mathfrak{sLie}$,

$$\Omega_{\bar{\pi}} : \text{B}\mathfrak{sLie}\text{-coalg} \xrightleftharpoons[\perp]{} \mathfrak{sLie}\text{-alg} : \text{B}_{\bar{\pi}},$$

produces the following cosimplicial complete dg Lie algebra:

$$\overline{\mathfrak{mc}}_\bullet := \Omega_{\overline{\pi}} C^\bullet.$$

The associated pair of adjoint functors

$$\mathfrak{Q} : \mathfrak{sSet} \xrightleftharpoons[\perp]{} \mathfrak{sLie}\text{-alg} : \overline{R}$$

is the one introduced in [BFMT16], but in another way. To see this, one just has to prove that XXX.

Proposition 2.12. *The two simplicial sets which are obtained from a complete dg Lie algebra respectively under the \mathfrak{sLie} and the \mathfrak{sL}_∞ -representation functors are isomorphic.*

Proof. to be finished

□

To sum up, the present \mathfrak{sL}_∞ -representation functor extends both the Dold–Kan functor and the \mathfrak{sLie} -representation functor of Buijs–Felix–Murillo–Tanré.

$$\begin{array}{ccc} & \mathfrak{sL}_\infty\text{-alg} & \\ & \uparrow & \\ \mathfrak{sSet} & \xleftarrow{\quad} \mathfrak{sLie}\text{-alg} & \\ & \uparrow & \\ & \mathfrak{dgMod} & \end{array}$$

3. RELATION WITH MAURER–CARTAN SPACES [DANIEL]

3.1. Statement. The literature has already seen several another ways to define Maurer–Cartan spaces, that is actually Kan complexes, out of a complete \mathfrak{sL}_∞ -algebra \mathfrak{g} . One can first consider the *Deligne–Hinich space* [Hin97] made up of the solutions to the Maurer–Cartan equation of the complete \mathfrak{sL}_∞ -algebra structure on the tensor product with the simplicial dg commutative algebra Ω_\bullet :

$$\mathrm{MC}_\bullet(\mathfrak{g}) := \mathrm{MC}(\mathfrak{g} \widehat{\otimes} \Omega_\bullet).$$

On can then use the Dupont simplicial contraction to transfer this \mathfrak{sL}_∞ -algebra structure onto $\mathfrak{g} \widehat{\otimes} C_\bullet$ and thus consider $\mathrm{MC}(\mathfrak{g} \widehat{\otimes} C_\bullet)$ references ? RN et Bandiera?. Finally, one can also consider the *Getzler space* [Get09] made up of the kernels of the Dupont’s homotopy:

$$\gamma_\bullet(\mathfrak{g}) := \{\alpha \in \mathrm{MC}_\bullet(\mathfrak{g}) \mid (\mathrm{id}_{\mathfrak{g}} \widehat{\otimes} h_\bullet)(\alpha) = 0\}.$$

Theorem 3.1. *Let \mathfrak{g} be a complete \mathfrak{sL}_∞ -algebra. The various Maurer–Cartan spaces associated to \mathfrak{g} are related by the following weak equivalences and isomorphisms:*

$$\mathrm{MC}_\bullet(\mathfrak{g}) \simeq \gamma_\bullet(\mathfrak{g}) \cong \mathrm{MC}(\mathfrak{g} \widehat{\otimes} C_\bullet) \cong \mathrm{R}(\mathfrak{g}).$$

E. Getzler first proved the weak equivalence in [Get09]. Then, R. Bandiera established the first isomorphism in his Ph.D. thesis [Ban14]. Finally, the first named author proved directly the weak equivalence $\mathrm{MC}_\bullet(\mathfrak{g}) \simeq \mathrm{MC}(\mathfrak{g} \widehat{\otimes} C_\bullet)$ in [RN17]. What remains to show here is the isomorphism with the present representation functor $\mathrm{R}(\mathfrak{g})$.

3.2. XXX.

3.3. YYY.

3.4. ZZZ.

4. HIGHER LAWRENCE–SULLIVAN ALGEBRA [DANIEL] [BRUNO]

4.1. Generalisation of the Lawrence–Sullivan algebra to the \mathfrak{sL}_∞ -case [Daniel]. ->Daniel : explain the method, different from Proposition 2.7. Describe \mathfrak{mc}_0 and \mathfrak{mc}_1 . Recover Lawrence–Sullivan Lie algebra from that.

4.2. **Homotopies between ∞ -morphisms [Bruno].** I will do it later, when you will have typed the section above

5. HOW TO FILL HORNS IN MAURER–CARTAN SPACES[Bruno]

The image of a topological space under the singular functor produces a Kan complex, that is a simplicial set whose horns can be filled. The issue is actually how to fill these horns. [A reprendre](#)

5.1. **Filling horns [Bruno].** The following result is the key ingredient of this section.

Lemma 5.1. *For $n \geq 2$, there is an isomorphism of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras*

$$\mathrm{mc}_n = \mathfrak{L}_\infty(\Delta^n) \cong \mathfrak{L}_\infty(\Lambda_k^n) \hat{\sqcup} \mathfrak{s}\mathcal{L}_\infty(u, du),$$

where u is a generator of degree n and where the differential on the right-hand side is given by $d(u) = du$.

Proof. We consider the following morphism of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras:

$$\begin{array}{ccc} \mathfrak{L}_\infty(\Lambda_k^n) \hat{\sqcup} \mathfrak{s}\mathcal{L}_\infty(u, du) & \xrightarrow{\varphi} & \mathfrak{L}_\infty(\Delta^n) \\ a_I & \mapsto & a_I \\ u & \mapsto & a_{0\dots n} \\ du & \mapsto & d(a_{0\dots n}) . \end{array}$$

In the other way round, we want to define a morphism of $\mathfrak{s}\mathcal{L}_\infty$ -algebras as follows

$$\begin{array}{ccc} \mathfrak{L}_\infty(\Delta^n) & \xrightarrow{\psi} & \mathfrak{L}_\infty(\Lambda_k^n) \hat{\sqcup} \mathfrak{s}\mathcal{L}_\infty(u, du) \\ a_I & \mapsto & a_I \\ a_{0\dots n} & \mapsto & u \\ a_{0\dots \widehat{k} \dots n} & \mapsto & x . \end{array}$$

For $\psi \circ \varphi = \mathrm{id}$ to be equal to the identity, we need X to satisfy the equation

$$(-1)^k x + \sum_{l \neq k} (-1)^l a_{0\dots \widehat{l} \dots n} - \sum_{\substack{\tau \in \overline{\mathrm{RT}} \\ I_1, \dots, I_m \subset [n]}} \lambda_{\tau, I_1, \dots, I_m} \tau(a_{I_1}, \dots, x, \dots, x, \dots, a_{I_m}) = du ,$$

where

$$d(a_{0\dots n}) = \sum_l (-1)^l a_{0\dots \widehat{l} \dots n} - \sum_{\substack{\tau \in \overline{\mathrm{RT}} \\ I_1, \dots, I_m \subset [n]}} \lambda_{\tau, I_1, \dots, I_m} \tau(a_{I_1}, \dots, a_{0\dots \widehat{k} \dots n}, \dots, a_{0\dots \widehat{k} \dots n}, \dots, a_{I_m}) .$$

This equation is equivalent to the following fixed-point equation:

$$(1) \quad x = du - (-1)^k \sum_{l \neq k} (-1)^l a_{0\dots \widehat{l} \dots n} + (-1)^k \sum_{\substack{\tau \in \overline{\mathrm{RT}} \\ I_1, \dots, I_m \subset [n]}} \lambda_{\tau, I_1, \dots, I_m} \tau(a_{I_1}, \dots, x, \dots, x, \dots, a_{I_m})$$

in the complete $\mathfrak{s}\mathcal{L}_\infty$ -algebra $\mathfrak{L}_\infty(\Lambda_k^n) \hat{\sqcup} \mathfrak{s}\mathcal{L}_\infty(u, du)$. This equation is a particular case of the fixed-point equation (2): for each $N \geq 1$, we consider the map

$$P_N(-, \dots, -) := (-1)^k \sum_{\substack{\tau \in \overline{\mathrm{RT}} \\ I_1, \dots, I_m \subset [n]}} \lambda_{\tau, I_1, \dots, I_m} \tau(a_{I_1}, \dots, -, \dots, -, \dots, a_{I_m}) ,$$

which is well-defined since for any x_1, \dots, x_N , the element $\tau(a_{I_1}, \dots, x_1, \dots, x_N, \dots, a_{I_m})$ lives in F_m and since, at for any $m \geq 2$, the number of rooted trees in RT_m is finite. It remains to prove that each P_N raises the degree filtration by 1, which amounts to show that every non-trivial term appearing in the sum involves at least one a_I , with $I \neq \{0, \dots, \widehat{k}, \dots, n\}$. According to the description given in the proof of Proposition 2.7, this is equivalent to prove that for any reduced rooted tree $\tau \in \overline{\mathrm{RT}}$, we have $\mu_\tau(\omega_{0\dots \widehat{k} \dots n}, \dots, \omega_{0\dots \widehat{k} \dots n}) \neq \lambda \omega_{0\dots n}$, with $\lambda \in \mathbb{K} \setminus \{0\}$, in the homotopy transferred $\Omega\mathrm{BCom}$ -algebra structure on $C(\Delta^n)^\vee$. For $n \geq 3$, we have $\omega_{0\dots \widehat{k} \dots n} \cdot \omega_{0\dots \widehat{k} \dots n} = 0$, so since every vertex of each reduced rooted trees have at least two inputs, we have actually $\mu_\tau(\omega_{0\dots \widehat{k} \dots n}, \dots, \omega_{0\dots \widehat{k} \dots n}) = 0$. [Bruno: do the case \$n = 2\$.](#)

So one can apply Proposition A.4 to get the existence of a solution x to the fixed-point equation (1). Forgetting the differentials, the map ψ is now a well-defined morphism of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebra,

which satisfies $\psi \circ \varphi = \text{id}$. The image under the morphism φ of Equation (1) produces the same kind of equation

$$\varphi(x) = d(a_{0\dots n}) - (-1)^k \sum_{l \neq k} (-1)^l a_{0\dots \widehat{l} \dots n} + (-1)^k \sum_{\substack{\tau \in \mathbb{RT} \\ I_1, \dots, I_m \subset [n]}} \lambda_{\tau, I_1, \dots, I_m} \tau(a_{I_1}, \dots, \varphi(x), \dots, \varphi(x), \dots, a_{I_m})$$

but in the complete \mathfrak{L}_∞ -algebra \mathfrak{mc}_n this time. A solution is known, it is $a_{0\dots \widehat{k} \dots n}$, and by the uniqueness of the solution to the fixed-point equation in Proposition A.4, we conclude that $\varphi(x) = a_{0\dots \widehat{k} \dots n}$. This shows that $\varphi \circ \psi = \text{id}$ and thus that ψ is a chain map. \square

Theorem 5.2. *Let \mathfrak{g} be a complete \mathfrak{L}_∞ -algebra and let $\Lambda_k^n \rightarrow R(\mathfrak{g})$ be a map of simplicial sets. The set of horn fillers is in natural bijection with the set \mathfrak{g}_n of elements of degree n :*

$$\left\{ \begin{array}{ccc} \Lambda_k^n & \longrightarrow & R(\mathfrak{g}) \\ \sim \downarrow & \nearrow & \\ \Delta^n & & \end{array} \right\} \cong \mathfrak{g}_n .$$

Proof. The isomorphism $\mathfrak{mc}_n \cong \mathfrak{L}_\infty(\Lambda_k^n) \hat{\sqcup} \mathfrak{L}_\infty(u, du)$ of Lemma 5.1 and the adjunction of Theorem 2.10 induce the following bijections

$$\begin{aligned} \text{Hom}_{\mathfrak{sSet}}(\Delta^n, R(\mathfrak{g})) &\cong \text{Hom}_{\mathfrak{L}_\infty\text{-alg}}(\mathfrak{L}_\infty(\Delta^n), \mathfrak{g}) \\ &\cong \text{Hom}_{\mathfrak{L}_\infty\text{-alg}}(\mathfrak{L}_\infty(\Lambda_k^n), \mathfrak{g}) \times \text{Hom}_{\mathfrak{L}_\infty\text{-alg}}(\mathfrak{L}_\infty(u, du), \mathfrak{g}) \\ &\cong \text{Hom}_{\mathfrak{L}_\infty\text{-alg}}(\mathfrak{L}_\infty(\Lambda_k^n), \mathfrak{g}) \times \mathfrak{g}_n \\ &\cong \text{Hom}_{\mathfrak{sSet}}(\Lambda_k^n, R(\mathfrak{g})) \times \mathfrak{g}_n , \end{aligned}$$

which is natural in the complete \mathfrak{L}_∞ -algebra \mathfrak{g} . The final bijection from left to right is given by the restriction along the horn inclusion $\Lambda_k^n \hookrightarrow \Delta^n$, which concludes the proof. \square

Corollary 5.3. *The simplicial representation $R(\mathfrak{g})$ of any complete \mathfrak{L}_∞ -algebra \mathfrak{g} is a Kan complex with a canonical algebraic ∞ -groupoid structure.*

Proof. This is a direct corollary of Theorem 5.2: since each \mathfrak{g}_n is none empty, every map $\Lambda_k^n \rightarrow R(\mathfrak{g})$ can be lifted to a map $\Delta^n \rightarrow R(\mathfrak{g})$. So $R(\mathfrak{g})$ is a Kan complex. One can however be more precise here by always considering the canonical choice provided by the element $0 \in \mathfrak{g}_n$. Therefore, the simplicial representation $R(\mathfrak{g})$ comes equipped with a canonical choice of filler for every horn, which is the definition of an algebraic ∞ -groupoid structure. \square

Remark 5.4. *Under the isomorphism of Theorem 3.1, the canonical horn fillers given by the element $0 \in \mathfrak{g}_n$ in the simplicial representation $R(\mathfrak{g})$ correspond to the unique thin fillers of [Get09, Theorem 5.4] in $\gamma_\bullet(\mathfrak{g})$. to be typed, straightforward.*

5.2. Applications: Higher BCH-formulae [Work in progress].

Definition 5.5 (Higher BCH maps). *The bijection of Theorem 5.2 produces a collection of maps denoted by*

$$\begin{aligned} \text{Hom}_{\mathfrak{sSet}}(\Lambda_k^n, R(\mathfrak{g})) \times \mathfrak{g}_n &\rightarrow \mathfrak{g}_{n-1} \\ (\{x_{0\dots \widehat{j} \dots n}\}_{j \neq k}, y) &\mapsto \Gamma_y(x_{0\dots \widehat{n}}, \dots, x_{\widehat{0} \dots n}) . \end{aligned}$$

The map associated to the canonical choice $y = 0 \in \mathfrak{g}_n$ is simply denoted by $\Gamma(x_{0\dots \widehat{n}}, \dots, x_{\widehat{0} \dots n})$.

Let us first show that the present theory recovers the classical Baker–Campbell–Hausdorff formula in the case of complete (dg) Lie algebra.

Proposition 5.6. *Let \mathfrak{g} be a complete dg Lie algebra. The canonical horn filler*

$$\begin{aligned} \text{Hom}_{\mathfrak{sSet}}(\Lambda_1^2, R(\mathfrak{g})) &\rightarrow \mathfrak{g}_1 \\ (x_{01}, x_{12}) &\mapsto \Gamma(x_{01}, x_{12}) = \text{BCH}(x_{01}, x_{12}) . \end{aligned}$$

is equal to the Baker–Campbell–Hausdorff formula.

$$\begin{array}{ccc}
 & x_1 & \\
 x_{01} \nearrow & 0 & \nwarrow x_{12} \\
 x_0 & \xrightarrow{\Gamma(x_{01}, x_{12})} & x_2
 \end{array}$$

Proof. See that the associativity is simple to get... □

6. HOMOTOPY THEORY [BRUNO]

6.1. The model category structure. Let us first recall the classical, aka Quillen–Kan, model category structure on the category \mathbf{sSet} of simplicial sets. We refer the reader to the monographs [Hov99, GJ09] for more details.

Theorem 6.1. [Qui67] *The following three classes of maps form a model category structure on the category \mathbf{sSet} of simplicial sets:*

- a morphism $f : X \rightarrow Y$ of simplicial sets is a weak equivalence if it induces a weak homotopy equivalence of topological spaces $|f| : |X| \xrightarrow{\sim} |Y|$ between the associated geometric realisation.
- a morphism $f : X \hookrightarrow Y$ of simplicial sets is a cofibration if it is made up of injective maps $f_n : X_n \hookrightarrow Y_n$ in each degree.
- a morphism $f : X \rightarrow Y$ of simplicial sets is a fibration if it satisfies the right lifting property with respect to (acyclic) horn inclusion:

$$\begin{array}{ccc}
 \Lambda_k^n & \longrightarrow & X \\
 \downarrow \sim & \nearrow \exists & \downarrow f \\
 \Delta^n & \longrightarrow & Y
 \end{array}$$

This model category structure is cofibrantly generated by the following classes of maps:

- the boundary inclusions $\partial \Delta^n \hookrightarrow \Delta^n$ are the generating cofibrations,
- the acyclic horn inclusions $\Lambda_k^n \hookrightarrow \Delta^n$ are the generating acyclic cofibrations.

We now transfer this cofibrantly generated model category structure to complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras under the \mathfrak{L}_∞ -realisation functor.

Theorem 6.2. *The following three classes of maps form a model category structure on the category of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras:*

- a morphism $f : \mathfrak{g} \rightarrow \mathfrak{h}$ of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras is a weak equivalence if it induces a bijection $\mathcal{MC}(f) : \mathfrak{g} \xrightarrow{\cong} \mathfrak{h}$ between the associated moduli spaces of Maurer–Cartan elements and quasi-isomorphisms $f^\alpha : \mathfrak{g}^\alpha \xrightarrow{\cong} \mathfrak{h}^{f(\alpha)}$, for any Maurer–Cartan element $\alpha \in \mathcal{MC}(\mathfrak{g})$.
- a morphism $f : \mathfrak{g} \rightarrow \mathfrak{h}$ of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras is a fibration if it is made up of epimorphisms $f_n : \mathfrak{g}_n \rightarrow \mathfrak{h}_n$ in degrees $n \geq 2$.
- a morphism $f : \mathfrak{g} \hookrightarrow \mathfrak{h}$ of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras is a cofibration if it satisfies the right lifting property with respect to acyclic fibrations.

This model category structure is cofibrantly generated by the following classes of maps:

- the boundary inclusions $\mathfrak{L}_\infty(\partial \Delta^n) \hookrightarrow \mathfrak{L}_\infty(\Delta^n)$ are the generating cofibrations,
- the horn inclusions $\mathfrak{L}_\infty(\Lambda_k^n) \hookrightarrow \mathfrak{L}_\infty(\Delta^n)$ are the generating acyclic cofibrations.

Proof. Let us begin by proving the last assertion. To do so, we use Quillen’s transfer theorem under left adjoint [Qui67, Section II.4]. The version used here follows from [BM03, Section 2.6]: we need to prove the following four facts.

- (1) *The category of complete $\mathfrak{s}\mathcal{L}_\infty$ -algebras is complete and cocomplete:* this was proved in Proposition 1.3.
- (2) *The functor \mathfrak{L}_∞ preserves small objects :* TBC

- (3) *Every complete \mathfrak{L}_∞ -algebra is fibrant*: this is a direct corollary of Lemma 5.1, see also Theorem 5.2.
- (4) *The category of complete \mathfrak{L}_∞ -algebra has functorial path objects*: for any complete \mathfrak{L}_∞ -algebra \mathfrak{g} , we consider the following composite of morphisms

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{\sim} & \mathfrak{g} \otimes \Omega_1 \longrightarrow \mathfrak{g} \times \mathfrak{g} \\ x & \longmapsto & x \otimes 1 \\ & & x \otimes P(t) \longmapsto (P(0)x, P(1)x) \\ & & x \otimes Q(t)dt \longmapsto 0, \end{array}$$

where we denote $t := t_1$ in Ω_1 . The first map is a quasi-isomorphism of complete \mathfrak{L}_∞ -algebras. By Theorem 3.1, we have a natural weak equivalence $R(\mathfrak{g}) \simeq MC_\bullet(\mathfrak{g})$, and [DR15, Theorem 1.1] shows that Deligne–Hinich functor MC_\bullet sends quasi-isomorphisms to weak equivalences. Therefore, the first map is a weak equivalence of complete \mathfrak{L}_∞ -algebras. The second map is an epimorphism and thus a fibration, see below. So all together this construction forms a functorial path object.

Let us now make the fibrations explicit: the fibrations are the morphisms $f : \mathfrak{g} \twoheadrightarrow \mathfrak{h}$ of complete \mathfrak{L}_∞ -algebras which are made up of epimorphisms $f_n : \mathfrak{g}_n \twoheadrightarrow \mathfrak{h}_n$ in degrees $n \geq 2$. In a transferred cofibrantly generated model category structure under a left adjoint functor, the class of fibrations is the class of maps $f : \mathfrak{g} \twoheadrightarrow \mathfrak{h}$ whose image $R(f)$ under the right adjoint functor is a fibration of simplicial sets, that is the ones which satisfy the right lifting property with respect to the acyclic horn inclusions. We use the same arguments as in Theorem 5.2. First, we consider the equivalent problem on the level of complete \mathfrak{L}_∞ -algebras under the adjunction of Theorem 2.10:

$$\begin{array}{ccc} \mathfrak{L}_\infty(\Lambda_k^n) & \longrightarrow & \mathfrak{g} \\ \downarrow \sim & \searrow \exists & \downarrow f \\ \mathfrak{L}_\infty(\Delta^n) & \longrightarrow & \mathfrak{h} \end{array}.$$

Then, we conclude using Lemma 5.1.

Let us now make weak equivalences explicit. Again, in a transferred cofibrantly generated model category structure under a left adjoint functor, the class of weak equivalences is the class of maps $f : \mathfrak{g} \xrightarrow{\sim} \mathfrak{h}$ whose image $R(f)$ under the right adjoint functor is a weak equivalence of simplicial sets. **TBC. Straightforward**

□

Remark 6.3. *The model category structure of [BFMT16] on complete shifted dg Lie algebras appears as a model sub-category of the present one. and the adjunction also restricts ... also chain complexes via Dold–Kan.*

Generalise here the paper of BFMT from complete dg Lie algebras to complete L_∞ -algebras. Quillen equivalence with simplicial sets. New model of $(\infty, 1)$ -category, examples \mathcal{P}_∞ -algebras.

6.2. Homotopy invariance of the representation functor.

Proposition 6.4.

Proof. **TBC**

□

Theorem 6.5.

Type the ultimate version of Goldman–Milson and Dolgushev–Rogers theorem for filtered ∞ -quasi-isomorphisms: use the π -rectification; just check the compatibility with the underlying filtrations. dit aussi invariance homotopique de "notre" espace de Maurer–Cartan

APPENDIX A. FORMAL EQUATIONS [BRUNO]

A.1. Formal fixed-point equations. Let (V, F) be a complete module, that is $V = F_0 V = F_1 V \supset F_2 V \supset \dots$.

Definition A.1 (Analytic function). *An analytic function $P : V \rightarrow V$ is an application of the form*

$$P(x) = \sum_{n=0}^{\infty} P_n(x^{\otimes n}) ,$$

with $P_n \in \mathcal{F}_1 \text{ hom}(V^{\otimes n}, V)$, that is a linear map satisfying

$$P_n(F_{i_1}, \dots, F_{i_n}) \subset F_{i_1 + \dots + i_n + 1} .$$


We use the notation $p_0 := P_0(1) \in F_1$, with 1 the unit of \mathbb{K} and we consider the set $\text{PT}^{[n]}$ of planar rooted trees with internal vertices having at least one input and such that the total number of vertices and leaves is equal to n .

Example A.2. *The set $\text{PT}^{[4]}$ is made up of the following 5 planar trees:*

$$\text{PT}^{[4]} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} , \begin{array}{c} \diagup \quad \diagdown \\ \bullet \\ | \\ \bullet \end{array} , \begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \\ \bullet \\ | \\ \bullet \end{array} , \begin{array}{c} \bullet \\ | \\ \bullet \\ \diagup \quad \diagdown \\ \bullet \end{array} , \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \end{array} \right\} .$$

To any tree $\tau \in \text{PT}^{[n]}$, we associate the element $P_\tau(p_0, \dots, p_0) \in F_n$ obtained by labelling the vertices of arity k with P_k , the leaves with p_0 , and by considering the global evaluation.

Example A.3. *The planar tree*



produces the element $P_\tau(p_0, \dots, p_0) = P_2(p_0, P_1(p_0)) \in F_4$.

Proposition A.4. *The fixed-point equation*

$$(2) \quad x = P(x) = p_0 + \sum_{n=1}^{\infty} P_n(x^{\otimes n}) .$$

associated to any analytic function admits the following unique solution:

$$x = \sum_{n=1}^{\infty} \sum_{\tau \in \text{PT}^{[n]}} P_\tau(p_0, \dots, p_0) .$$

Proof. Suppose that there exists a solution and let us write it as a series $x = \sum_{n=1}^{\infty} x_n$, with $x_n \in F_n$. The projection of Equation (2) onto V/F_{n+1} , for $n \geq 1$, gives

$$(*) \quad x_1 + \dots + x_n = p_0 + P_1(x_1 + \dots + x_{n-1}) + \dots + P_{n-1}(x_1, \dots, x_1) .$$

By induction on $n \geq 1$, this shows that the projection of x onto V/F_{n+1} is unique, which proves that x is unique since (V, F) is complete. In the other way round, let us consider

$$x_n := \sum_{\tau \in \text{PT}^{[n]}} P_\tau(p_0, \dots, p_0) \in F_n \quad \text{and} \quad x := \sum_{n=1}^{\infty} x_n .$$

We now prove, by induction on $n \geq 1$, that x satisfies the projection $(*)$ of Equation (2) onto V/F_{n+1} . For $n = 1$, the set $\text{PT}^{[0]}$ is made up of the trivial tree $|$, so $x_1 = p_0$. Suppose now that the result holds true for n , and let us prove it for $n + 1$:

$$\begin{aligned} p_0 + P_1(x_1 + \dots + x_n) + \dots + P_n(x_1, \dots, x_1) &= p_0 + P_1(x_1 + \dots + x_{n-1}) + \dots + P_{n-1}(x_1, \dots, x_1) \\ &\quad + P_1(x_n) + P_2(x_1, x_{n-1}) + \dots + P_2(x_{n-1}, x_1) + \dots \\ &\quad + P_{n+1}(x_1, \dots, x_1) \end{aligned}$$

$$\begin{aligned}
&= x_1 + \cdots + x_n + \sum_{\tau \in \mathcal{PT}^{[n+1]}} P_\tau(p_0, \dots, p_0) \\
&= x_1 + \cdots + x_{n+1} .
\end{aligned}$$

□

A.2. Formal differential equations.

REFERENCES

- [Ban14] R. Bandiera. *Higher Deligne groupoids, derived brackets and deformation problems in holomorphic Poisson geometry*. PhD thesis, Università degli studi di Roma “La Sapienza”, 2014.
- [BFMT16] U. Buijs, Y. Félix, A. Murillo, and D. Tanré. Homotopy theory of complete Lie algebras and Lie models of simplicial sets. *ArXiv e-prints*, 2016.
- [BM03] C. Berger and I. Moerdijk. Axiomatic homotopy theory for operads. *Comment. Math. Helv.*, 78(4):805–831, 2003.
- [DR15] Vasily A. Dolgushev and Christopher L. Rogers. A version of the Goldman-Millson theorem for filtered L_∞ -algebras. *J. Algebra*, 430:260–302, 2015.
- [DSV18] Vladimir Dotsenko, Sergei Shadrin, and Bruno Vallette. The twisting procedure. *arXiv:???*, 2018.
- [Dup76] Johan L. Dupont. Simplicial de Rham cohomology and characteristic classes of flat bundles. *Topology*, 15(3):233–245, 1976.
- [Get09] Ezra Getzler. Lie theory for nilpotent L_∞ -algebras. *Ann. of Math. (2)*, 170(1):271–301, 2009.
- [GJ09] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Modern Birkhäuser Classics. Birkhäuser Verlag, Basel, 2009.
- [Hin97] Vladimir Hinich. Descent of Deligne groupoids. *Internat. Math. Res. Notices*, (5):223–239, 1997.
- [Hov99] Mark Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [Kan58] Daniel M. Kan. Functors involving c.s.s. complexes. *Trans. Amer. Math. Soc.*, 87:330–346, 1958.
- [LV12] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*, volume 346 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2012.
- [Nik11] Thomas Nikolaus. Algebraic models for higher categories. *Indag. Math. (N.S.)*, 21(1-2):52–75, 2011.
- [Qui67] Daniel G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [RN17] D. Robert-Nicoud. Representing the Deligne–Hinich–Getzler ∞ -groupoid. 2017. [arXiv:1702.02529](#).
- [RNW17] D. Robert-Nicoud and F. Wierstra. Homotopy morphisms between convolution homotopy Lie algebras. 2017. [arXiv:1712.00794](#).

TO DO LIST:

- Applications : comparer avec ces notions d’équivalences des ∞ -morphisms. Liens avec Dotsenko–Poncin
- Etendre au cas a courbure [Bruno]
- Comprendre BCH (supérieur)
- Holonomie ???
- Quillen equivalence ?
- Lien avec Henriques

LABORATOIRE ANALYSE, GÉOMÉTRIE ET APPLICATIONS, UNIVERSITÉ PARIS 13, SORBONNE PARIS CITÉ, 99 AVENUE JEAN BAPTISTE CLÉMENT, 93430 VILLETANEUSE, FRANCE
E-mail address: robert-nicoud@math.univ-paris13.fr

LABORATOIRE ANALYSE, GÉOMÉTRIE ET APPLICATIONS, UNIVERSITÉ PARIS 13, SORBONNE PARIS CITÉ, 99 AVENUE JEAN BAPTISTE CLÉMENT, 93430 VILLETANEUSE, FRANCE
E-mail address: vallette@math.univ-paris13.fr