

# Assignment 9

PRABHAV SINGH (BT21BTECH11004)

01 June 2022

# Outline

1 Question

2 Solution

# Question

Suppose the Conditional distribution of  $x$  given  $y = n$  is binomial with parameters  $n$  and  $p_1$ . Further,  $Y$  is a binomial random variable with parameters  $M$  and  $p_2$ . Show that the distribution of  $x$  is also binomial. Find its parameters.

# Solution

Using the symbols in their Standard definition and given as in question ,  
 $P(X = k|Y = n) = \binom{n}{k} p_1^k q_1^{n-k}; k = 1, 2, 3, \dots, n$

$$E(e^{j\omega k} | Y = n) = \sum_{k=0}^n e^{j\omega k} P(X = k | Y = n) \quad (1)$$

$$= (p_1 e^{j\omega} + q_1)^n \quad (2)$$

Also we know that ,

$$\phi_x(\omega) = E[e^{j\omega X}] \quad (3)$$

$$= E(E[e^{j\omega x} | Y = n]) \quad (4)$$

$$= \sum_{n=0}^M E(E[e^{j\omega x} | Y = n]) P(Y = n) \quad (5)$$

$$= \sum_{n=0}^{\infty} (p_1 e^{j\omega} + q_1)^n \binom{M}{n} p_2^n q_2^{M-n} \quad (6)$$

$$= \sum_{n=0}^M \binom{M}{n} (p_2(p_1 e^{j\omega} + q_1))^n q_2^{M-n} \quad (7)$$

$$= (p_1 p_2 e^{j\omega} + q_1 p_2 + q_2)^M \quad (8)$$

But

$$(1 - p_1 p_2) = 1 - (1 - q_1)(1 - q_2) \quad (9)$$

$$= q_1 p_2 + q_2 \quad (10)$$

Hence  $\phi_{\omega} = (p e^{j\omega} + q)^m$  where  $p = p_1 p_2$

Thus  $X \sim \text{Binomial}(M, p_1 p_2)$