

Master Theorem in Another Form

Master Theorem

- The order of growth of its solution $T(n)$ depends on the values of the constants a and b and *the order of growth of the function $f(n)$* .

$$T(n) = aT(n/b) + f(n) \quad \text{where } f(n) \in \Theta(n^d), \quad d \geq 0$$

- The efficiency analysis of many divide-and-conquer algorithms is greatly simplified by the following theorem:

Master Theorem: If $a < b^d$, $T(n) \in \Theta(n^d)$
 If $a = b^d$, $T(n) \in \Theta(n^d \log n)$
 If $a > b^d$, $T(n) \in \Theta(n^{\log_b a})$

Note: The same results hold with O instead of Θ .

Master Theorem

$$T(n) = aT(n/b) + f(n) \quad \text{where } f(n) \in \Theta(n^d), \quad d \geq 0$$

$$\text{If } a < b^d, \quad T(n) \in \Theta(n^d)$$

$$\text{If } a = b^d, \quad T(n) \in \Theta(n^d \log n)$$

$$\text{If } a > b^d, \quad T(n) \in \Theta(n^{\log_b a})$$

Solve these recurrence:

$$\cdot \quad T(n) = 4T(n/2) + n^2$$

$$\cdot \quad T(n) = 16T(n/4) + n$$

Master Theorem

$$T(n) = aT(n/b) + f(n) \quad \text{where } f(n) \in \Theta(n^d), \quad d \geq 0$$

$$\text{If } a < b^d, \quad T(n) \in \Theta(n^d)$$

$$\text{If } a = b^d, \quad T(n) \in \Theta(n^d \log n)$$

$$\text{If } a > b^d, \quad T(n) \in \Theta(n^{\log_b a})$$

Solve these recurrence:

$$. \quad T(n) = 4T(n/2) + n^2$$

$$T(n) = 4T(n/2) + n^2 \implies T(n) = \Theta(n^2 \log n)$$

$$. \quad T(n) = 16T(n/4) + n$$

$$T(n) = 16T(n/4) + n \implies T(n) = \Theta(n^2)$$