

Asthma Patient Guide: Breathing Well and Preventing Attacks

Plain-language guide to asthma control, inhaler technique, trigger reduction, action plans, and urgent symptoms

Document ID	PE-PULM-ASTH-003	Version	1.0
Owner	Pulmonary Medicine and Respiratory Therapy	Approved by	Patient Education Committee and Medical Executive Committee
Effective date	16 July 2026	Review date	16 July 2028
Audience	Patients, families, caregivers, nurses, educators, pharmacists, dietitians, and outpatient clinicians	Classification	Patient education reference

Dataset status

This document is a realistic synthetic artifact created to test document ingestion, chunking, retrieval, metadata filtering, reranking, citation, and question-answering workflows. It has not been reviewed or approved for patient care.

Document navigation

1. Understanding Asthma
 3. Controller and Reliever Medicines
 5. Using Dry-Powder, Soft-Mist, and Other Devices
 7. Triggers and Irritants
 9. Your Written Asthma Action Plan
 11. Colds, Flu, COVID-19, and Other Illness
 13. Follow-up and Lung Testing
2. Recognizing Control and Loss of Control
 4. Using a Metered-Dose Inhaler and Spacer
 6. Preventing Local Side Effects
 8. Exercise, Sports, and Daily Activity
 10. Handling an Asthma Attack
 12. Pregnancy, Work, School, and Travel
 14. Common Questions and Personal Checklist

Controlled use

Printed copies are uncontrolled. Users should verify the effective version, approved local order sets, current formulary information, and governing law before applying any content.

1. Understanding Asthma

Asthma is a long-term condition in which the breathing tubes become inflamed and more sensitive. They can narrow because of muscle tightening, swelling, and mucus, causing symptoms that vary over time. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed. [1,2]

Key considerations are: Common symptoms are wheeze, cough, chest tightness, and shortness of breath, but the pattern and intensity differ among people. Use the advice with the individual care plan, because medicines, other illnesses, cost, access, and personal goals can change what is safest. Symptoms may be intermittent while airway inflammation is still present, so feeling well on one day does not always mean preventive treatment is no longer needed. A useful plan states what to do, how often, what result is expected, and which change should prompt a call or urgent assessment. Asthma can usually be controlled with the right inhaled treatment, correct technique, trigger management, and a written action plan. Choose a small change that can be repeated, note practical barriers, and bring the experience to follow-up so the plan can be adjusted.

Within the SVMH care pathway, Ask the clinician to explain the diagnosis, the expected symptom pattern, and how control will be measured. At follow-up, review what was actually done, what was difficult, and whether the

recommendation improved symptoms or measurements. The asthma diagnosis and severity should be reviewed when symptoms, test results, or response to treatment do not fit the expected pattern. Materials should use respectful language and account for disability, culture, literacy, finances, food access, and caregiving duties.

Practice note
This synthetic guide does not replace the patient-specific prescription or written asthma action plan.

2. Recognizing Control and Loss of Control

Good asthma control means few daytime symptoms, little or no night waking, normal or near-normal activity, infrequent reliever use, and low risk of attacks and medicine side effects. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Assessment should address the following: Increasing cough, wheeze, chest tightness, breathlessness, night waking, activity limitation, or reliever use may signal worsening control. A useful plan states what to do, how often, what result is expected, and which change should prompt a call or urgent assessment. A person can have few daily symptoms yet remain at high attack risk because of a prior severe attack, low lung function, smoking, poor access, or inadequate anti-inflammatory treatment. Choose a small change that can be repeated, note practical barriers, and bring the experience to follow-up so the plan can be adjusted. Home peak-flow values may support the action plan for selected patients but should be interpreted against the personal best and symptoms. Consider timing, severity, associated symptoms, recent illness, missed doses, food and fluid intake, and any newly started medicine or product.

For routine implementation, Track symptoms and reliever use for at least one or two weeks before follow-up when control is uncertain. Before discharge, confirm access to prescriptions and supplies, after-hours contact information, and the documented follow-up location and time frame. Repeated oral steroid courses, emergency visits, or reliever overuse require prompt clinician review rather than repeated refills alone. The current prescription label, action plan, discharge instructions, and clinician's directions take priority over general examples.

Table 1. Questions that suggest asthma may not be controlled

Question	Concerning pattern	What to do
Daytime symptoms?	More than occasionally or increasing	Review technique, adherence, triggers, and treatment
Night waking?	Any recurrent asthma-related waking	Arrange assessment and update the action plan
Reliever needed?	Increasing frequency or poor effect	Use action plan; seek urgent help if severe
Activity limited?	Avoiding play, exercise, stairs, or work	Assess control and alternative diagnoses

3. Controller and Reliever Medicines

Controller treatment reduces airway inflammation and future attacks, while reliever treatment is used for symptoms according to the prescribed plan. Some inhalers can serve both roles when specifically prescribed. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed. [1,2]

The practical interpretation is as follows: Inhaled corticosteroid-containing treatment is central to reducing severe attacks and should not be stopped simply because symptoms improve. Choose a small change that can be repeated, note practical barriers, and bring the experience to follow-up so the plan can be adjusted. A short-acting bronchodilator opens the airways temporarily but does not treat the underlying inflammation and should not be the only long-term strategy for most patients. Consider timing, severity, associated symptoms, recent illness, missed doses, food and fluid intake, and any newly started medicine or product. Combination inhalers, leukotriene medicines, biologic therapy, oral corticosteroids, and other treatments have specific

indications, benefits, and risks. Use teach-back: explain the plan in your own words and demonstrate any device or technique before leaving the teaching visit.

During assessment and treatment, Keep a current list that labels each inhaler as daily controller, as-needed reliever, or both, with the exact maximum use and contact threshold. When the plan changes, update the medication list and patient instructions together and remove or clearly mark outdated copies. Pharmacists and clinicians should check dose counter, refill pattern, side effects, device compatibility, and actual use at every review. Repeated difficulty should trigger assessment of burden, side effects, understanding, distress, and access rather than blame.

Warning

Do not borrow another person's inhaler or assume that inhalers with a similar color contain the same medicine. Read the label and follow the individual prescription.

4. Using a Metered-Dose Inhaler and Spacer

Correct inhaler technique determines how much medicine reaches the lungs. A pressurized metered-dose inhaler is often easier to use correctly with a spacer or valved holding chamber. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Relevant clinical features include: Typical steps include removing the cap, checking the mouthpiece, shaking when required, attaching the spacer, breathing out, pressing one puff, and inhaling slowly and deeply or using the taught tidal-breath method. Consider timing, severity, associated symptoms, recent illness, missed doses, food and fluid intake, and any newly started medicine or product. Only one puff should be placed in the chamber at a time unless the device instructions say otherwise, and the chamber should be cleaned and dried as directed. Use teach-back: explain the plan in your own words and demonstrate any device or technique before leaving the teaching visit. A whistle from some spacers may mean the breath is too fast, and visible spray escaping may signal a poor seal or timing problem. Keep the written plan accessible and include current medicines, allergies, contact details, and the symptoms that require emergency evaluation.

At each transition of care, Demonstrate the complete technique with the actual device, then ask the patient to repeat it without coaching. Urgent concerns require direct communication; routine portal messages are not substitutes for emergency services when red-flag symptoms are present. Technique should be rechecked after every device change and periodically even when the patient has used an inhaler for years. The education team reviews questions and safety events to identify wording or access problems that require improvement.

- Sit or stand upright.
- Seal lips around the mouthpiece.
- Hold the breath as comfortably possible after a deep inhalation.

5. Using Dry-Powder, Soft-Mist, and Other Devices

Different inhalers require different preparation and breathing speed. A technique that is correct for one device may be wrong for another. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

The core elements are: Many dry-powder inhalers require a quick deep inhalation and should not be shaken or exposed to moisture, while soft-mist devices have their own loading and coordination steps. Use teach-back: explain the plan in your own words and demonstrate any device or technique before leaving the teaching visit. Capsules intended for an inhaler must not be swallowed, and device-specific dose preparation should be completed exactly as taught. Keep the written plan accessible and include current medicines, allergies, contact details, and the symptoms that require emergency evaluation. Nebulizers may be appropriate in selected situations but require cleaning, maintenance, and an infection-control plan. Monitoring should guide a decision rather than create anxiety; record the result with symptoms and relevant treatment or activity.

When the guideline is applied in practice, Use the manufacturer instructions and a trained demonstration for each exact brand and model rather than relying on a generic inhaler video. At follow-up, review what was actually done, what was difficult, and whether the recommendation improved symptoms or measurements. If hand strength, vision, cognition, inspiratory flow, or coordination limits use, select a more suitable device and document who can assist. Materials should use respectful language and account for disability, culture, literacy, finances, food access, and caregiving duties.

Table 2. Technique differences by device type

Device	Typical inhalation pattern	Common mistake
Metered-dose inhaler	Slow deep breath coordinated with actuation; spacer often helpful	Breathing in too fast or firing before/after the breath
Dry-powder inhaler	Forceful deep inhalation after loading dose	Exhaling into device or inadequate inspiratory effort
Soft-mist inhaler	Slow deep inhalation coordinated with mist release	Incorrect cartridge loading or priming
Nebulizer	Relaxed breathing through fitted mouthpiece or mask	Poor cleaning, leakage, or stopping before dose complete

6. Preventing Local Side Effects

Inhaled medicines are generally well tolerated when used correctly. Local side effects can often be reduced through technique, mouth care, and use of the prescribed device. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Key considerations are: Rinsing the mouth and spitting after inhaled corticosteroid use can reduce oral thrush and hoarseness when recommended for the product. Keep the written plan accessible and include current medicines, allergies, contact details, and the symptoms that require emergency evaluation. A spacer can reduce medicine deposition in the mouth for many pressurized inhalers and may improve delivery to the lungs. Monitoring should guide a decision rather than create anxiety; record the result with symptoms and relevant treatment or activity. Persistent mouth soreness, white patches, voice change, tremor, palpitations, or other suspected effects should be reviewed rather than leading to unplanned discontinuation. Tell the care team early when transport, food, housing, work, language, insurance, supplies, or medicine cost may prevent the plan from working.

Within the SVMH care pathway, Report side effects with the exact product, dose, timing, technique, and effect on daily use so the treatment can be adjusted safely. Before discharge, confirm access to prescriptions and supplies, after-hours contact information, and the documented follow-up location and time frame. The care team should use the lowest treatment intensity that maintains control after stability is confirmed and risk is reviewed. The current prescription label, action plan, discharge instructions, and clinician's directions take priority over general examples.

7. Triggers and Irritants

Triggers do not cause all asthma, and complete avoidance is rarely possible. The goal is to identify exposures linked to symptoms and reduce those that are important and feasible. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Assessment should address the following: Common triggers include viral infection, tobacco or vape aerosol, air pollution, allergens, cold air, exercise, strong fumes, occupational exposure, and some medicines. Monitoring should guide a decision rather than create anxiety; record the result with symptoms and relevant treatment or activity. Routine expensive environmental measures are not helpful unless there is evidence that the specific exposure matters for that patient. Tell the care team early when transport, food, housing, work, language, insurance, supplies, or medicine cost may prevent the plan from working. Smoking cessation and avoiding indoor smoke or aerosol exposure are high-priority interventions because they affect symptoms, treatment response,

and long-term lung health. Use the advice with the individual care plan, because medicines, other illnesses, cost, access, and personal goals can change what is safest.

For routine implementation, Use a symptom and exposure diary to test suspected patterns and discuss targeted changes with the clinical team. When the plan changes, update the medication list and patient instructions together and remove or clearly mark outdated copies. Work-related symptom patterns require timely occupational assessment because continued exposure can worsen disease and affect employment decisions. Repeated difficulty should trigger assessment of burden, side effects, understanding, distress, and access rather than blame.

- Check local air-quality alerts when pollution worsens symptoms.
- Treat allergic rhinitis when present.
- Keep vaccines current according to the clinical plan.

8. Exercise, Sports, and Daily Activity

People with well-controlled asthma should be able to participate in exercise and sports. Symptoms during activity may indicate inadequate control, poor warm-up, cold-air exposure, or another condition. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

The practical interpretation is as follows: A gradual warm-up, appropriate controller treatment, and use of the prescribed pre-exercise reliever can prevent symptoms for many people. Tell the care team early when transport, food, housing, work, language, insurance, supplies, or medicine cost may prevent the plan from working. Stopping all activity can reduce fitness and increase breathlessness from deconditioning, making it harder to distinguish asthma from poor conditioning. Use the advice with the individual care plan, because medicines, other illnesses, cost, access, and personal goals can change what is safest. Chest pain, fainting, severe breathlessness, bluish color, or symptoms that do not improve with the action plan require urgent evaluation. A useful plan states what to do, how often, what result is expected, and which change should prompt a call or urgent assessment.

During assessment and treatment, Record which activities cause symptoms, how quickly they begin, the reliever response, and whether symptoms occur at rest or overnight. Urgent concerns require direct communication; routine portal messages are not substitutes for emergency services when red-flag symptoms are present. Athletes should check applicable competition rules for restricted medicines or documentation requirements without compromising needed care. The education team reviews questions and safety events to identify wording or access problems that require improvement.

9. Your Written Asthma Action Plan

An asthma action plan describes routine treatment and the exact steps for worsening symptoms. It is often organized into green, yellow, and red zones using symptoms, reliever use, and sometimes peak flow. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Relevant clinical features include: The green zone lists daily medicines and the signs of good control, while the yellow zone lists early worsening and temporary treatment changes authorized by the clinician. Use the advice with the individual care plan, because medicines, other illnesses, cost, access, and personal goals can change what is safest. The red zone lists severe symptoms, emergency medicine instructions, and when to call emergency services or go to an emergency department. A useful plan states what to do, how often, what result is expected, and which change should prompt a call or urgent assessment. The plan should include the patient's usual best peak flow when used, medicine names rather than colors alone, maximum doses, contact information, and the date. Choose a small change that can be repeated, note practical barriers, and bring the experience to follow-up so the plan can be adjusted.

At each transition of care, Complete the plan with the prescribing clinician, practice a sample worsening scenario, and provide copies for home, school, work, or caregivers as appropriate. At follow-up, review what was actually done, what was difficult, and whether the recommendation improved symptoms or measurements. Update the action plan after an emergency visit, hospitalization, oral steroid course, pregnancy, major medicine change, or recurring loss of control. Materials should use respectful language and account for disability, culture, literacy, finances, food access, and caregiving duties.

Table 3. Action-plan zones

Zone	Typical meaning	Required information in personal plan
Green	Doing well; normal activity and sleep	Routine controller/reliever plan and prevention measures
Yellow	Symptoms increasing or peak flow below personal target	Temporary medicine steps, monitoring, and call threshold
Red	Severe symptoms, poor response, or very low peak flow	Emergency medicine, emergency contact, and immediate destination

10. Handling an Asthma Attack

An asthma attack may progress quickly. Early use of the written plan and prompt escalation are safer than waiting until speaking or walking becomes difficult. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

The core elements are: Sit upright, stay with the person, use the prescribed reliever exactly as directed, and monitor response and ability to speak, breathe, and remain alert. A useful plan states what to do, how often, what result is expected, and which change should prompt a call or urgent assessment. Call emergency services for severe breathlessness, inability to speak in full phrases, bluish lips, drowsiness, collapse, a silent chest, or poor response to initial reliever treatment. Choose a small change that can be repeated, note practical barriers, and bring the experience to follow-up so the plan can be adjusted. Do not drive yourself when severe symptoms, faintness, confusion, or repeated reliever doses make travel unsafe. Consider timing, severity, associated symptoms, recent illness, missed doses, food and fluid intake, and any newly started medicine or product.

When the guideline is applied in practice, After any significant attack, arrange clinical review to identify the trigger, assess technique and access, and adjust preventive treatment. Before discharge, confirm access to prescriptions and supplies, after-hours contact information, and the documented follow-up location and time frame. Emergency and discharge teams should provide an updated action plan, controller supply, device education, and time-specific follow-up. The current prescription label, action plan, discharge instructions, and clinician's directions take priority over general examples.

Warning

Severe asthma can be life-threatening even without a loud wheeze. A quiet or silent chest with increasing effort is an emergency.

11. Colds, Flu, COVID-19, and Other Illness

Respiratory infections commonly worsen asthma. Preventive medicines usually continue during illness unless a clinician gives different instructions. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Key considerations are: Follow the yellow-zone plan early when cough, wheeze, chest tightness, night symptoms, or reliever use increases. Choose a small change that can be repeated, note practical barriers, and bring the experience to follow-up so the plan can be adjusted. Maintain hydration, monitor fever and breathing, and seek advice about testing or antiviral treatment when relevant to the illness and risk level. Consider timing, severity, associated symptoms, recent illness, missed doses, food and fluid intake, and any newly started medicine or product. Use separate devices when possible and clean spacers or nebulizer equipment according to instructions to reduce infection spread. Use teach-back: explain the plan in your own words and demonstrate any device or technique before leaving the teaching visit.

Within the SVMH care pathway, Ask the clinician how illness changes the action plan, including when an oral corticosteroid or urgent assessment is appropriate. When the plan changes, update the medication list and patient instructions together and remove or clearly mark outdated copies. Vaccination recommendations should be reviewed based on age, risk, current public-health guidance, and contraindications. Repeated difficulty should trigger assessment of burden, side effects, understanding, distress, and access rather than blame.

12. Pregnancy, Work, School, and Travel

Asthma control remains important during pregnancy and in changing environments. Planned treatment is generally safer than uncontrolled asthma or abrupt medicine discontinuation. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Assessment should address the following: Pregnant patients should continue prescribed asthma treatment and obtain prompt review for worsening symptoms, because low oxygen can affect both parent and fetus. Consider timing, severity, associated symptoms, recent illness, missed doses, food and fluid intake, and any newly started medicine or product. Schools and workplaces may need an action plan, permission for access to rescue medicine, trigger controls, and clear emergency contacts. Use teach-back: explain the plan in your own words and demonstrate any device or technique before leaving the teaching visit. Travel planning includes adequate inhaler supply, prescriptions, spacer, action plan, insurance information, and awareness of altitude, climate, smoke, or pollen exposure. Keep the written plan accessible and include current medicines, allergies, contact details, and the symptoms that require emergency evaluation.

For routine implementation, Arrange review before pregnancy, extended travel, relocation, or a workplace exposure change when possible. Urgent concerns require direct communication; routine portal messages are not substitutes for emergency services when red-flag symptoms are present. Confidentiality should be respected while ensuring that designated people know how to respond to an emergency. The education team reviews questions and safety events to identify wording or access problems that require improvement.

13. Follow-up and Lung Testing

Asthma should be reviewed after starting or changing treatment and periodically when stable. Follow-up assesses symptoms, attacks, lung function, technique, adherence, side effects, triggers, and goals. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

The practical interpretation is as follows: Spirometry or another lung-function test may confirm variable airflow limitation, assess risk, or investigate symptoms that do not match the expected pattern. Use teach-back: explain the plan in your own words and demonstrate any device or technique before leaving the teaching visit. Bring all inhalers, spacer, action plan, refill history, symptom record, and peak-flow log if used so the clinician can evaluate actual treatment. Keep the written plan accessible and include current medicines, allergies, contact details, and the symptoms that require emergency evaluation. Treatment may be stepped up when control is poor and stepped down cautiously after sustained control, but only after checking technique, adherence, exposure, and alternative diagnoses. Monitoring should guide a decision rather than create anxiety; record the result with symptoms and relevant treatment or activity.

During assessment and treatment, Schedule early review after an attack and routine review at the interval in the action plan. At follow-up, review what was actually done, what was difficult, and whether the recommendation improved symptoms or measurements. Specialist referral is appropriate for uncertain diagnosis, severe or difficult-to-treat asthma, repeated attacks, major side effects, occupational concerns, or consideration of biologic treatment. Materials should use respectful language and account for disability, culture, literacy, finances, food access, and caregiving duties.

Table 4. What to bring to an asthma review

Item	What it shows	Common decision supported
All inhalers and spacer	Exact device, dose counter, technique, condition	Correct technique or change device

Item	What it shows	Common decision supported
Symptom/reliever record	Frequency, timing, triggers, activity and sleep impact	Assess control and action-plan zone
Attack history	Emergency visits, steroids, admissions, intensive care	Estimate future risk and need for referral
Test results	Baseline and variability in airflow	Confirm diagnosis and monitor risk

14. Common Questions and Personal Checklist

Asthma care is more reliable when the patient can explain the diagnosis, identify each inhaler, demonstrate technique, and describe the action-plan response to worsening symptoms. This section explains the practical choices a patient or caregiver may need to make about asthma self-management and identifies when professional help is needed.

Relevant clinical features include: Patients should know whether the reliever contains an inhaled corticosteroid, how many inhalations may be used, and what maximum or call threshold applies. Keep the written plan accessible and include current medicines, allergies, contact details, and the symptoms that require emergency evaluation. Patients should know how to check remaining doses, obtain refills before running out, store devices, clean equipment, and replace expired rescue medicine. Monitoring should guide a decision rather than create anxiety; record the result with symptoms and relevant treatment or activity. Patients should know the emergency warning signs and which person, clinic, or emergency service to contact during the day and after hours. Tell the care team early when transport, food, housing, work, language, insurance, supplies, or medicine cost may prevent the plan from working.

At each transition of care, Use teach-back and a device demonstration before leaving the visit, and write unresolved questions for the next contact. Before discharge, confirm access to prescriptions and supplies, after-hours contact information, and the documented follow-up location and time frame. Care teams should address cost, language, dexterity, cognition, housing, smoke exposure, and transportation when these barriers interfere with control. The current prescription label, action plan, discharge instructions, and clinician's directions take priority over general examples.

Practice note

A well-controlled patient still keeps reliever medicine and an action plan available, because attacks can occur unexpectedly.

15. References

References are provided to make the synthetic document realistic and to support citation and provenance tests. They should be checked against the most current source before any real-world use.

[1] Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention, 2026 update.

[2] Global Initiative for Asthma. Summary Guide for Asthma Management and Prevention, 2026.

[3] National Heart, Lung, and Blood Institute. Asthma Management Guidelines and patient resources.

[4] U.S. Food and Drug Administration. Current prescribing information and device instructions for inhaled asthma medicines.

[5] Centers for Disease Control and Prevention. Asthma and respiratory infection patient resources. Accessed 2026.

Reference status

External guidelines and regulations may change between scheduled reviews. The document owner is responsible for checking high-impact updates and initiating an interim revision when necessary.