Data Representing Atmospheric Conditions on Mars

Pradyumnan R

JNCASR

May 24, 2024



- **1** General Conditions
- 2 Descent Characteristics
- 3 Sample Analysis at Mars (SAM)
- 4 References

- **1** General Conditions
- 2 Descent Characteristics
- Sample Analysis at Mars (SAM)
- 4 References

Pradyumnan R

Surface temperatures generally range between 166 K and 256 K.

- Surface temperatures generally range between 166 K and 256 K.
- Mean atmospheric pressure is about 600 Pa.

- Surface temperatures generally range between 166 K and 256 K.
- Mean atmospheric pressure is about 600 Pa.
- Very low thermal inertia, temperature swings of 100 K.

- Surface temperatures generally range between 166 K and 256 K.
- Mean atmospheric pressure is about 600 Pa.
- Very low thermal inertia, temperature swings of 100 K.
- Presence of katabatic winds

- \bullet Surface temperatures generally range between 166 K and 256 K.
- Mean atmospheric pressure is about 600 Pa.
- Very low thermal inertia, temperature swings of 100 K.
- Presence of katabatic winds
- Atmospheric electricity

- General Conditions
- 2 Descent Characteristics
- Sample Analysis at Mars (SAM)
- 4 References

Description

 The descent characteristics are obtained from the data collected by NASA's Curiosity rover from the Planetary Data System [1].

Description

- The descent characteristics are obtained from the data collected by NASA's Curiosity rover from the Planetary Data System [1].
- The descent trajectory is between $-3.91^{\circ}N$ to $-4.59^{\circ}N$ and $126.56^{\circ}E$ to $137.32^{\circ}E$.



- The descent characteristics are obtained from the data collected by NASA's Curiosity rover from the Planetary Data System [1].
- The descent trajectory is between $-3.91^{\circ}N$ to $-4.59^{\circ}N$ and $126.56^{\circ}E$ to $137.32^{\circ}E$.
- The variations of temperature, pressure and density during descent are highlighted in the next few slides.

Temperature




Fig. 1: Temperature During Descent

Descent Characteristics of Pressure Logarithmic Pressure (in Pa) 5.0 Fitted Logarithmic Pressure (in Pa) Logarithm of Atmospheric Pressure (in Pa) 2.5 0.0 -2.5 -5.0 -7.5 -10.0 20 120 140 Height Above CoM of Mars (in km) →

Fig. 2: Pressure During Descent

Density




Fig. 3: Density During Descent

•00000

- 1 General Conditions
- 2 Descent Characteristics
- 3 Sample Analysis at Mars (SAM)
- 4 References

Description

 All the data is collected by NASA's Curiosity rover and stored in the Mars Science Laboratory's Reduced Data Records repository [2].

Description

 All the data is collected by NASA's Curiosity rover and stored in the Mars Science Laboratory's Reduced Data Records repository [2].

000000


 The final data presented are the volume mixing ratios of various compounds in the Martian atmosphere obtained using the Quadrupole Mass Spectrometer (QMS) present on the rover.


INCASR Pradvumnan R

Description


- All the data is collected by NASA's Curiosity rover and stored in the Mars Science Laboratory's Reduced Data Records repository [2].
- The final data presented are the volume mixing ratios of various compounds in the Martian atmosphere obtained using the Quadrupole Mass Spectrometer (QMS) present on the rover.
- The next few slides give few representative plots of the volume mixing ratios of various compounds collected across 29 QMS experiments.

Volume Mixing Ratios






Volume Mixing Ratios



Volume Mixing Ratios



- General Conditions
- **3** Sample Analysis at Mars (SAM)
- 4 References

- C Holstein-Rathlou, A Maue, and P Withers.
 Atmospheric studies from the mars science laboratory entry, descent and landing atmospheric structure reconstruction. Planetary and Space Science, 120:15–23, 2016.
- [2] SAM Reduced Data Record RDR. Mars science laboratory (msl) software interface specification. 2013.