

✓ What is/are advantage(s) of Locally Weighted Regression? *

1/1

☐ (A) Pointwise approximation of complex target function

☐ (B) Earlier data has no influence on the new ones

☒ (C) Both A & B

☐ (D) none of these



✓ The quality of the result depends on (LWR) *

1/1

☐ Choice of the function

☐ Choice of the kernel function K

☐ Choice of the hypothesis space H

☒ All of these



✓ Algorithm step used by locally weighted regression. *

1/1

- ☐ 1. Given training data $D=\{x_i, y_i\}$, Kernel function $K(\cdot, \cdot)$ and input x
- ☐ 2. Fit weighted regression $w^{\wedge}(x) = \operatorname{argmin}_w \sum_{i=1}^n K(x, x_i) (w^T x_i - y_i)^2$
- ☐ 3. Return regression prediction $w^{\wedge}(x)^T x$.
- ☐ 4. Fit weighted logistic regression $w^{\wedge}(x) = \operatorname{argmin}_w \sum_{i=1}^n K(x, x_i) \log(1 + \exp\{-y_i w^T x_i\})$
- ☐ 5. Return logistic regression prediction $\operatorname{sign}(w^{\wedge}(x)^T x)$.
- ☒ step 1, 2, and 3 ✓
- ☐ step 1, 4 and 5
- ☐ step 1, 2, 4, and 5
- ☐ all the 5 steps need

✓ True-False: Is Logistic regression a supervised machine learning algorithm?

1/1

- ☒ True ✓
- ☐ False



✓ What is “K” in KNN algorithm? *

1/1

- ☐ K = Number of nearest neighbors you want to select to train the class of a given item
- ☐ K = Number of nearest neighbors you want to select to test the class of a given item
- ☐ K = Number of nearest neighbors you want to select to cluster the class of a given item
- ☒ K = Number of nearest neighbors you want to select to predict the class of a given item ✓

✓ Disadvantages of KNN *

1/1

- ☐ Does not work well with large dataset:
- ☐ Does not work well with high dimensions:
- ☐ Need feature scaling
- ☐ Sensitive to noisy data, missing values and outliers
- ☒ all the above ✓



✓ How do we decide the value of “K” in KNN algorithm? *

1/1

- ☐ 1.K should be the square root of n .
- ☐ 2.. K should be odd so that there are no ties. If square root is even, then add or subtract 1 to it.
- ☐ 3.1 and 4
- ☐ 4. K should be even so that there are no ties.
- ☒ 1 and 2



✓ Which of the following distance measure do we use in case of categorical variables in k-NN? 1.Hamming Distance 2.Euclidean Distance 3.Manhattan Distance *

1/1

- ☒ 1
- ☐ 2
- ☐ 3
- ☐ 1 and 2
- ☐ 2 and 3
- ☐ 1,2 and 3



This form was created inside of M S Engineering College.

Google Forms

