

# Agroforestry is paying off – Economic evaluation of ecosystem services in European landscapes with and without agroforestry systems

Sonja Kay<sup>a,\*</sup>, Anil Graves<sup>b</sup>, João H.N. Palma<sup>c</sup>, Gerardo Moreno<sup>d</sup>, José V. Rocas-Díaz<sup>a,e</sup>,  
Stéphanie Aviron<sup>f</sup>, Dimitrios Chouvardas<sup>g</sup>, Josep Crous-Duran<sup>c</sup>, Nuria Ferreiro-Domínguez<sup>h</sup>,  
Silvestre García de Jalón<sup>b</sup>, Vlad Măcicășan<sup>i</sup>, María Rosa Mosquera-Losada<sup>h</sup>, Anastasia Pantera<sup>j</sup>,  
Jose Javier Santiago-Freijanes<sup>h</sup>, Erich Szerencsits<sup>a</sup>, Mario Torralba<sup>k</sup>, Paul J. Burgess<sup>b</sup>,  
Felix Herzog<sup>a</sup>

<sup>a</sup> Agroscope, Department of Agroecology and Environment, Reckenholzstrasse 191, 8046 Zurich, Switzerland

<sup>b</sup> Cranfield University, Cranfield, Bedfordshire MK43 0AL, United Kingdom

<sup>c</sup> Forest Research Centre, School of Agriculture, University of Lisbon, 1349-017 Lisbon, Portugal

<sup>d</sup> Forestry Research Group, Universidad de Extremadura, 10600 Plasencia, Spain

<sup>e</sup> Department of Geography, College of Science, Swansea University, Singleton Park, Swansea SA2 8PP, UK

<sup>f</sup> UMR BAGAP Inra-Agrocampus Ouest-ESA, 35042 Rennes Cedex, France

<sup>g</sup> Laboratory of Rangeland Ecology, Department of Forestry and Natural Environment, Aristotle University, 54 124 Thessaloniki, Greece

<sup>h</sup> Crop Production and Project Engineering Department, Universidad de Santiago de Compostela, 27002 Lugo, Spain

<sup>i</sup> Faculty of Environmental Science and Engineering, Babeș-Bolyai University, 400294 Cluj-Napoca, Romania

<sup>j</sup> Agricultural University of Athens, Karpenissi 36100, Greece

<sup>k</sup> University of Kassel, Faculty of Organic Agricultural Sciences, 37213 Witzenhausen, Germany

## ARTICLE INFO

### Keywords:

Biomass production  
Carbon storage  
Soil loss  
External cost  
Nutrient loss  
Pollination deficit

## ABSTRACT

The study assessed the economic performance of marketable ecosystem services (ES) (biomass production) and non-marketable ecosystem services and dis-services (groundwater, nutrient loss, soil loss, carbon sequestration, pollination deficit) in 11 contrasting European landscapes dominated by agroforestry land use compared to business as usual agricultural practice. The productivity and profitability of the farming activities and the associated ES were quantified using environmental modelling and economic valuation. After accounting for labour and machinery costs the financial value of the outputs of Mediterranean agroforestry systems tended to be greater than the corresponding agricultural system; but in Atlantic and Continental regions the agricultural system tended to be more profitable. However, when economic values for the associated ES were included, the relative profitability of agroforestry increased. Agroforestry landscapes: (i) were associated to reduced externalities of pollution from nutrient and soil losses, and (ii) generated additional benefits from carbon capture and storage and thus generated an overall higher economic gain. Our findings underline how a market system that includes the values of broader ES would result in land use change favouring multifunctional agroforestry. Imposing penalties for dis-services or payments for services would reflect their real world prices and would make agroforestry a more financially profitable system.

## 1. Introduction

The European agricultural economy relies on revenue from the sale of its agricultural products and thus its success is strongly linked to global prices (Hill and Bradley, 2015). The minimum price at which it is profitable to supply these products depends on production costs such as labour, machinery, and fertilisers and other agrochemical inputs. The negative environmental effects or dis-services associated with

agricultural production, such as pollution from fertiliser, soil degradation, and biodiversity losses (Zhang et al., 2007), also known as external costs, are not included in the prices paid for agricultural products, and are often experienced by third parties (Tilman et al., 2002; Zander et al., 2016).

During recent decades, the European Common Agricultural Policy (CAP) has provided financial support for agricultural production and rural development (European Commission, 2016). Although an

\* Corresponding author.

E-mail address: [Sonja.kay@agroscope.admin.ch](mailto:Sonja.kay@agroscope.admin.ch) (S. Kay).

<https://doi.org/10.1016/j.ecoser.2019.100896>

Received 13 March 2018; Received in revised form 14 January 2019; Accepted 23 January 2019

Available online 02 February 2019

2212-0416/ © 2019 Elsevier B.V. All rights reserved.

increasing share of those payments is linked to environmental performance of farming (pillar II, cross compliance), the effectiveness and efficiency of those financial instruments is regularly questioned (Pe'er et al., 2017). It is therefore anticipated that the next funding period (post 2020) will further strengthen the link between financial support and the improvement of the environment and social well-being, as well as addressing climate change (Council of the European Union, 2017).

Agroecological practices, often based on lower agrochemical inputs and higher labour inputs, are increasingly highlighted as promising agricultural systems to reach the goal of environmental and social improvement and favour ecosystem services (ES) (Wezel et al., 2014). ES are defined as the provisioning, regulating and cultural benefits human-beings obtain from ecosystems (MEA, 2003; Haines-Young and Potschin, 2013). However, these agro-ecological systems are often less profitable than intensive production systems under current subsidy and price schemes and this can hamper their adoption (Ponisio et al., 2014). One example of an agro-ecological multi-functional approach are agroforestry systems. Agroforestry is the incorporation of woody elements on agricultural fields; it simultaneously generates food, fodder, and woody material (Somarriba, 1992; European Commission, 2013). Moreover, agroforestry can provide ES and multi-environmental functions such as erosion control, reduced nutrient loss, and carbon storage (Torralba et al., 2016) and is thus valued by farmers (García de Jalón et al., 2018a; Rois-Díaz et al., 2018).

Currently, these environmental benefits from agro-ecological approaches that promote ES are typically not monetarized and hence are not included in the market value of the most profitable production system. Palma et al. (2007) integrated monetary and environmental benefits in a multicriteria analysis and concluded that – if they were well designed – agroforestry systems are the preferable land use when environmental benefits are accounted for. In 2010, the Economics of Ecosystems and Biodiversity Report (TEEB, 2010) valued services perceived as goods by human beings and distinguished between use and non-use values. According to neoclassical economics the use value was separated into (i) direct use value, (ii) indirect use value and (iii) (quasi) option value. The first two features are premised on market-based cost methods, the last one uses mitigation or non-market cost methods. The ES valuation approach adopted a use value perspective, evolved to a monetary valuation and ended as exchange value or commodity (Costanza et al., 2017). The question remains of how to cash ES in markets (Gómez-Baggethun et al., 2010; Muradian et al., 2010). E.g. Costanza et al. (1997) proposed general values per biome and ecosystem service derived from “willingness-to-pay” studies. The authors already pointed out that the approach is limited for valuing public goods of which people might not be aware such as clean air, clean water, or climate regulation. In recent literature, valuing schemes for ES are divided into payments for ES such as price-based incentives for watershed protection (Bennett et al., 2014) or carbon sequestration (Caparros et al., 2007) and markets for ES e.g. carbon emission trading (Boyce, 2018). These payment schemes suffer the problem that e.g. the causal relationship between land use and its service is difficult to define (Muradian et al., 2010) and incomplete information leads to uncertainties and estimations of values (Gómez-Baggethun et al., 2010). However, prices are a tool to value products or services and summarize different ES into one common unit. In the case of carbon markets, prices are also used to regulate emissions (Boyce, 2018). Transparent comparisons including both market and non-market values associated with agricultural production are therefore needed for socially beneficial decision-making (e.g. Brenner et al., 2010; Zander et al., 2016).

This study assessed the monetary use values composed of producer surplus and mitigation costs of provisioning and regulating ES for landscapes with and without agroforestry systems. Taking eleven traditional agroforestry landscapes in Europe as an example, we assessed one marketable ES (biomass production) and five non-marketable ES and dis-services (groundwater, nutrient loss, soil loss, carbon sequestration, and pollination deficit) in landscape test sites with and without

agroforestry in each region. This research investigated three specific questions: 1) Can sales of marketable ES from agroforestry landscapes match those of landscapes dominated by “business-as-usual” agriculture under current market conditions in different parts of Europe? 2) Do these results change when valuing the (non-market) regulating ES services and dis-services? 3) How sensitive are the results to changes in ES prices?

## 2. Material and methods

In order to capture the environmental variability and the diversity of agroforestry systems, the study was undertaken in eleven case study regions ( $> 50 \text{ km}^2$ ) across the Mediterranean, Continental, and Atlantic regions of Europe. In each case study region, eight landscape test sites (LTS) of  $1 \text{ km} \times 1 \text{ km}$  were randomly selected, of which four LTS were dominated by agricultural land (NAF, non-agroforestry) and the other four were dominated by agroforestry land (AF). In the NAF LTS the typical agricultural practice of the specific region was analysed and assessed as economic baseline and represents the “business as usual (BAU) alternative”. The selection process and further data on each case study region are presented by Moreno et al. (2017).

A total of 88 LTS were assessed, of which 44 NAF LTS provided the economic BAU baseline. In all LTS, the habitats and agroforestry trees were mapped, and ES indicators modelled. In this context the landscape scale represents the aggregation of the four NAF and the four AF LTS, respectively, in a case study region.

### 2.1. Case study regions

The study regions represent a wide range of agroforestry systems in Europe including scattered wood pastures (e.g. broadleaf-trees in dehesas in Spain or coniferous trees in Switzerland), high value trees systems (e.g. cherry orchards in Switzerland, olives groves in Greece), and wind break systems (e.g. bocage in France or hedgerows in the United Kingdom) as listed in Table 1 and shown in Fig. 1.

### 2.2. Ecosystem service indicators

One marketable (biomass production) and five non-marketable ES and dis-services (groundwater, nutrient loss, soil loss, carbon sequestration, and pollination deficit) were assessed. The EcoYield-SAFE agroforestry model (Palma et al., 2018) was used to predict biomass production [Unit:  $\text{t ha}^{-1} \text{ a}^{-1}$  separately for crop and/or woody material] and aboveground carbon storage [Unit:  $\text{t C ha}^{-1} \text{ a}^{-1}$ ]. Below-ground carbon storage was predicted by YASSO 0.7 (Liski et al., 2005) [Unit:  $\text{t C ha}^{-1} \text{ a}^{-1}$ ]. The groundwater recharge [Unit:  $\text{mm ha}^{-1} \text{ a}^{-1}$ ] was based on the general water balance including the evapotranspiration equation by FAO (Allen et al., 1998). Nutrient leaching [Unit:  $\text{kg N ha}^{-1} \text{ a}^{-1}$ ] was determined by the MODIFFUS 2.0 model (Hürdlér et al., 2015), the RUSLE equation (Renard et al., 1997) was used to assess soil loss [Unit:  $\text{t ha}^{-1} \text{ a}^{-1}$ ], and the pollination service assessment was based on the Lonsdorf equation (Lonsdorf et al., 2009). A spatially explicit model (resolution  $2 \times 2 \text{ m}$ ) was used to model these six indicators in 88 LTS (8 LTS  $\times$  11 regions) (Kay et al., 2018a,b).

The economic assessment was based on the biophysical evaluation of the six modelled ES indicators. A potential double counting of ES values as highlighted by Fu et al. (2011) was narrowed by using as far as possible independent and static models for each indicator. They were estimated as summarized in the two following sections.


### 2.3. Valuation and prices of market ecosystem services

The market value of biomass production for food, fodder and woody components was calculated using FAO's compendium “Producer Prices – Annual per Country” for each crop (FAO, 2017a), the UNECE/FAO TIMBER database “Wood Prices” (UNECE/FAO, 2017) for timber and

**Table 1**

Case study regions and the dominating agricultural (NAF, business as usual) and agroforestry (AF, alternative) system.

Biogeo-graphical region	Country	Abb.	Agricultural NAF landscapes (Business as Usual, BAU Baseline)	Agroforestry, AF landscapes (Alternative I)
Mediterranean	Portugal	PT	Open pasture	Montado – Wood pasture (Cork oak, <i>Quercus suber</i> L.)
	Greece	GR	Intensive olive groves ( <i>Olea europaea</i> L.)	Intercrop olive groves ( <i>Olea europaea</i> L.)
	Spain	ES1	Open pasture	Dehesa – Wood pasture (Holm oak, <i>Quercus ilex</i> L.)
	Spain	ES2	Arable farming	Intercrop oak (Holm oak, <i>Quercus ilex</i> L.)
Continental	Romania	RO	Open pasture	Wood pasture (Common Oak, <i>Quercus robur</i> L.)
	Switzerland	CH1	Open pasture and arable farming	Fruit orchard (Cherry, <i>Prunus avium</i> L.)
	Germany	GE	Arable farming	Hedgerow landscape with arable farming (mixed species)
	Switzerland	CH2	Open pasture	Wood pasture (Spruce, <i>Picea abies</i> L.)
Atlantic	France	FR	Mixed arable-pasture systems	Bocage - Mixed arable-pasture systems fenced by hedgerows (mixed species)
	Spain	ES3	Open pasture and arable farming	Chestnut soutos ( <i>Castanea sativa</i> Miller)
	United Kingdom	UK	Arable farming	Hedgerow landscape with arable farming (mixed species)

**Fig. 1.** Location of the eleven case study regions.

the farm accountancy data network (FADN) index “Total output / Total input (SE132)” (FADN, 2017). The FADN index accounts for the monetary benefit of crop and livestock production and the specific costs. Further information can be found in [Supplementary material](#). Overheads are provided on an annual basis for each European country. This was then used to recalculate the general net profit of crop and timber products by excluding machinery and labour input, which were included in the price datasets. All values are mean values of the years 2010–2014.

The net financial benefit of biomass production per unit weight (Units: € t<sup>-1</sup>) was determined from the difference between the total output and the total input, which was derived from the total output divided by the FADN index (Eq. (1)).

$$B_{\text{Biomass}} = T_o - \frac{T_o}{i} \quad (1)$$

$B_{\text{Biomass}}$  = Benefits of biomass production per tonne [€ t<sup>-1</sup>]  
 $T_o$  = total output = FAO Producer Prices per crop [€ t<sup>-1</sup>]  
 $i$  = FADN index (Farm Accountancy Data Network)

Based on these assumptions, the net financial benefit of biomass production ranged from 0.43 € t<sup>-1</sup> for wood chips in Switzerland to 802.6 € t<sup>-1</sup> for walnuts in Greece.

## 2.4. Valuation and potential prices of non-market ecosystem services and dis-services

### 2.4.1 Groundwater recharge

Depending on the availability and quality of water resources, the prices per unit indicated in literature varied from 0 to > 4 € m<sup>-3</sup> depending on the specific country (Roo et al., 2012; JRC Water Portal, 2017).

### 2.4.2 Carbon storage

During recent years, the value of a tonne of carbon dioxide (CO<sub>2eq</sub>, 3.7C) traded on the European Energy Exchange (EEX) ranged from 2.95 to 8.54 € t<sup>-1</sup>, with a mean value of about 5 € t<sup>-1</sup> CO<sub>2eq</sub> or 18.5 € t<sup>-1</sup> C (EEX, 2017). Worldwide carbon pricing initiatives use internal prices between 1 and 140 € t<sup>-1</sup> C (Zechter et al., 2016), the social cost of CO<sub>2</sub> was estimated to range between 5 and 65 \$ t<sup>-1</sup> (around 5–55 € t<sup>-1</sup> CO<sub>2</sub>, Greenstone et al., 2013), and the UN recommend a minimum of 100 \$ t<sup>-1</sup> C (approximately 85 € t<sup>-1</sup> C) to maintain global warming within the 1.5–2-degree Celsius pathway (United Nations Global Compact, 2016).

### 2.4.3 Nutrient loss

The environmental costs associated with the dis-services of nitrate losses into groundwater are summarized by Brink et al. (2011) and range from 0 to 4 € kg<sup>-1</sup> N. Recent studies from Denmark and United Kingdom used values of 8 € and 8.4 € kg<sup>-1</sup> N respectively (OXERA,

2006; Jacobsen, 2017).

#### 2.4.4 Soil loss

Soil is an important component of agricultural production. Its degradation can lead to a loss of productivity and cause additional off-site (external) costs for compensation and reparation. For the UK, [OXERA \(2006\)](#) used the value of  $6.41 \text{ € t}^{-1}$  that it costs to remove sediment from domestic water supplies. [Schwegler \(2014\)](#) found that the environmental cost of this dis-service was between 0.9 and  $23 \text{ € t}^{-1}$ .

#### 2.4.5 Pollination deficits

The dis-service assumed here is assessed in those parts of the LTS where pollination services are deficient. In these areas, crop yield was reduced by the specified requirement for pollination. For example, cherry production is 65% dependent on pollination ([Gallai et al., 2009](#)); in pollination deficit areas, the cherry yield was thus assumed to decline up to 65%. For each crop within pollination deficit areas, the biophysical demand for pollination, based on [Gallai et al. \(2009\)](#), was multiplied by the biomass benefit.

### 2.5. Summary of the net ecosystem service value

Eq. (2) describes the benefits and costs associated with the modelled ES. The value (V) of the indicator (I) for the benefit or cost of a particular ES is the product of the annual quantity of that indicator (Q) multiplied by the monetary value calculated for one unit of that indicator. [Table 2](#) shows the price range and the monetary value (MV<sub>I</sub>) of each assessed indicator.

$$V_I = Q_I * MV_I \quad (2)$$

In the final step of the analysis, the services (S) and dis-services (D) were aggregated to provide a net economic value of the combined impact of the ES (NET ES<sub>value</sub>) by applying Eq. (3).

$$NETES_{value} = S_{Biomass} + S_{Water} + S_{Carbon} - D_{Nutrient} - D_{Soil} - D_{Pollination} \quad (3)$$

with the benefits of biomass production service ( $S_{Biomass}$ ), groundwater ( $S_{Water}$ ), carbon storage ( $S_{Carbon}$ ), and the costs for dis-services nutrient loss ( $D_{Nutrient}$ ), soil loss ( $D_{Soil}$ ) and yield losses caused by reduced pollination ( $D_{Pollination}$ ). The result was expressed for each LTS [Units:  $\text{€ ha}^{-1} \text{ a}^{-1}$ ]. [Fig. 2](#) shows an example of the Greek case study region (GR) with four AF (AF1, AF2, etc.) and four NAF LTS (NAF1, NAF2, etc.). The biogeographical comparison was done for the Atlantic, Continental, and Mediterranean regions. Detailed results for each case study region can be found in the [Research data](#).

### 2.6. Evaluation of threshold prices

In order to identify a threshold cost for each pollutant where the

benefit of the non-agroforestry landscape (NAF) matched the agroforestry (AF) landscape for nutrient emissions, soil losses, and carbon storage, we conducted a detailed analysis of the range of prices found in literature. The intersection points – where landscapes with and without agroforestry systems (AF vs NAF LTS) are on equal economic terms – were determined.

Nutrient loss expressed as nitrate pollution costs were examined in the range between 0 and  $8 \text{ € kg}^{-1} \text{ N}$ , soil degradation costs were examined from 0 to  $20 \text{ € t}^{-1}$  soil and carbon prices were assessed in a range between 0 and  $100 \text{ € t}^{-1} \text{ C}$ .

The analyses were conducted using [R Development Core Team \(2016\)](#). The figures were created with the R packages [ggplot2 \(Wickham et al., 2016\)](#) and [plotly \(Sievert et al., 2016\)](#) and the maps with [QGIS \(QGIS Development Team, 2015\)](#).

## 3. Results

### 3.1. Valuation of ecosystem services

#### 3.1.1. Net benefit from biomass production

The mean value for the annual net financial benefit of biomass production (crop and timber products) tended to be higher in agricultural NAF landscapes. On average across all study regions, the mean profit was  $36 \text{ € ha}^{-1} \text{ a}^{-1}$  in the NAF landscapes as compared to  $29 \text{ € ha}^{-1} \text{ a}^{-1}$  in the AF landscapes ([Fig. 3](#)). Large differences were found among the biogeographical regions. The oak and olive systems of the Mediterranean landscapes had a mean financial net benefit of  $76 \text{ € ha}^{-1} \text{ a}^{-1}$ , and the AF landscapes provided a greater financial revenue from biomass than the NAF landscapes. Atlantic and Continental landscapes were less lucrative, and NAF LTS generated slightly greater financial net benefits than the AF landscapes ([Fig. 3](#)).

#### 3.1.2. Monetary valuation of individual ecosystem services and dis-services


In terms of benefits, the market value of biomass production was greater than the monetary values assigned to groundwater, carbon storage, nutrient and soil losses, and pollination service deficits across all the LTS, reaching as much as  $160 \text{ € ha}^{-1} \text{ a}^{-1}$  in some cases ([Fig. 4](#)). The financial benefit of groundwater recharge was typically less than  $2 \text{ € ha}^{-1} \text{ a}^{-1}$ . Carbon sequestration benefits ranged between 15 and  $30 \text{ € ha}^{-1} \text{ a}^{-1}$ . In terms of costs, nutrient pollution in water caused costs as great as  $150 \text{ € ha}^{-1} \text{ a}^{-1}$  and soil loss costs ranged between 15 and  $30 \text{ € ha}^{-1} \text{ a}^{-1}$ . The market value of reduced pollination service was typically minimal across the LTS.

[Fig. 4](#) also illustrates the relative performance in monetary terms depending on the proportion of agroforestry in a LTS. Whilst the dis-service nutrient loss was higher in LTS without agroforestry, only a slight difference appeared in the case of the market value of biomass production. The highest values, both positive and negative, occurred in LTS without agroforestry.

**Table 2**  
Summary of prices-ranges for ecosystem service indicators and the used monetary values.

Indicators	Unit	Price range	References	Used monetary value (MV <sub>I</sub> )
Services	Biomass production	$\text{€ t}^{-1}$ 0.43–802.6 depending on crop and country	FADN (2017), FAO (2017b), UNECE/FAO (2017)	0.43–802.6 depending on crop and country
	Groundwater recharge	$\text{€ m}^{-3}$ 0.0–4.0 depending on country	JRC Water Portal (2017); Roo et al. (2012)	0.0–4.0 depending on country
	Carbon storage	$\text{€ t C}^{-1}$ 1.0–140.0	European Energy Exchange (EEX) (2017), Zechter et al. (2016)	5
Dis-Services	Nutrient loss	$\text{€ kg N}^{-1}$ 0.0–8.4	García de Jalón et al. (2018b), Jacobsen (2017), OXERA (2006)	4
	Soil loss	$\text{€ t}^{-1}$ 0.9–23.0	García de Jalón et al. (2018b), Schwegler (2014)	6.41
	Pollination deficits	$\text{€ t}^{-1}$ 0.43–802.6 depending on crop and country	FADN (2017), FAO (2017b), Gallai et al. (2009), UNECE/FAO (2017)	0.43–802.6 depending on crop and country






**Fig. 2.** Visualisation of net ecosystem services value (NET ES<sub>value</sub>) composition including service and dis-service indicators of biomass production, ground-water, carbon storage, nutrient loss, soil loss, and pollination deficit. Indicators were assessed in each landscape test site (LTS) and summarized to NET ES<sub>value</sub> (black cross). The figure shows an example of the Greek case study region with four agroforestry (AF1, AF2, etc.) and four non-agroforestry LTS (NAF1, NAF2, etc.) as Business-As-Usual baseline.

### 3.1.3. Integrated assessment of monetary valuation of all ecosystem services and dis-services


The net value of the ES for each LTS was also summed up for the case study regions (Fig. 5). The net value of the AF landscapes tended to be greater in all three biogeographical regions, indicating that they provided greater economic welfare to society in comparison to the NAF landscapes. However, in nearly all regions, the net societal values of both, the agricultural and the agroforestry landscapes, were calculated to be negative when externalities were included in the economic analysis. The only exception were the Mediterranean agroforestry landscapes. The highest negative values were found in agricultural landscapes in the Atlantic regions. All results per ES and case study regions are listed in [Research data](#).

### 3.2. Threshold prices


Building on the previous results, threshold values were calculated to identify the ES price level that would be needed for AF systems to become as profitable as the NAF systems (Fig. 6). This was done for nutrient loss, soil loss, and carbon storage using the revenue from biomass production to provide a baseline for the NAF LTS, whilst the external



**Fig. 4.** Monetary values [€ ha<sup>-1</sup> a<sup>-1</sup>] of ES indicators, depending on the percentage of agroforestry in the landscape test sites (LTS). The coloured lines are the regression line of the measurements.



**Fig. 3.** Average net financial benefit of biomass production [€ ha<sup>-1</sup> a<sup>-1</sup>] of all 11 cases study regions (I) and divided into biogeographical regions (II) based on landscape test sites [LTS] grouped by land cover categories into agroforestry (AF) and non-agroforestry (NAF, Business as Usual) sites.



**Fig. 5.** Net ecosystem service value in € ha<sup>-1</sup> a<sup>-1</sup> of all 11 cases study regions (I) and divided into biogeographical regions (II) based on landscape test sites [LTS] grouped according to dominating land cover categories into agroforestry (AF) and non-agroforestry (NAF, Business as Usual) LTS.

costs or benefits for each ES were added individually to the baseline of the AF LTS. In this analysis, cost and prices of the other ES were not accounted for.

### 3.2.1. Nutrient loss

Fig. 6a shows how the economic performance (€ ha<sup>-1</sup> a<sup>-1</sup>) in the biogeographic regions decreased as the cost of nutrient losses (€ kg<sup>-1</sup> N) increased. As the economic output of biomass was used as baseline it remained unchanged. The AF landscapes generally showed a slower decrease in overall profitability as costs of nutrient losses increased, which indicates an overall greater robustness of these systems. The NAF landscapes showed negative economic outcomes at a nutrient emission cost of 3 € kg<sup>-1</sup> N, whereas AF LTS provided positive returns up to a nutrient emission cost of approximately 5 € kg<sup>-1</sup> N.

These results differed in the three biogeographical regions. Whilst the AF LTS in the Atlantic and Continental systems were slightly less profitable than NAF when nutrient emission costs were 0 € kg<sup>-1</sup> N, AF and NAF were equally profitable when the nutrient emission cost was 2.5 € kg<sup>-1</sup> N. This shows that even though economic output of biomass production is generally lower in Atlantic and Continental AF (Atlantic: AF 32.3 € ha<sup>-1</sup> a<sup>-1</sup>, NAF 42.7 € ha<sup>-1</sup> a<sup>-1</sup>; Continental: AF 26.0 € ha<sup>-1</sup> a<sup>-1</sup>, NAF 34.7 € ha<sup>-1</sup> a<sup>-1</sup>), introducing even fairly low costings for nutrient emission would reverse the relationship due to lower nitrate losses in the AF areas. In all three regions, the relative benefit of AF systems increased as the cost of nutrient emission increased.

### 3.2.2. Soil loss

The soil loss assessment (Fig. 6b) showed similar results to the nutrient emission assessment. In general, a rise in the cost of soil erosion resulted in declining economic performance of both AF and NAF relative to the economic output of the biomass only scenario. Again, the economic performance of the AF landscapes suggested greater robustness as decreases in economic performance were less than for NAF as the cost of soil losses increased. While in Atlantic and Continental regions, economic performance of AF was lower at low soil loss costs compared to NAF, the economic performance of AF benefitted from rising costs of soil loss relative to NAF. At values for soil loss of 12 € t<sup>-1</sup> soil (Continental biogeographic region) and 17 € t<sup>-1</sup> soil (Atlantic biogeographic region), AF and NAF landscapes produced the same economic outcome. Raising the cost for soil loss by another 5–10 € made

all landscapes (AF, NAF) unprofitable in those two regions, whilst in the Mediterranean region, both landscape types remained profitable, at least within the price range investigated.

### 3.2.3. Carbon sequestration


The results for carbon sequestration (Fig. 6c) showed that increasing the value of stored carbon resulted in increases in the economic performance of both AF and NAF systems across all the biogeographic regions. However, the patterns were comparable to the results for nutrient emissions and soil loss. Generally, AF was more profitable than NAF even at modest carbon prices. In Atlantic and Continental biogeographic regions particularly, AF profited from an increasing carbon value and exceeded the economic performance of NAF at most carbon values (thresholds were at approximately 10 € t<sup>-1</sup> C in the Continental biogeographic regions and 30 € t<sup>-1</sup> C in the Atlantic biogeographic region; the Mediterranean AF was more profitable at all carbon values).

## 4. Discussion

This research investigated three questions: 1) How does the societal value of agroforestry landscapes compare with landscapes dominated by agriculture in different parts of Europe if only the values of the products are considered? 2) Do these results change if the values of selected regulating services are included? and 3) How sensitive are the results to changes in ES prices?

The trends we identified are in line with former findings on ES and mitigation provided by agroforestry systems (e.g. Jose, 2009; Moreno et al., 2018; Tsonkova et al., 2012). Biophysical and economic differences between agroforestry and non-agroforestry practices are clearer at the plot scale (Graves et al., 2007; Palma et al., 2007; Sereke et al., 2015). Our investigation related to the landscape scale because some ES such as soil conservation or pollination services involve spatial interactions that cannot be evaluated at the plot scale. Yet, as we investigated mixed landscapes there were some agroforestry trees even in NAF LTS and vice-versa, which somehow “blurred” the differences between the landscape test sites. Also, the proportion of the land use categories agroforestry, agriculture, forest and others differed from region to region, which led to the high variability observed.

In response to the first research question, in Atlantic and Continental regions of Europe, the market values of the products from agroforestry landscapes were calculated to be generally lower than for



**Fig. 6.** Economic performance of agroforestry (AF) and non-agroforestry (NAF, Business as Usual) for different ecosystem services (a) nutrient emission costs, (b) soil loss costs and (c) carbon prices together with the current sales revenues of biomass production in  $\text{€ ha}^{-1} \text{a}^{-1}$  (I) over all 11 cases study regions and (II) divided into biogeographical regions based on landscape test sites [LTS] grouped by dominating land cover categories into AF and NAF (Business as Usual) LTS.

non-agroforestry systems. The opposite was observed in Mediterranean regions, where the market value of the products from agroforestry landscapes were calculated to be higher than for the non-agroforestry cases. This was mainly due to the multiple tree products (olives or acorns in addition to timber) and the use of the FADN index in the calculation of the market value of the biomass production, which was between 1.28 and 1.31 in Portugal, Spain and Greece; while in northern and central European countries values around 1.0 were obtained. The agroforestry olive groves in our Greek case study region were already fully productive and therefore profitable (producer price:  $\sim 2000 \text{ € t}^{-1}$ ; net benefit:  $\sim 470 \text{ € t}^{-1}$ ; yield:  $100 \text{ kg tree}^{-1}$ ;  $1 \text{ t ha}^{-1} \text{ a}^{-1}$  olives,  $0.2 \text{ t ha}^{-1} \text{ a}^{-1}$  olive oil; European Commission, 2012; FAO, 2017b; Pantera et al., 2016). According to the European Commission, 2012 (intensive) olive production is one of the most important and profitable agricultural activities in southern marginal regions. Whilst olive groves produce in average  $2.5 \text{ t ha}^{-1}$ , agroforestry production is around  $1 \text{ t ha}^{-1}$ . After five to seven years, olive systems start to become fully productive and after around year 20 the initial costs are covered and they obtain revenues (Stillitano et al., 2016). This resulted in AF landscapes to have higher sales revenues in Mediterranean regions than NAF landscapes. The multiple gains of dehesas are reflected in their land prices for lease or sale. While open pastures in Spain cost around  $5000 \text{ € ha}^{-1}$  and are leased for  $53.50 \text{ € ha}^{-1}$ , dehesas are on sale for  $\sim 8000 \text{ € ha}^{-1}$  and leased for  $78.70 \text{ € ha}^{-1}$  (Consejería de agricultura, 2014; FEDEHESA, 2017). This positive economic performance for AF relative to NAF is also reflected in the spread and extent of agroforestry in Mediterranean regions. Den Herder et al. (2017) identified the current extent of agroforestry in Europe and found that the largest areas were in Spain (5.6 million ha), Greece (1.6 million ha), France (1.6 million ha), Italy (1.4 million ha) and Portugal (1.2 million ha).

For agroforestry in Continental and Atlantic regions the situation is different. Sereke et al. (2015), Nerlich et al. (2013) and Eichhorn et al. (2006) have stated that many traditional agroforestry systems are in decline. Highlighting and valuing their environmental role was related to the second research question. Actually, the decision of managing the land as an agroforestry system is not only related to financial profitability but also to other criteria such as to increase the diversification of products, improve biodiversity, animal health and welfare as described by García de Jalón et al., 2018a, Rois-Díaz et al., 2018, and Sereke et al. (2016). This indicates that (some) farmers value ES even if they don't provide financial benefit. At the policy level, the European environmental (e.g. Water Framework Directive) and agricultural policies (CAP with greening and cross compliance) focus was on the impact of environmental pollution, notably nutrient emissions and soil losses. Here, even small monetary benefits associated with reduced nutrient and soil losses, and – in addition – modest carbon sequestration payments favoured the economic performance of the assessed systems in favour of agroforestry. These findings are echoed by Zander et al. (2016) in their evaluation of the performance of grain legumes, and La Notte et al. (2017) in their evaluation of in-stream nitrogen and reflect the failure of markets to pass costs back to polluters.

The third research question focused on the sensitivity of the outcomes to price changes. Unexpectedly, the value of nutrient emissions was the most important factor affecting the economic performance of the assessed systems, since small changes in prices charged for nutrient losses led to relatively large changes in economic performance. Compared to this, soil losses were of lesser importance, as also observed by García de Jalón et al. (2018b). Even though water pollution by nitrates is addressed by several environmental regulations (e.g. Nitrate Directive, Water Framework Directive), European water prices for irrigation or domestic purposes are surprisingly low. In comparison with the costs and prices assigned to other ES indicators, they thus had a negligible impact on economic performance.

The decline in pollinators and its possible consequence on pollination service has been a key issue at European scale (Zulian et al., 2013; Breeze et al., 2014). However, as enough nesting and foraging resource

for wild pollinators were available in all case study landscapes (Kay et al., 2018b), the cost of potentially reduced pollination services had no impact.

Regarding the European climate policy (e.g. EU 2030 Climate and Energy Framework), carbon storage and emission reduction are the most important ES. Agroforestry has the potential to store carbon on agricultural land (Zomer et al., 2016). The United Nations Global Compact (2016) proposes the use of a carbon value of  $\$100 \text{ t}^{-1}$  (approximately  $85 \text{ € t}^{-1} \text{ C}$ ). If such high carbon prices could be obtained by farmers, this would drastically change the economic performance of many land use systems. Even with a carbon price of  $30 \text{ € t}^{-1} \text{ C}$ , landscapes with AF were more profitable compared to NAF LTS.

## 5. Conclusion

In many parts of Europe, agroforestry systems such as wood pastures and hedgerows remain under threat either due to land abandonment or an increase in mechanization and decline in labour availability. In this study, agroforestry landscapes in Atlantic and Continental regions showed slightly lower market outputs than agricultural areas if the focus was only on marketable provisioning ecosystem services. However, in Mediterranean regions, the marketable outputs from the considered agroforestry systems were typically greater than the associated agricultural system.

When the societal values of regulating ES and dis-services were also accounted for, the aggregated landscape profitability of AF was generally higher than NAF in each region. This was driven by a reduction in societal costs related to lower nutrient and soil losses, and the societal benefits of carbon sequestration. Overall, our study underlined that relatively low costs per ES unit (nutrient emission:  $> 2.5 \text{ € kg}^{-1} \text{ N}$ ; soil loss:  $> 17 \text{ € t}^{-1}$  soil; carbon sequestration  $> 30 \text{ € t}^{-1} \text{ C}$ ) would be sufficient to render AF profitable, at least to match NAF profitability.

Our results show that there is a critical gap in economic assessments that fails to account for ecological and social benefits. This issue needs to be imperatively addressed if international agreements (e.g. European Commission, 2011; UNFCCC, 2015; United Nations, 1992) should have any effect. New methods of accounting for externalities e.g. payments for ecosystem services or other incentives to stimulate farmers and land users to turn towards more socially beneficial forms of land use should be strengthened.

## Acknowledgements

We acknowledge funding through Grant 613520 from the European Commission (Project AGFORWARD, 7th Framework Program), the Forest Research Centre strategic project (PEst OE/AGR/UI0239/2014) and the Portuguese Foundation for Science and Technology through the contract SFRH/BD/52691/2014. We would like to thank three anonymous reviewers for their constructive comments, which helped to improve an earlier version of the manuscript.

## Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ecoser.2019.100896>.

## References

- Allen, R.G., Pereira, L.S., Raes, D., et al., 1998. Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. Rom.
- Bennett, D.E., Gosnell, H., Lurie, S., Duncan, S., 2014. Utility engagement with payments for watershed services in the United States. *Ecosyst. Serv.* 8, 56–64. <https://doi.org/10.1016/J.ECOSER.2014.02.001>.
- Boyce, J.K., 2018. Carbon pricing: effectiveness and equity. *Ecol. Econ.* 150, 52–61.
- Breeze, T.D., Vaissière, B.E., Bommarco, R., et al., 2014. Agricultural policies exacerbate honeybee pollination service supply-demand mismatches across Europe. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0082996>.
- Brenner, J., Jiménez, J.A., Sardá, R., Garola, A., 2010. An assessment of the non-market



- value of the ecosystem services provided by the Catalan coastal zone, Spain. *Ocean Coast. Manage.* 53, 27–38. <https://doi.org/10.1016/j.ocecoaman.2009.10.008>.
- Brink, C., van Grinsven, H., Jacobsen, B.H., et al., 2011. Costs and benefits of nitrogen in the environment. In: *The European Nitrogen Assessment*. Cambridge University Press, Cambridge, pp. 513–540.
- Caparrós, A., Cerdá, E., Ovando, P., Campos, P., 2007. Sequestration with Reforestations and Biodiversity – Scenic Values.
- Consejería de agricultura pesca y desarrollo rural, 2014. Encuesta cánones de arrendamiento rústico. <http://www.juntadeandalucia.es/organismos/agriculturapescaydesarrollorural/consejeria/sobre-consejeria/estadisticas/paginas/agrarias-precios-encuesta-canon.html>.
- Costanza, R., Arge, R., De Groot, R., et al., 1997. The value of the world's ecosystem services and natural capital. *Nature* 387, 253–260. <https://doi.org/10.1038/387253a0>.
- Costanza, R., de Groot, R., Braat, L., et al., 2017. Twenty years of ecosystem services: how far have we come and how far do we still need to go? *Ecosyst. Serv.* 28, 1–16.
- Council of the European Union, 2017. Outcome of the council meeting “Agriculture and Fisheries”, Brussels, 17 and 18 July 2017.
- Roo, A. De, Burek, P., Gentile, A., et al., 2012. A multi-criteria optimisation of scenarios for the protection of water resources in Europe. Support to the EU Blueprint to Safeguard Europe's Waters.
- den Herder, M., Moreno, G., Mosquera-Losada, R.M., et al., 2017. Current extent and stratification of agroforestry in the European Union. *Agric. Ecosyst. Environ.* 241, 121–132. <https://doi.org/10.1016/j.agee.2017.03.005>.
- European Energy Exchange (EEX), 2017. European Emission Allowances Auction (EUA) Global Environmental Exchange. In: Homepage. <https://www.eex.com/de/marktdaten/umweltprodukte/a>.
- Eichhorn, M.P., Paris, P., Herzog, F., et al., 2006. Silvoarable systems in Europe – past, present and future prospects. *Agrofor. Syst.* 67, 29–50. <https://doi.org/10.1007/s10457-005-1111-7>.
- European Commission, 2011. The EU Biodiversity Strategy to 2020.
- European Commission, 2012. Economic analysis of the olive sector. *Dir. Agric. Rural Dev.* 10 p.
- European Commission, 2013 Regulation (EU) No 1305/2013 on support for rural development by the European Agricultural Fund for Rural Development (EAFRD) and repealing Council Regulation (EC) No 1698/2005. *Off J Eur Union* 1698:487–548.
- European Commission, 2016. Agriculture – a partnership between Europe and farmers. The EU's common agricultural policy (CAP): for our food, for our countryside, for our environment.
- FADN, 2017. FADN Public Database – Total output/total input (SE132). In: dataset. [http://ec.europa.eu/agriculture/rica/database/report\\_en.cfm?dwh=50](http://ec.europa.eu/agriculture/rica/database/report_en.cfm?dwh=50).
- FAO, 2017a. Producer Price – Annual. In: Dataset. <http://www.fao.org/faostat/en/#data/PP>.
- FAO, 2017b. FAO Statistic – CROPS. In: Dataset. <http://www.fao.org/faostat/en/#data/QC>.
- FEDEHESA, 2017. Land prices. <http://fedehesa.org/>.
- Fu, B.J., Su, C.H., Wei, Y.P., et al., 2011. Double counting in ecosystem services valuation: causes and countermeasures. *Ecol. Res.* 26, 1–14. <https://doi.org/10.1007/s11284-010-0766-3>.
- Gallai, N., Salles, J.M., Settele, J., Vaissière, B.E., 2009. Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecol. Econ.* <https://doi.org/10.1016/j.ecolecon.2008.06.014>.
- García de Jalón, S., Graves, A., Palma, J.H., et al., 2018b. Modelling and valuing the environmental impacts of arable, forestry and agroforestry systems: a case study. *Agrofor. Syst.* 92, 1059–1073. <https://doi.org/10.1007/s10457-017-0128-z>.
- García de Jalón, S., Burgess, P.J., Graves, A., et al., 2018a. How is agroforestry perceived in Europe? An assessment of positive and negative aspects by stakeholders. *Agrofor. Syst.* 92, 829–848. <https://doi.org/10.1007/s10457-017-0116-3>.
- Gómez-Baggethun, E., de Groot, R., Lomas, P.L., Montes, C., 2010. The history of ecosystem services in economic theory and practice: from early notions to markets and payment schemes. *Ecol. Econ.* 69, 1209–1218. <https://doi.org/10.1016/j.ecolecon.2009.11.007>.
- Graves, A., Burgess, P.J., Palma, J.H.N., et al., 2007. Development and application of bio-economic modelling to compare silvoarable, arable, and forestry systems in three European countries. *Ecol. Eng.* 29, 434–449. <https://doi.org/10.1016/j.ecoleng.2006.09.018>.
- Greenstone, M., Kopits, E., Wolverton, A., 2013. Developing a social cost of carbon for us regulatory analysis: a methodology and interpretation. *Rev. Environ. Econ. Policy* 7, 23–46. <https://doi.org/10.1093/reep/res015>.
- Haines-Young, R., Potschin, M., 2013. Common International Classification of Ecosystem Services (CICES): Consultation on Version 4, August–December 2012. EEA Framework Contract No EEA/IEA/09/003 34 p.
- Hill, B., Bradley, B.D., 2015. Comparison of farmers' incomes in the EU member states. *Hürdler, J., Prasuhn, V., Spiess, E., 2015. Abschätzung diffuser Stickstoff- und Phosphoreinträge in die Gewässer der Schweiz.* 117 p.
- Jacobsen, B.H., 2017. Beregning af kvælstofsygepris med udgangspunkt i Fødevarer- og Landbrugspakken. IFRO Udrødning 08, 27 p.
- Jose, S., 2009. Agroforestry for ecosystem services and environmental benefits: an overview. *Agrofor. Syst.* 76, 1–10. <https://doi.org/10.1007/s10457-009-9229-7>.
- JRC Water Portal, 2017. Map of costs of water abstraction for irrigation. In: Dataset. <http://water.jrc.ec.europa.eu/waterportal>.
- Kay, S., Crous-Duran, J., García de Jalón, S., et al., 2018b. Spatial similarities between European agroforestry systems and ecosystem services at the landscape scale. *Agrofor. Syst.* 92, 1075–1089. <https://doi.org/10.1007/s10457-017-0132-3>.
- Kay, S., Crous-Duran, J., García de Jalón, S., et al., 2018a. Landscape-scale modelling of agroforestry ecosystem services in Swiss orchards: a methodological approach. *Landscape Ecol.* 33, 1633–1644. <https://doi.org/10.1007/s10980-018-0691-3>.
- La Notte, A., Maes, J., Dalmazzone, S., et al., 2017. Physical and monetary ecosystem service accounts for Europe: a case study for in-stream nitrogen retention. *Ecosyst. Serv.* 23, 18–29. <https://doi.org/10.1016/j.ecoser.2016.11.002>.
- Liski, J., Palosuo, T., Peltoniemi, M., Sievänen, R., 2005. Carbon and decomposition model Yasso for forest soils. *Ecol. Model.* 189, 168–182. <https://doi.org/10.1016/j.ecolmodel.2005.03.005>.
- Lonsdorf, E., Kremen, C., Ricketts, T., et al., 2009. Modelling pollination services across agricultural landscapes. *Ann. Bot.* 103, 1589–1600. <https://doi.org/10.1093/aob/mcp069>.
- MEA, 2003. Ecosystems and human well-being: a framework for assessment. In: *Millennium Ecosystem Assessment*. Millenium Ecosystem Assessment. Island Press, Washington, DC, pp. 1–25.
- Moreno, G., Aviron, S., Berg, S., et al., 2018. Agroforestry systems of high nature and cultural value in Europe: provision of commercial goods and other ecosystem services. *Agrofor. Syst.* 92, 877–891. <https://doi.org/10.1007/s10457-017-0126-1>.
- Moreno, G., Aviron, S., Burgess, P., et al., 2017. AGFORWARD EU Project Milestone 33 (7.3): Spatial characterization of sample landscapes.
- Muradian, R., Corbera, E., Pascual, U., et al., 2010. Reconciling theory and practice: an alternative conceptual framework for understanding payments for environmental services. *Ecol. Econ.* 69, 1202–1208. <https://doi.org/10.1016/j.ecolecon.2009.11.006>.
- Nerlich, K., Graeff-Hönniger, S., Claupein, W., 2013. Agroforestry in Europe: a review of the disappearance of traditional systems and development of modern agroforestry practices, with emphasis on experiences in Germany. *Agrofor. Syst.* 87, 1211. <https://doi.org/10.1007/s10457-013-9618-9>.
- OXERA, 2006. What is the Cost of Reducing Ammonia, Nitrates and BOD in Sewage Treatment Works Effluent? OFWAT, pp. 36.
- Palma, J., Graves, A., Crous-Duran, J., et al., 2018. EcoYield-SAFE.
- Palma, J., Graves, A., Burgess, P., et al., 2007. Integrating environmental and economic performance to assess modern silvoarable agroforestry in Europe. *Ecol. Econ.* 63, 759–767. <https://doi.org/10.1016/j.ecolecon.2007.01.011>.
- Pantera, A., Papadopoulos, A., Kitsikopoulos, D., et al., 2016. System report: olive agroforestry in Molos, Central Greece. AGFORWARD Deliv. 3 (7), 9.
- Pe'er, G., Lakner, S., Müller, R., et al., 2017. Is the CAP Fit for purpose? An evidence-based fitness-check assessment – An evidence based fitness-check assessment. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig.
- Ponisio, L.C., M'Gonigle, L.K., Mace, K.C., et al., 2014. Diversification practices reduce organic to conventional yield gap. *Proc. R. Soc. B Biol. Sci.* <https://doi.org/10.1098/rspb.2014.1396>.
- QGIS Development Team. 2015. QGIS Geographic Information System. Open Source Geospatial Found. Proj.
- R Development Core Team, 2016. R Software. R: A language and environment for statistical computing.
- Renard, K., Foster, G., Weesies, G., et al., 1997. Predicting soil erosion by water: a guide to conservation planning with the Revised Universal Soil Loss Equation (RUSLE). *Agricultural Handbook No. 703*. U.S. Department 1997 of Agriculture (USDA).
- Rois-Díaz, M., Lovric, N., Lovric, M., et al., 2018. Farmers' reasoning behind the uptake of agroforestry practices: evidence from multiple case-studies across Europe. *Agrofor. Syst.* 92, 811–828. <https://doi.org/10.1007/s10457-017-0139-9>.
- Schwegler, P., 2014. Economic valuation of environmental costs of soil erosion and the loss of biodiversity and ecosystem services caused by food wastage. *Sci. Pap. Award* 2014 20 p.
- Sereke, F., Graves, A.R., Dux, D., et al., 2015. Innovative agroecosystem goods and services: key profitability drivers in Swiss agroforestry. *Agron Sustain Dev* 35, 759–770. <https://doi.org/10.1007/s13593-014-0261-2>.
- Sereke, F., Dobricki, M., Wilkes, J., et al., 2016. Swiss farmers don't adopt agroforestry because they fear for their reputation. *Agrofor. Syst.* 90, 385–394. <https://doi.org/10.1007/s10457-015-9861-3>.
- Sievert, C., Farmer, C., Hocking, T., et al., 2016. plotly: Create Interactive Web Graphics via “plotly.js”.
- Somarriba, E., 1992. Revisiting the past: an essay on agroforestry definition. *Agrofor. Syst.* 19, 233–240. <https://doi.org/10.1007/BF00118781>.
- Stillitano, T., De Luca, A.I., Falcone, G., et al., 2016. Economic profitability assessment of mediterranean olive growing systems. *Bulg J Agric Sci* 22, 517–526.
- TEEB, 2010. The Economics of Ecosystems and Biodiversity (TEEB). In: Kumar, P. (Ed.), *Ecological and Economic Foundations*. Earthscan, UNEP/Earthprint, London.
- Tilman, D., Cassman, K.G., Matson, P.A., et al., 2002. Agricultural sustainability and intensive production practices. *Nature*. <https://doi.org/10.1038/nature01014>.
- Toralba, M., Fagerholm, N., Burgess, P.J., et al., 2016. Do European agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis. *Agric. Ecosyst. Environ.* 230, 150–161. <https://doi.org/10.1016/j.agee.2016.06.002>.
- Tsonkova, P., Böhm, C., Quinkenstein, A., Freese, D., 2012. Ecological benefits provided by alley cropping systems for production of woody biomass in the temperate region: a review. *Agrofor. Syst.* 85, 133–152. <https://doi.org/10.1007/s10457-012-9494-8>.
- UNECE/FAO, 2017. UNECE/FAO TIMBER database - Historical Prices, Export Unit Price. In: Dataset. <http://www.unece.org/forests/output/prices.html>.
- UNFCCC (2015) Paris Agreement.
- United Nations, 1992. United Nations Framework Convention on Climate Change. United Nations 1:270–277. <https://doi.org/10.1111/j.1467-9388.1992.tb00046.x>.
- United Nations Global Compact, 2016. UN Global Compact Calls on Companies to Set \$100 Minimum Internal Price on Carbon. In: April 22, 2016. <https://www.unglobalcompact.org/news/3381-04-22-2016>.
- Wezel, A., Casagrande, M., Celette, F., et al., 2014. Agroecological practices for sustainable agriculture. A review. *Agron. Sustain. Dev.* 34, 1–20. <https://doi.org/10.1007/>

- s13593-013-0180-7.
- Wickham, H., Winston, C., RStudio, 2016. *ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics*. Cran.
- Zander, P., Amjath-Babu, T.S., Preissel, S., et al., 2016. Grain legume decline and potential recovery in European agriculture: a review. *Agron. Sustain. Dev.* <https://doi.org/10.1007/s13593-016-0365-y>.
- Zechter, R., Kerr, T.M., Kossoy, A., et al., 2016. Carbon Pricing Watch 2016.
- Zhang, W., Ricketts, T.H., Kremen, C., et al., 2007. Ecosystem services and dis-services to agriculture. *Ecol. Econ.* 64, 253–260. <https://doi.org/10.1016/j.ecolecon.2007.02.024>.
- Zomer, R.J., Neufeldt, H., Xu, J., et al., 2016. Global tree cover and biomass carbon on agricultural land: the contribution of agroforestry to global and national carbon budgets. *Sci. Rep.* 6, 29987. <https://doi.org/10.1038/srep29987>.
- Zulian, G., Maes, J., Paracchini, M., 2013. Linking land cover data and crop yields for mapping and assessment of pollination services in Europe. *Land* 2, 472–492. <https://doi.org/10.3390/land2030472>.