

AMENDMENT NO. 1 JULY 2025
TO
IS 18947 : 2024 GLOBULES FOR USE IN HOMOEOPATHY — SPECIFICATION

(Foreword, para 7) — Substitute the following for the existing:

‘The composition of the Committee responsible for the formulation of this standard is given in Annex C.’

(Page 1, clause 2) — Insert the following new references at the end:

<i>IS No.</i>	<i>Title</i>
IS 5887 (Part 3/Sec 1) : 2020/ISO 6579-1 : 2017	Methods for detection of bacteria responsible for food poisoning: Part 3 Horizontal method for the detection, enumeration and serotyping of <i>Salmonella</i> , Section 1 Detection of <i>Salmonella</i> spp. (<i>third revision</i>)
IS 15279 : 2003	Sugar and sugar products — Methods of test
IS 18946 : 2024	Non-medicated tablets for use in homoeopathy — Specification

(Page 1, clause 5) — Substitute ‘**OTHER REQUIREMENTS AND TEST METHODS**’ for ‘**TEST METHODS**.’

(Page 1, clause 5.1) — Substitute the following for the existing:

‘5.1 General Appearance

The globules shall be white-coloured uniform spheroids on inspecting.’

(Page 1, clause 5.2.1) — Insert the following at the end:

‘Dissolve 5 g of globules in 100 ml of distilled water at room temperature. Stir continuously and observe for complete dissolution. No residue shall remain.’

(Page 1, clause 5.2.2) — Insert the following at the end:

‘Add 5 g of globules to 100 ml of alcohol (90 percent to 95 percent) at room temperature. Stir and observe. The globules shall not dissolve.’

(Page 2, clause 5.7.1) — Substitute the following for the existing:

‘5.7.1 Sucrose Content in Globules

It shall not be less than 99 percent on dried basis, when tested by either Polarimetry or HPLC method as prescribed in Annex B.’

(Page 2, clause 5.8) — Insert the following at the end:

‘Measure the optical rotation at 20 °C using the Polarimetry method when tested as per 6 of IS 15279’

(Page 3, clause 5.11) — Substitute the following for the existing:

‘5.11 Loss on Drying

The loss on drying shall not be more than 4 percent w/w, when tested as per 4 of IS 15279.’

Amendment No. 1 to IS 18947 : 2024

(Page 3, clause 5.13) — Substitute the following for the existing:

‘5.13 Foreign Matter

Globules shall be free from any flavour, foreign matter like kaolin, chalk, talc, starch, animal dander, inorganic or any synthetic whitening agents. Prepare a 5 percent w/v Solution A from powdered globules and the test methods as prescribed in Table 3 shall be followed for the testing. Globules shall comply with the requirements given in Table 3.’

(Page 3, clause 5.13, Table 3) — Substitute the following for the existing.

Table 3 Test Methods and Requirements

(Clause 5.13)

Sl No.	Test	Method
(1)	(2)	(3)
i)	Kaolin test	Take 5 ml to 10 ml of Solution A in a conical flask, add a small amount of dilute ammonia solution and allow to stand for 5 min. No white precipitate shall be observed.
ii)	Chalk test	Take 5 ml to 10 ml of Solution A, add a small amount of dilute ammonia oxalate solution and allow to stand for 5 min. No white precipitate shall be observed. Take some powdered samples and add dilute hydrochloric acid. No effervescence shall be observed.
iii)	Talc test	Take 5 ml to 10 ml of Solution A, add a small amount of ammonium carbonate solution and boil. No white precipitate shall be observed.
iv)	Starch test	Take 5 ml to 10 ml of sample Solution A, add iodine tincture solution to it, a violet to deep blue color shall not appear.
v)	Animal dander test	Place a few drops of Solution A on a clean glass slide. Use a microscope to examine the slide at varying magnifications (for example, 40x, 100x, 400x). Observe for the presence of animal dander, such as skin flakes, hair, or saliva. No animal dander shall be present. If dander is present, it will appear as small, irregular, or flattened particles, often with a fibrous structure.
vii)	Inorganic or synthetic whitening agents test	Expose a sample of Solution A to UV light (wavelength of 254 nm to 365 nm). Observation of fluorescence, typically as a bright blue or greenish glow, indicates the presence of inorganic or synthetic whitening agents (for example, optical brighteners); otherwise sample is considered free from inorganic or synthetic whitening agents.

(Page 5, clause 5.16, line 7 of para 1) — Substitute the following for the existing:

‘The acceptance criteria for microbial contamination as given in Table 4’ shall be ensured.’

(Page 5, clause **5.16**, Table 4) — Substitute the following for the existing:

Table 4 Microbial Contamination

(Clause 5.16)

Sl No.	Parameter	Requirement	Methods of Test
(1)	(2)	(3)	(4)
i)	TAMC	10 ² CFU/g, <i>Max</i>	Annex B of IS 18946
ii)	TYMC	10 ¹ CFU/g, <i>Max</i>	
iii)	Microorganism		
	a) <i>Staphylococcus aureus</i>	Absent	
	b) <i>Pseudomonas aeruginosa</i>	Absent	
	c) <i>E. coli</i>	Absent	
	d) <i>Salmonella spp.</i>	Absent	IS 5887 (Part 3/Sec 1)/ ISO 6579-1

(Page 5, clause 5.16) — Insert the following new clause at the end:

‘5.17 Sulphated Ash

Sulphated ash shall not be more than 0.1 percent, when tested as per **10** of IS 15279.’

(Page 7) — Substitute ‘ANNEX C’ for ‘ANNEX B’.

(Page 7) — Insert the following new ANNEX B after ‘ANNEX A’

‘ANNEX B

(Clause 5.7.1)

METHODS OF DETERMINING SUGAR CONTENT IN GLOBULES

B-1 POLARIMETRY METHOD

B-1.1 Equipment

- a) Analytical balance;
- b) 100 ml flask;
- c) Water R;
- d) Stirring rod/magnetic stirrer;
- e) Vacuum filtration setup;
- f) Polarimetry; and
- g) Drying oven and desiccator.

Amendment No. 1 to IS 18947 : 2024

B-1.2 Procedure

Weigh 10 g of ground globules in a 100 ml flask and make up to 100 ml with water R. Stir and decant. Filter under vacuum to obtain a clear solution. Measure the angle of optical rotation **5.8** and calculate the sucrose percentage content using the following expression:

$$\frac{10^6 \times \alpha}{66.5 \times l \times m \times (100 - H)}$$

where

α = angle of rotation;

l = length of the polarimeter tube, in decimetres;

m = exact mass of the sample, in grams;

H = loss on drying; and

Water R = Reagent grade water.

B-2 HPLC METHOD

B-2.1 Apparatus

- Liquid chromatograph — Equipped with an isocratic solvent delivery system, manual injector, refractive index detector, and recording/computing integrator;
- Column — 300 × 4 (id) mm μ -Bondapak/Carbohydrate (Waters Associates, No. 84038);
- Test solution clarification kit — Available in kit form from chromatography suppliers; 0.45 μ m filters, stable in organic solvents, are suitable; and
- Syringes — 10 μ l (No. 701-N) point style No. 1, 2 × 0.020 in. od, 25 gauge needle.

B-2.2 Reagents

- Mobile phase — LC grade acetonitrile diluted with H₂O (83 + 17). The mobile phase is degassed daily by magnetic stirring for 15 min under vacuum; and
- Sugar standard solution — Place 3.804 g fructose, 3.010 g glucose, and 0.602 g sucrose into a 100 ml volumetric flask, dissolve in 50 ml H₂O, and add CH₃CN to volume. Composition of standard approximates 5 g of honey dissolved in 50 ml aqueous CH₃CN (1 + 1).

B-2.3 Operating Conditions

Fructose, glucose, and sucrose are baseline separated and quantitated in 20 min under the following conditions: flow rate, 1.0 ml/min (3.45 Mpa; ca 500 psi); temperature, ambient (ca 23 °C); detector set so that 380 μ g fructose gives full-scale peak. Mono-, di-, and trisaccharides are eluted from the column in order of MW.

B-2.4 Preparation of Test Solution

Weigh 5 g test portion in a small beaker and transfer to a 50 ml volumetric flask with 25 ml H₂O. Immediately dilute to volume with CH₃CN and filter through 0.45 μ m filter, using a clarification kit.

B-2.5 Chromatography

Inject 10 µl of standard solution into the chromatograph. Establish retention times, measure peak heights, and check reproducibility. Repeat for the test solution. Calculate glucose, fructose, and sucrose from integrator values or peak heights as follows:

B-2.6 Calculation

$$Assay = \frac{\text{Sample peak area} \times \text{Standard weight} \times \text{Sample dilution} \times \text{Potency} \times 100}{\text{Standard peak area} \times \text{Standard dilution} \times \text{Sample weight} \times 100}$$

$$\text{Weight percent sugar} = 100 (PH/PH') (V/V') (W'/W)$$

where

PH and PH' = peak heights (or integrator values) of test solution and standard, respectively;

V and V' = (ml) test and standard (50 and 100) solutions, respectively; and

W and W' = (g) test portion (5.0) and standard, respectively.

NOTE — In case of dispute, the HPLC method shall be the referee method.