

AMENDMENT NO. 1 NOVEMBER 1979

TO

IS:1824-1978 SPECIFICATION FOR
INSECTICIDAL SPACE SPRAY

(Second Revision)

Alterations

(Page 7, clause 3.1.1) - Substitute the following for the existing clause:

'3.1.1 The packing shall also conform to the Tariff Rules as in force from time to time for packing of goods falling in the category of materials of low flash point.'

(Page 25, clause E-2.1.1 to E-2.1.3) - Substitute the following for the existing clauses:

'E-2.1.1 Weigh accurately a quantity of the material containing about 0.015 g of the lindane in a tared 250-ml erlenmeyer flask. Add 25 ml of isopropyl alcohol and shake the flask. Introduce 2.5 g of metallic sodium into the erlenmeyer flask, connect the flask to a reflux condenser and reflux for at least one hour on a water-bath. Shake the flask frequently during the refluxing period. Dissolve the excess metallic sodium by cautiously adding 10 ml of 50 percent isopropyl alcohol through the condenser at the rate of 1 to 2 drops per second. Boil for an additional 10 minutes and then add 60 ml of water.

E-2.1.2 Cool, add 2-3 drops of phenolphthalein indicator. Neutralize by adding dilute nitric acid dropwise and then add 10 ml in excess. Cool the flask to room temperature and quantitatively transfer the contents of the flask to a 250-ml separating funnel. Transfer the aqueous layer to a 500-ml erlenmeyer flask. Wash the contents of the separating funnel thrice using 20 ml water each time. Transfer the washings to the erlenmeyer flask. Reject the washed solvent. Mix well.

E-2.1.3 Add, with a pipette, 25 ml of standard silver nitrate solution. Coagulate the precipitated silver chloride on a water-bath for 30 minutes, cool the flask and, if necessary, filter the contents of the flask through a fast qualitative filter paper collecting the filtrate quantitatively in an erlenmeyer flask. Add 5 ml of ferric ammonium sulphate solution either to the cooled unfiltered mixture or to the filtrate as the case may be, and titrate the excess of silver nitrate with standard potassium thiocyanate solution (the end-point is the appearance of red ferric thiocyanate colour).'

(Page 26, clause E-3.2, formula) - Substitute the following for the existing formula:

$$\begin{array}{l} \text{Organic chlorine content} \\ \text{of the material, percent} \\ \text{by mass} \end{array} = D = \frac{3.546 \times (B - A) \times N}{M} - C$$

(AFCD 6)