

AMENDMENT NO. 6 DECEMBER 2010
TO
IS 1051 : 1980 SPECIFICATION FOR PYRETHRUM EXTRACTS

(Second Revision)

(Page 4, clause 2.2) — Substitute the following for the existing:

‘2.2 Colour and Odour — The material shall possess the characteristic odour of the commercial pyrethrum flowers. When determined by the method prescribed in Appendix D, the optical density of the material should be less than 0.900.’

(Page 17, Appendix C, clause C-4) — Add the following new Appendix after **C-4**:

‘APPENDIX D
(Clause 2.2)

DETERMINATION OF COLOUR

D-0 PRINCIPLE

The natural colouring matter content of pyrethrum extract is extracted with mineral turpentine oil. The absorbance of the solution obtained is measured by using a UV-Visible Spectrometer at a wavelength of 435 nm.

D-1 APPARATUS

D-1.1 UV-Visible Spectrophotometer suitable for measuring at wavelength of 435 nm and having a band width of 2 nm.

D-1.2 Volumetric Flask, 20 ml capacity.

D-1.3 Measuring Cylinder

D-1.4 Cuvettes

D-2 REAGENTS

D-2.1 Solvent, mineral turpentine oil of known purity.

D-3 PROCEDURE

Take 1 ml of the pyrethrum extract in the 20-ml volumetric flask. Add to it mineral turpentine oil and make up the volume up to the mark with mineral turpentine oil. Shake the flask to mix the contents thoroughly. Take a pair of clean cuvettes. Switch on the Spectrophotometer and set the wavelength at 435 nm. In one cuvette take the solvent and place this cuvette in the reference cell of the spectrophotometer. Take the diluted sample from the volumetric flask in the other cuvette and place the cuvette in the sample cell of the spectrophotometer. Close the chamber and measure the absorbance at 435 nm.’

(FAD 1)