

AMENDMENT NO. 2 MARCH 2006
TO
IS 7874 (PART 2) : 1975 METHODS OF TESTS FOR
ANIMAL FEEDS AND FEEDING STUFFS
PART 2 MINERALS AND TRACE ELEMENTS

(Page 4, clause 4.2.1) — Substitute the following for the existing text:

‘Weigh accurately about 1 g of the material and transfer to 250 ml graduated flask. Add 50 ml of the ferric sulphate solution with a pipette, gently swirling the flask add 100 ml of the ammonium hydroxide solution. Add sufficient quantity of glass distilled water to bring it to the mark. Swirl the flask enough to ensure thorough mixing but avoid vigorous shaking. Allow to settle for 10 minutes. Filter through 11-cm Whatman filter paper No. 41 or its equivalent. Transfer from the filtrate, an aliquot of 25 ml to a 250-ml beaker. Add 10 ml of nitric acid and 10 ml of the ferric sulphate indicator solution. Then add with constant stirring, known quantity of the standard silver nitrate solution in slight excess. Heat the solution to the boil and cool to room temperature and stir to coagulate the precipitate. Titrate the excess of silver nitrate with the standard potassium thiocyanate solution. The end point is indicated by the first appearance of reddish tint that persists for 15 seconds.’

(Page 5, clause 4.3.1) — Substitute the following for the existing text:

$$\text{‘Chlorine as sodium chloride (on moisture free basis), percent by mass} = \frac{58.45 \times (AN_1 - BN_2)}{m(100 - M)} \times 100$$

where

- A = volume in ml of the standard silver nitrate solution used,
- N_1 = normality of the standard silver nitrate solution,
- B = volume in ml of the standard potassium thiocyanate solution used up by the excess silver nitrate solution,
- N_2 = normality of the standard potassium thiocyanate solution,
- m = mass in g of the material taken for the test (see 4.2.1), and
- M = percent moisture content.”

(FAD 5