

AMENDMENT NO. 2 MAY 1996 TO IS 880 : 1956 SPECIFICATION FOR TARTARIC ACID

(Page 2, *Foreword*, clause 0.1) — Insert the following new clauses after 0.1 and renumber the subsequent clauses:

0.2 Tartaric acid (2, 3-dihydroxysuccinic acid) having empirical formula $C_4H_6O_6$ and molecular mass 150.00 is a byproduct of grape fermentation industry. Commercially it may be synthesized by epoxidation of maleic acid followed by cleavage of epoxide. Its molecular formula is $[CH(OH)COOH]_2$.

0.3 Tartaric acid by virtue of having an asymmetric carbon atom possesses optical isomerism and as expected, is present in four forms. Two of these forms, namely, D (-) tartaric acid and L (+) tartaric acid exhibit optical rotation (each of these differing in signs of rotation) whereas the other two forms, namely, *racemic* and *meso* tartaric acid being 1:1 mixture of the former two-optically active forms, do not exhibit any rotation. The *racemic* variety can be resolved in optical isomers after special treatments whereas *meso* variety cannot be resolved by any method. Synthetic tartaric acid, except when resolved, is obtained in *racemic* form and hence does not have any optical rotation. The natural tartaric acid, depending upon its optical purity will show some specific rotation.

0.4 Tartaric acid is used as acidulant in soft drink, confectionery, bakery, gelation dessert industries, textile industry and silvering glass mirrors. It is also used in photography, tanning, ceramic industries, besides as raw material for various tartrates.'

[Page 3, Table 1, *Sl No.* (i), col 4 and 5] — Add '* (204 to 206)'.

[Page 3, Table 1, *Sl No.* (x)] — Insert a new requirement as follows:

(1)	(2)	(3)	(4)	(5)	(6)
xi)	†Optical rotation in 20 percent solution in water	←	+11.9° to 12.3°	→	IS 326 (Part 4). 1980

(Page 3, Table 1) — Insert at the end of the table:

*This value is applicable to synthetic material only.

†This requirement shall be nil in the case of synthetic material.'

(PCD 9)

Reprography Unit, BIS, New Delhi, India