

AMENDMENT NO. 1 SEPTEMBER 1998
TO
IS 918 : 1985 SPECIFICATION FOR CALCIUM
CARBONATE, PRECIPITATED FOR
COSMETIC INDUSTRY
(Second Revision)

(Page 16, clause A-10.0) — Substitute the following for the existing:

“A-10.0 General — Two methods have been specified for the determination of heavy metals. In ‘Method A’ the colour produced with hydrogen sulphide solution is compared with that obtained with standard lead solution while in ‘Method B’, the colour produced with thioacetamide reagent is compared with that obtained with standard lead solution.”

(Page 16, clause A-10.0) — Insert ‘METHOD A’ after clause A-10.0.

(Page 16, clause A-10.3.1) — Insert ‘METHOD B’ after clause A-10.3.1:

METHOD B

A-10.4 Apparatus

A-10.4.1 Weighing Balance — 0.000 1 g accuracy.

A-10.4.2 Nessler Cylinders — 25/50 ml.

A-10.4.3 Measuring Cylinder — 10 ml, 25 ml.

A-10.4.4 Pipette — 10 ml, 2 ml.

A-10.4.5 Volumetric Flask — 100 ml.

A-10.4.6 Evaporating Dish

A-10.5 Reagents

A-10.5.1 Acetate Buffer, pH — 3.5.

Dissolve 25 g of ammonium acetate in 25 ml of water and add 38 ml of 7M hydrochloric acid. Adjust the pH to 3.5 with either 2M hydrochloric acid or 6M ammonia and dilute to 100 ml with water.

A-10.5.2 Glycerol Mixture — To 15 ml of sodium hydroxide, 1M, add 5 ml of water and 20 ml of 85 percent glycerol. Mix well.

Amend No. 1 to IS 918 : 1985

A-10.5.3 Thioacetamide Reagent — Weigh 80 mg of thioacetamide and add to it 2 ml of water. Shake to dissolve. Add to it 10 ml glycerol mixture, heat in water bath for 20 seconds, cool and use immediately.

A-10.5.4 Acetic Acid Solution — 6M

342 ml of glacial acetic acid with sufficient water to produce 1 000 ml.

A-10.5.5 Lead Nitrate (Stock) Solution — Dissolve 0.159 8 g of lead nitrate in 100 ml of water to which 1 ml of nitric acid has been added. Dilute to 1 000 ml.

A-10.5.6 Standard Lead Solution — (10 ppm/ml).

Dilute 10 ml of lead nitrate stock solution with water to 100 ml. Each ml is equivalent to 10 ppm of lead.

A-10.6 Procedure

Into 50-ml Nessler cylinder, pipette 1 ml of standard lead solution and dilute with water to 25 ml, add 2 ml acetate buffer (pH 3.5), mix. Add 1.2 ml of thioacetamide reagent, mix immediately. Dilute with water to 50 ml and allow to stand for 2 minutes.

Dissolve 1 g of sample in 25 ml of 6M acetic acid. When effervescence ceases, boil the solution for 2 minutes, cool, filter into Nessler cylinder. Add 2 ml of acetate buffer (pH 3.5), mix. Add 1.2 ml of thioacetamide reagent, mix immediately. Dilute with water to 50 ml, allow to stand for 2 minutes.

A-10.6.1 Any brown colour produced is not more intense than that produced in the standard solution.

(PCD 19)