


The Botanical Magazine, Vol. 08


William Curtis



Project Gutenberg

The Botanical Magazine, Vol. 08

William Curtis



Project Gutenberg

The Project Gutenberg eBook of The Botanical Magazine, Vol. 08

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The Botanical Magazine, Vol. 08

Author: William Curtis

Release date: February 22, 2008 [eBook #24670]

Most recently updated: January 3, 2021

Language: English

Other information and formats: www.gutenberg.org/ebooks/24670

Credits: Produced by Jason Isbell, Janet Blenkinship and the Online Distributed Proofreading Team at <https://www.pgdp.net> (This file made using scans of public domain works at the University of Georgia.)

*** START OF THE PROJECT GUTENBERG EBOOK THE
BOTANICAL MAGAZINE, VOL. 08 ***

THE

BOTANICAL MAGAZINE;

OR,

FLOWER-GARDEN DISPLAYED:

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the Open Ground, the Green-House, and the Stove, are accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according to the celebrated LINNÆUS; their Places of Growth, and Times of Flowering:

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A WORK

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as wish to become scientifically acquainted with the Plants they cultivate.

By WILLIAM CURTIS,

Author of the FLORA LONDINENSIS.

VOL. VIII.

"Much I love
To see the fair one bind the straggling pink,
Cheer the sweet rose, the lupin, and the stock,
And lend a staff to the still gadding pea.
Ye fair, it well becomes you. Better thus
Cheat time away, than at the crowded rout,
Rustling in silk, in a small room, close-pent,
And heated e'en to fusion; made to breathe
A rank contagious air, and fret at whist,
Or sit aside to sneer and whisper scandal."

Village Curate, p. 74.

LONDON:

PRINTED BY STEPHEN COUCHMAN, For W. CURTIS,
N^o 3, *St. George's-Crescent*, Black-Friars-Road;
And Sold by the principal Booksellers in Great-Britain and Ireland, M DCC
XCIV.

CONTENTS

- [253]—LATHYRUS ARTICULATUS. JOINTED-PODDED LATHYRUS.
- [254]—LOPEZIA RACEMOSA. MEXICAN LOPEZIA.
- [255]—CYTISUS SESSILIFOLIUS. SESSILE-LEAV'D, OR COMMON CYTISUS.
- [256]—IXIA LONGIFLORA. LONG-FLOWER'D IXIA.
- [257]—LYCHNIS CHALCEDONICA. SCARLET LYCHNIS.
- [258]—CORONILLA VARIA. PURPLE CORONILLA.
- [259]—LILIUM CATESBÆI. CATESBY'S LILY.
- [260]—METROSIDEROS CITRINA. HARSH-LEAV'D METROSIDEROS.
- [261]—ERODIUM INCARNATUM. FLESH-COLOURED CRANE'S-BILL.
- [262]—MESEMBRYANTHEMUM AUREUM. GOLDEN FIG-MARIGOLD.
- [263]—GLYCINE BIMACULATA. PURPLE GLYCINE.
- [264]—CISTUS FORMOSUS. BEAUTIFUL CISTUS.
- [265]—IXIA BULBOCODIUM. CROCUS-LEAV'D IXIA.
- [266]—RANUNCULUS AMPLEXICAULIS. PLANTAIN-LEAVED CROWFOOT.
- [267]—PYRUS SPECTABILIS. CHINESE APPLE TREE.
- [268]—GLYCINE RUBICUNDA. DINGY-FLOWERED GLYCINE.
- [269]—ORNITHOGALUM NUTANS. NEAPOLITAN STAR OF BETHLEHEM.
- [270]—GLYCINE COCCINEA. SCARLET GLYCINE.
- [271]—CYRTANTHUS ANGUSTIFOLIUS. NARROW-LEAVED CYRTANTHUS.
- [272]—GLADIOLUS TRISTIS. SQUARE-LEAVED CORN-FLAG.
- [273]—DIOSMA UNIFLORA. ONE-FLOWERED DIOSMA.
- [274]—BORBONIA CRENATA. HEART-LEAVED BORBONIA.
- [275]—LIRIODENDRON TULIPIFERA. COMMON TULIP-TREE.
- [276]—BLITUM VIRGATUM. STRAWBERRY BLITE.
- [277]—MAHERNIA PINNATA. WINGED MAHERNIA.
- [278]—LILIUM CANDIDUM. WHITE LILY.
- [279]—PLUMERIA RUBRA. RED PLUMERIA.
- [280]—APOCYNUM ANDROSÆMIFOLIUM. TUTSAN-LEAV'D, OR FLY-CATCHING DOGSBANE.
- [281]—TURNERA ANGUSTIFOLIA. NARROW-LEAV'D TURNERA.

[282]—HEDYSARUM OBSCURUM. CREEPING-ROOTED HEDYSARUM.

[283]—MIMULUS RINGENS. NARROW-LEAVED MONKEY-FLOWER.

[284]—ROSA SEMPERFLORENS. EVER-BLOWING ROSE.

[285]—JASMINUM ODORATISSIMUM. SWEETEST JASMINE.

[286]—PORTLANDIA GRANDIFLORA. GREAT-FLOWERED PORTLANDIA.

[287]—GOODENIA LÆVIGATA. SMOOTH GOODENIA.

[288]—PASSIFLORA CILIATA. FRINGED-LEAVED PASSION-FLOWER.

INDEX I.

INDEX II.

[253]

LATHYRUS ARTICULATUS. JOINTED-PODDED LATHYRUS.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Stylus planus, supra villosus, superne latior. *Cal.* laciniae superiores 2-breviores.

Specific Character and Synonyms.

LATHYRUS *articulatus* pedunculis subunifloris, cirrhis polyphyllis; foliolis alternis. *Linn. Syst. Veg. ed. 14. Murr. p. 662. Ait. Kew. v. 2. p. 41.*

CLYMENUM hispanicum, flore vario, siliqua articulata. *Tourn. Inst. 396.*

LATHYRUS hispanicus, pedunculis bifloris, cirrhis polyphyllis foliolis alternis. *Mill. Dict. ed. 6. 4to.*



Nº253.

[View larger image](#)

The seed-vessels are of the first importance in ascertaining the several species of *Lathyrus*, some being naked, others hairy, some long, others short, some having a smooth and perfectly even surface, others, as in the present instance, assuming an uneven or jointed appearance.

Of this genus we have already figured three annual species, common in flower-gardens, viz. *odoratus*, *tingitanus*, and *sativus*; to these we now add the *articulatus*, not altogether so frequently met with, but meriting a place on the flower-border, as the lively red and delicate white so conspicuous in its blossoms, causes it to be much admired.

It is a native of Italy, and was cultivated at the Chelsea Garden, in the time of Mr. RAND, anno 1739.

It is a hardy annual, requiring support, and rarely exceeding the height of two feet, flowering in July and August, and is readily raised from seeds, which should be sown in the open border at the beginning of April.

LOPEZIA RACEMOSA. MEXICAN LOPEZIA.

Class and Order.

MONANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* irregularis, pentapetala, duo superiora geniculata, quintum inferne declinatum, plicatum, ungue arcuata.

Specific Character and Synonyms.

LOPEZIA *racemosa* caule herbaceo ramoso; foliis alternis ovato-lanceolatis, serratis; floribus racemosis. *Cavanilles Ic. et descr. Pl.*



Nº254.

[View larger image](#)

Some plants have a claim on our attention for their utility, some for their beauty, and some for the singularity of their structure, and the wonderful nature of their œconomy; in the last class we must place the present plant, the flowers of which we recommend to the examination of such of our readers as may have an opportunity of seeing them; to the philosophic mind, not captivated with mere shew, they will afford a most delicious treat.

We first saw this novelty in flower, towards the close of the year 1792, at the Apothecaries Garden, Chelsea, where Mr. FAIRBAIRN informed me, that he had that season raised several plants of it from seeds, communicated by Dr. J. E. SMITH, who received them from Madrid, to which place they were sent from South-America, and where the plant as Mons. CAVANILLE informs us, grows spontaneously near Mexico. In October 1793, we had the pleasure of seeing the plant again in blossom in the aforesaid garden, raised from seeds which ripened there the preceding year, but unfortunately from the lateness of their flowering, and the very great injury the plants had sustained from the Cobweb Mite (*Acarus telarius*) vulgarly called the red Spider, there seemed little prospect that the seed-vessels would arrive at perfection.

The seeds were sown by Mr. FAIRBAIRN, in March, and the plants kept in the green-house till very late in the summer, when to accelerate their blowing, they were removed into the dry stove: it is worthy of remark, that these plants, even late in the autumn, shew no signs of blossoming, but the flowers at length come forth with almost unexampled rapidity, and the seed-vessels are formed as quickly, so that if the flowers were not very numerous, their blossoming period would be of very short duration; future experience may perhaps point out the means of making the plant blow earlier: in Spain, the blossoms appeared later than here, Mons. CAVANILLE observed them in the Royal Garden, in November and December, most probably in the open ground, as no mention is made of the plants having been preserved from the weather.

It was not till long after our description was taken, that we had an opportunity of seeing Mons. CAVANILLE's most accurate and elegant work,

above quoted, in which this plant is first figured and described; we have selected the most essential parts of his generic character, and adopted his specific description: there is one point, however, in which we differ from him; the part which he regards as the fifth Petal, we are inclined to consider rather as that indescribable something, called by LINNÆUS the Nectary, it is indeed of little moment whether we call it a Petal or a Nectary, but there are several reasons why, strictly speaking, we cannot regard it as a Petal: in general the number of Petals correspond with the number of the leaves of the Calyx, those of the latter are four; the base of this Nectary originates deeper than the claws of the Petals, springing in fact from the same part as the Filament, its structure, especially the lower part of it, is evidently different from that of the Petals, corresponding indeed as nearly as possible with that of the base of the filament.—*Vid.* DESCER.

Mons. CAVANILLE was induced to call this plant *Lopezia*, in compliment to TH. LOPEZ, a Spaniard^[1].

DESCRIPTION OF THE LOPEZIA.

ROOT annual.

STALK five or six feet high, branched almost to the bottom, square, of a deep red colour, smooth towards the bottom, slightly hairy above: *Branches* like the stalk.

LEAVES alternate, ovate, pointed, toothed on the edges, more so on the larger leaves, slightly beset with soft hairs, veins prominent on the under side, usually running parallel to each other and unbranched: *Leafstalks* hairy.

FLOWERS numerous, from the alæ of the leaves, growing irregularly on hairy leafy racemi, standing on long slender peduncles, which hang down as the seed-vessels are produced: in this and some others of its characters, the plant shews some affinity to the *Circæa*.

CALYX: a *Perianthium* of four leaves, sitting on the Germen, leaves narrow, concave, reddish, with green tips, the lowermost one widely separated from the others, and placed immediately under the Nectary, *fig. 1*.

COROLLA four *Petals* of a pale red colour, forming in their mode of growth the upper half of a circle, the two uppermost linear, of a deeper colour near the apex, jointed below the middle, with a small green gland on each joint, standing on short round footstalks, which are hairy when magnified, the two side *Petals* nearly orbicular with long narrow claws, the part between the base of the Petal and the claw of a deeper red or crimson, *fig. 2*.

NECTARY situated below the *Petals*, perfectly white, somewhat ovate, the sides folding together, before the flower fully expands, nearly upright, embracing and containing within

it the Pistillum and Stamen, on touching it ever so slightly with the point of a pin, while in this state, it suddenly springs back and quits the Pistillum, the lower elastic part of it is then bent in the form represented in a magnified view of the flower on the plate, *fig. 4.* this curious phænomenon has not been noticed by CAVANILLE.

STAMEN: *Filament* one, tapering and very slender just below the Anthera, arising from the same part as (and placed opposite to the base of) the Nectary the lower part of it broader, somewhat fleshy, cartilaginous, and of the same nature as the inferior part of the Nectary, with a groove as that has on the inside, so that before the flower expands, the bases of each are like two half tubes, the sides of which, nearly touching each other, wholly enclose the Pistillum; as the fructification goes forward, the Filament, endowed also with an elastic power, bends back soon after the flower is open, betwixt the two uppermost Petals, and becomes invisible to an inattentive observer; the Anthera, which is large, is at first yellow, and afterwards dark brown, *fig. 5.*

PISTILLUM: *Germen* below the Calyx, round, smooth, and green; *Style* filiform, white, length of the Filament; *Stigma* forming a small villous head, *fig. 6.* in some of the flowers the Pistillum appears imperfect, being much shorter than usual, and wanting the Stigma, perhaps such have not acquired their full growth, *fig. 6.*

PERICARPIUM (from CAVANILLE) a round *Capsule*, of four cells, and four valves, the cells many-seeded.

SEEDS very minute, ovate, affixed to a four-cornered receptacle.

CYTISUS SESSILIFOLIUS. SESSILE-LEAV'D, OR COMMON CYTISUS.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus: 2/3 *Legumen* basi attenuatum.

Specific Character and Synonyms.

CYTISUS *sessilifolius* racemis erectis, calycibus bractæa triplici, foliis floralibus sessilibus. *Linn. Syst. Vegetab. ed. 14. Murr. p. 666. Ait. Kew. v. 2. p. 50.*

CYTISUS *glabris* foliis subrotundis, pediculis brevissimis. *Bauh. Pin. p. 390.*

CYTISUS *vulgatior*, the common Tree Trefoile. *Park. Parad. p. 440.*



Nº255.

[View larger image](#)

The term *sessilifolius* has been given to this species of *Cytisus*, because the leaves are for the most part sessile, that is sit close to the branches, without any or very short footstalks; such they are at least on the flowering branches when the shrub is in blossom, but at the close of the summer they are no longer so, the leaves acquiring very evident footstalks.

It is a native of the more southern parts of Europe, and though in point of size and elegance it cannot vie with its kindred *Laburnum*, it is a deciduous shrub of considerable beauty, rarely exceeding the height of five or six feet, and producing a great profusion of bright yellow flowers, which continue in blossom a long while; they make their appearance in May and June, and are usually succeeded by seed-vessels which produce ripe seeds, by these the plant is readily propagated.

It is one of the most common shrubs we have, as well as one of the oldest inhabitants of our shrubberies, being mentioned by PARKINSON in his *Parad.*

Terrestris.

[256]

IXIA LONGIFLORA. LONG-FLOWER'D IXIA.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Cor. 6-partita, campanulata, regularis. *Stigmata* 3.

Specific Character and Synonyms.

IXIA *longiflora* foliis ensiformibus linearibus strictis, tubo filiformi longissimo. *Ait. Kew. v. 4. p. 58.*

GLADIOLUS *longiflorus* caule tereti, tubo longissimo, spathis foliisque linearibus glabris. *Linn. Suppl. p. 96. Gmel. Syst. Vegetab. ed. 14. Murr. p. 86.*



Nº256.

[View larger image](#)

We are not acquainted with a tribe of plants which stand more in need of elucidation than those of this genus; of the vast numbers imported from the Cape within these few years, where they are chiefly natives, and that for the most part by way of Holland, few comparatively are well ascertained; some of them appear subject to great variation, both in the size and colour of their blossoms (whether in their wild state they are thus inconstant, or whether there are seminal varieties raised by the persevering industry of the Dutch Florists, we have not yet had it in our power satisfactorily to ascertain); others like the present one have their characters strongly marked, and less variable; in general they are plants of easy culture, requiring chiefly to be protected from the effects of frost, the least degree of which is presently fatal to most of them.

The treatment recommended for the *Ixia flexuosa* is applicable to this and the other Cape species.

According to the *Hort. Kew.* this species was introduced by Mr. MASSON in the year 1774.

It flowers from April to June.

[257]

LYCHNIS CHALCEDONICA. SCARLET LYCHNIS.

Class and Order.

DECANDRIA PENTAGYNIA.

Generic Character.

Cal. 1-phyllus, oblongus, lævis. *Petala* 5-unguiculata. *Limbo* sub-bifido. *Caps.* 5-locularis.

Specific Character and Synonyms.

LYCHNIS *chalcedonica* floribus fasciculatis fastigiatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 435. Ait. Kew. v. 2. p. 16.*

LYCHNIS *hirsuta* flore coccineo major. *Bauh. Pin. 203.*

FLOS Constantinopolitanus. *Dod. Pempt. 178.*

LYCHNIS *chalcedonica* flore simplici miniato. Single Nonsuch, or Flower of Bristow or Constantinople. *Parkins. Parad. 253.*



Nº257.

[View larger image](#)

The Scarlet Lychnis appears to have been a great favourite with PARKINSON, he calls it a glorious flower, and in a wooden print of him prefixed to his *Paradisus Terrestris*, we see him represented with a flower of this sort in his hand of the double kind.

It grows spontaneously in most parts of Russia, and is one of our most hardy perennials.

The extreme brilliancy of its flowers renders it a plant, in its single state highly ornamental; when double, its beauty is heightened, and the duration of it increased.

It flowers in June and July.

The single sort may be increased by parting its roots in autumn, but more abundantly by seeds, which should be sown in the spring; the double sort may also be increased by dividing its roots, but more plentifully by cuttings

of the stalk, put in in June, before the flowers make their appearance; in striking of these, however, there requires some nicety.

This plant is found to succeed best in a rich, loamy, soil; and certain districts have been found to be more favourable to its growth than others.

A white and a pale red variety of it in its single state were known to CLUSIUS, and similar varieties of the double kind are said to exist; it is of little moment whether they do or not, every variation in this plant from a bright scarlet is in every sense of the word a degeneracy.

CORONILLA VARIA. PURPLE CORONILLA.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus: 2/3: dentibus superioribus connatis. *Vexillum* vix alis longius. *Legumen* isthmis interceptum.

Specific Character and Synonyms.

CORONILLA *varia* herbacea, leguminibus erectis teretibus torosis numerosis, foliolis plurimis glabris. *Linn. Syst. Vegetab. ed. 14. Murr. p. 670. Ait. Kew. v. 3. p. 59.*

SECURIDACA *dumetorum* major, flore vario, siliquis articulatis. *Bauh. Pin. p. 349.*

SECURIDACA II. altera species. *Clus. Hist. 2. ccxxxvij.* The greater joynted Hatched Vetch. *Park. Theat. p. 1088.*



Nº258.

[View larger image](#)

CLUSIUS, in his work above referred to, informs us that he found this plant growing wild in various parts of Germany, in meadows, fields, and by road sides; that it flowered in June, sometimes the whole summer through, and ripened its seeds in July and August; the blossoms he found subject to much variation of colour, being either deep purple, whitish, or even wholly white: CASP. BAUHINE notices another variety, in which the alæ are white and the rostrum purple; this variety, which we have had the honour to receive from the Earl of EGREMONT is the most desirable one to cultivate in gardens, as it is more ornamental than the one wholly purple, most commonly met with in the nurseries, and corresponds also better with its name of *varia*; it is to be noted however that this variety of colour exists only in the young blossoms.

The *Coronilla varia* is a hardy, perennial, herbaceous plant, climbing, if supported, to the height of four or five feet, otherwise spreading widely on the ground, and frequently injuring less robust plants growing near it; on this account, as well as from its having powerfully creeping roots whereby

it greatly increases, though a pretty plant, and flowering during most of the summer, it is not to be introduced without caution, and is rather to be placed in the shrubbery, or outskirts of the garden, than in the flower border.

It will grow in any soil or situation, but blossoms and seeds most freely in a soil moderately dry.

PARKINSON in his *Theater of Plants*, mentions its being cultivated, as an ornamental plant. *Ait. Kew.*

Its bitterness, will be an objection to its being cultivated for the use of cattle, for which purpose it has been recommended.

LILIUM CATESBÆI. CATESBY'S LILY.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. 6-petala campanulata: linea longitudinali nectarifera. *Caps.* valvulis pilo cancellato connexis.

Specific Character and Synonyms.

LILIUM *Catesbæi* caule unifloro, petalis erectis unguiculatis.
Linn. Syst. Nat. ed. 13. Gmel. p. 545.

LILIUM *Catesbæi* foliis sparsis, bipedali, flore unico erecto,
corolla campanulata, petalis unguibus angustis longis.
Walt. Fl. Carol. p. 123.

LILIUM *Spectabile* foliis sparsis; floribus solitariis erectis;
petalorum unguibus angustis, alternis extus utrinque
sulcatis, laminis revolutis. *Salisb. Ic. Stirp. rar. t. 5.*



Nº259.

[View larger image](#)

At the close of the year 1787, Mr. ROBERT SQUIBB, sent me from South-Carolina roots of the Lily here figured, many of which have since flowered with various persons in this kingdom.

CATESBY in his Natural History of Carolina, gives a figure and short account of it; WALTER in his *Flora Caroliniana* describes it under the name of *Lilium Catesbæi*; Mr. SALISBURY in the first number of his very magnificent work, lately published, presents us with a very highly finished likeness of this lily, accompanied by a most accurate and minute description of it, and judging from some appearances in CATESBY's figure, that it was not the *Lilium Catesbæi* of WALTER, names it *spectabile*; but as we are assured by Mr. SQUIBB, who assisted his friend WALTER in his publication, that it was the lily figured by CATESBY, we have continued the name given in honour of that Naturalist.

Of the different Lilies cultivated in this country, this is to be numbered among the least, the whole plant when in bloom being frequently little more than a foot high; in its native soil it is described as growing to the height of two feet; the stalk is terminated by one upright flower, of the form and colour represented on the plate; we have observed it to vary considerably in the breadth of its petals, in their colour, and spots.

It flowers usually in July and August.

This plant may be raised from seeds, or increased by offsets, which, however, are not very plentifully produced, nor is the plant to be made grow in perfection without great care, the roots in particular are to be guarded against frost; the soil and situation may be the same as recommended for the *Cyclamen Coum.* p. 4. v. 1.

[260]

METROSIDEROS CITRINA. HARSH-LEAV'D METROSIDEROS.

Class and Order.

ICOSANDRIA MONOGYNIA.

Generic Character.

Cal. 5-dentatus, sinu germen fovens. *Petala* 5, caduca. *Stam.* discreta, petalis multoties longiora. *Caps.* 3-4 locularis, polysperma. *Banks. Gærtner.*

Specific Character and Synonyms.

METROSIDEROS *citrina* foliis lineari-lanceolatis rigentibus.



Nº260.

[View larger image](#)

Though many species of this genus have been raised from seeds, brought within these few years from the South Seas, where they are said to be very numerous; this is, we believe, the only one that as yet has flowered in this country: our drawing was made from a plant which blossomed toward the close of last summer at Lord CREMORNES, the root of which had been sent from Botany-Bay; previous to this period we have been informed, that the same species flowered both at Kew and Sion-House: as it is without difficulty raised both from seeds and cuttings, young plants of it are to be seen in most of the Nurseries near town; it would seem that they do not flower till they are at least five or six years old.

Metrosideros is a name given originally by RUMPHIUS in *Herb. Amboin* to some plants of this genus, the term applies to the hardness of their wood, which by the Dutch is called Yzerhout (Ironwood): FORSTER in his *Gen. Pl.* figures this and another genus on the same plate, under the name of *Leptospermum*; SCHREBER in his edition of the *Gen. Pl.* of LINNÆUS, unites *Metrosideros*, *Melaleuca*, *Leptospermum*, and *Fabricia*, under the genus *Melaleuca*; GÆRTNER in his elaborate work on the seeds of plants, makes separate genera of these, agreeably to the ideas of Sir JOSEPH BANKS and Mr. DRYANDER, who on this subject can certainly boast the best information.

We cannot, without transgressing the allotted limits of our letter-press, give a minute description of the plant figured; suffice it to say, that it is an ever-green shrub, growing to the height of from four to six or more feet, that its leaves on the old wood feel very harsh or rigid to the touch, and when bruised give forth an agreeable fragrance, the flowers grow in spikes on the tops of the branches, and owe their beauty wholly to the brilliant colour of the filaments.

[261]

ERODIUM INCARNATUM. FLESH-COLOURED CRANE'S-BILL.

Class and Order.

MONADELPHIA PENTANDRIA.

Generic Character.

Cal. 5-phyllus. *Cor.* 5-petala. *Nect.* *Squamulæ* 5 cum filamentis alternantes; et glandulæ melliferæ, basi staminum insidentes. *Fructus* 5-coccus, rostratus; rostra spiralia, introrsum barbata. *L'Herit. Geran.*

Specific Character and Synonyms.

ERODIUM *incarnatum* pedunculis paucifloris, foliis tripartitis ternatisve trifidis scabris, caule fruticuloso. *L'Herit. n. 21. tab. 5. Ait. Kew. v. 2. p. 415.*

GERANIUM *incarnatum* pedunculis bifloris, foliis tripartitis trifidis glabris, petalis integris, arillis glabris. *Linn. Suppl. Pl.*

GERANIUM *incarnatum* foliis incisis quinquelobis punctatis; petiolis longissimis, pedunculis trifloris. *Cavanill. diff. 4. p. 223. n. 314. t. 97. f. 3.*



Nº261.

[View larger image](#)

In the 80th number of this work we gave a figure of the *Pelargonium tricolor*, a plant very generally regarded as the most beautiful of the genus; we now present our readers with the representation of an *Erodium*, which has to boast nearly an equal share of admiration.

This species, as we learn from the *Hortus Kewensis*, is a native of the Cape, and was introduced by Mr. MASSON in the year 1787.

Its usual time of flowering is July and August; in this point it is inferior to the *Pelargonium tricolor*, which blossoms through the spring as well as summer months.

It produces seeds but sparingly; cuttings of the plant are struck with less difficulty than those of the *Pelargonium* above mentioned, the same treatment is applicable to both plants, they must be regarded as green-house plants of the more tender kind, which are liable to be destroyed in the winter season by a moist cold atmosphere.

[262]

MESEMBRYANTHEMUM AUREUM. GOLDEN FIG-MARIGOLD.

Class and Order.

ICOSANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-fidus. *Petala* numerosa linearia. *Caps.* carnosae inferae polyspermae.

Specific Character and Synonyms.

MESEMBRYANTHEMUM *aureum* foliis cylindrico-triquetris punctatis distinctis, pistillis atro purpurascentibus. *Linn. Syst. Nat. ed. 10. p. 1060. Ait. Kew. v. 2. p. 190.*



Nº262.

[View larger image](#)

This Mesembryanthemum is one of the taller and more upright species, as well as the earliest in point of flowering, producing its blossoms from February to May; these are large and of a bright orange hue, the pistilla in the centre are purple, and serve at once to distinguish and embellish them.

It was first described in the *10th ed.* of *Linn. Syst. Nat.* and afterwards inserted in the *Hort. Kew.* of Mr. AITON, who informs us that it is a native of the Cape, and was cultivated by Mr. MILLER, in the year 1750. Prof. MURRAY omits it in his *12th ed.* of the *Syst. Vegetab.* of LINNÆUS, as does Prof. GMELIN in the last edition of *Linn. Syst. Nat.*

The facility with which this tribe in general is increased by cuttings is well known; this is raised as readily as the others.

[263]

GLYCINE BIMACULATA. PURPLE GLYCINE.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus. *Corollæ* carina apice vexillum reflectens.

Specific Character.

GLYCINE *bimaculata* caule volubili lævi, foliis simplicibus cordato-oblongis, racemis multifloris.



Nº263.

[View larger image](#)

Of the many plants which within these few years have been raised from Botany-Bay seeds, this is one of the first which flowered in this country, and one of the most ornamental; to the greenhouse it is indeed an invaluable acquisition: we regret that the size of our paper and the imperfection of the colouring art, will not admit of our giving a representation of it more adequate to its beauty.

It rises up with a twining shrubby stalk to the height of six, eight, or more feet; these multiplying greatly by age, become loaded with a profusion of purple flowers, growing in racemi, the richness of which is enlivened by the appearance of two green spots at the base of the vexillum; for the most part the blossoms go off with us without producing any seed-vessels; in some instances, however, perfect seeds have been produced, and we have seen a plant in bloom raised from such in the charming retreat of JOHN ORD, Esq. Walham-Green.

A great excellence of this plant is the duration of its flowering period, it begins to put forth its blossoms in February, and continues to do so during most of the summer.

In the Nurseries about town, it is known by the name of *Glycine virens*, a name given the plant originally by Dr. SOLANDER; the latter of these terms we have taken the liberty of changing to *bimaculata*, as being more expressive of an obvious character in the flower: we might, perhaps, been justified in altering the genus, as its characters do not appear to be peculiarly expressive of a *Glycine*, nor indeed of any other genus in this numerous natural order.

It is raised readily from seeds.

We think it highly probable, that in warm sheltered situations, this climber might grow in the open ground; to such as have it in abundance, we recommend them to make the experiment.

[264]

CISTUS FORMOSUS. BEAUTIFUL CISTUS.

Class and Order.

POLYANDRIA MONOGYNIA.

Generic Character.

Cor. 5-petala. *Cal.* 5-phyllus, foliolis duobus minoribus.
Capsula.



N^o264.

[View larger image](#)

Mr. LEE, Nurseryman of Hammersmith, informs me, that in the year 1780, he raised the Cistus here figured from seeds, the produce of Portugal, and as

its flowers were uncommonly beautiful, he was induced to name it *formosus*.

It approaches so near to the *Cistus halimifolius* in point of habit, in the form and colour of its leaves and flowers, that we are inclined rather to regard it as a variety of that plant, than as a distinct species; at the same time it must be allowed to be a very striking variety, the flowers being at least thrice as large as those of the *halimifolius* usually are, and the whole plant more hairy: as an ornamental shrub, it is highly deserving a place in all curious collections.

It will grow very well in the open border in warm sheltered situations, it may be kept also in a pot, by which means it may more readily be sheltered during the winter, either in the greenhouse or under a frame.

It flowers early in May, and may be increased by cuttings.

[265]

IXIA BULBOCODIUM. CROCUS-LEAV'D IXIA.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Cor. 1-petala, tubulosa; tubo recto, filiformi; limbo 6-partito, campanulato, æquali. *Stigmata* tria, simplicia. *Thunb. Diss. de Ixia.*

Specific Character and Synonyms.

IXIA *Bulbocodium* scapo unifloro brevissimo, foliis angulatis caulinis, stigmatibus sextuplicibus. *Linn. Syst. Nat. ed. 13. p. 76.*

IXIA *Bulbocodium* scapo ramoso, floribus solitariis, foliis sulcatis reflexis. *Thunb. Diss. n. 3.*

CROCUS vernus angustifolius. 1. 11. *Clus. Hist. i. p. 207. violaceo flore, 208. ejusd.*



Nº265.

[View larger image](#)

There are three plants cultivated in the gardens of the curious to which *Bulbocodium* is applied, either as a generic or a trivial name, viz. *Narcissus Bulbocodium*, *Bulbocodium vernum*, already figured, and the present plant: the *Ixia Bulbocodium* and *Bulbocodium vernum* are given in this work, not so much for their beauty as their rarity, not so much to gratify the eye, as to communicate a knowledge of two plants but little known, and liable to be confounded from a similarity of their names.

This is one of the few hardy species of the genus, and grows wild in many parts of Spain and Italy; it is said to have been found in Guernsey: it affects hilly and dry situations, will grow readily in almost any soil, especially if fresh, and not infested with vermin: it flowers about the middle of April, the blossoms do not expand fully unless exposed to the sun, and are not of long duration: authors describe the wild plants as varying greatly in colour, *vid. Clus.* they are most commonly pale blue.

Like the Crocus, it increases readily by offsets.

Was cultivated by Mr. MILLER, in 1739, *Ait. Kew. Bulbocodium*, 1. in the 6th edition of his Dictionary in 4to, is not this plant, but the *Anthericum scrobinum*, *Jacq. Fl. Austr. v. 5. app. t. 38.*

[266]

RANUNCULUS AMPLEXICAULIS. PLANTAIN-LEAVED CROWFOOT.

Class and Order.

POLYANDRIA POLYGYNIA.

Generic Character.

Cal. deciduus. 5 phyllus, (rarius 3-phyllus) *Petala* 5, (rarius 2, 3, aut 8) intra ungues squamula vel poro mellifero. *Styli* persistentes. *Sem.* incrustata, erecta. *Linn. Syst. Nat. ed. 13. Gmel.*

Specific Character and Synonyms.

RANUNCULUS *amplexicaulis* foliis ovatis acuminatis amplexicaulibus, caule multifloro, radice fasciculata. *Linn. Syst. Vegetab. ed. 14. Murr. p. 515. Ait. Kew. v. 2. p. 265.*

RANUNCULUS *montanus* foliis plantaginis. *Bauh. Pin. 180.*

RANUNCULUS *pyrenæus* albo flore. *Clus. app. alt. auct. ic. 4 ta. Ger. emac. 963. fig. 2.*



Nº266.

[View larger image](#)

The leaves of the *Ranunculus amplexicaulis* in part surround the stalk at their base, whence its trivial name; in colour they differ from most others of the genus, being of a greyer or more glaucous hue, which peculiarity joined to the delicate whiteness of the flowers, renders this species a very desirable one to add to a collection of hardy, ornamental, herbaceous plants, more especially as it occupies but little space, and has no tendency to injure the growth of others.

It is a native of the Apennine and Pyrenean mountains, and flowers in April and May.

CLUSIUS is the first author who describes and figures this species. JOHNSON in his *ed.* of GERARD copies his figure, and mentions it as being then made a denizen of our gardens.

It is readily propagated by parting its roots in Autumn, and provided it has a pure air will succeed in most soils an situations.

[267]

PYRUS SPECTABILIS. CHINESE APPLE TREE.

Class and Order.

ICOSANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-fidus. *Petala* 5. *Pomum* inferum, 5-loculare, polyspermum.

Specific Character and Synonyms.

PYRUS *spectabilis* umbellis sessilibus, foliis ovali oblongis serratis lævibus, unguibus calyce longioribus, stylis basi lanatis. *Ait. Kew.* v. 2. p. 175. *Linn. Syst. Nat. ed. 13.* *Gmel.* p. 842.



Nº267.

[View larger image](#)

The Chinese Apple-Tree when it blossoms in perfection, answers truly to the name of *spectabilis*; a more shewy or ornamental tree can scarcely be introduced to decorate the shrubbery or plantation; its beauty like that of most trees, whose ornament consists chiefly in their blossoms, is however but of short duration, and depends in some degree on the favourableness of the season at the time of their expansion, which usually takes place about the end of April or beginning of May; the flowers are large, of a pale red when open, and semi-double, the buds are of a much deeper hue, the fruit is of little account, and but sparingly produced. Trees of this species are to be met with in some gardens of the height of twenty or thirty feet.

Dr. FOTHERGILL is regarded as the first who introduced this Chinese native, he cultivated it in the year 1780; such plants of it as were in his collection, passed at his decease into the hands of Messrs. GORDON and THOMPSON, in whose rich and elegant Nursery, at Mile-End, this tree may be seen in great perfection.

Though perfectly hardy, as its blossoms are liable to be injured by cutting winds, it will be most proper to plant it in a shelter'd situation.

It is usually increased by grafting it on the Crab stock.

[268]

GLYCINE RUBICUNDA. DINGY-FLOWERED GLYCINE.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus. *Corollæ* carina apice vexillum reflectens.

Specific Character.

GLYCINE *rubicunda* caule perenni volubili, foliis ternatis, foliolis subovalibus integerrimis, pedunculis subtrifloris.



N^o268.

[View larger image](#)

The plant here figured, and very generally known to the Nurserymen, in the neighbourhood of London, by the name of *Glycine rubicunda*, is a native of New South-Wales, and was introduced to this country about the same time as the *Glycine bimaculata* already figured.

It is a shrubby, twining plant, running up to the height of five, six, or more feet, producing blossoms abundantly from April to June, which are usually succeeded by seed-vessels which ripen their seeds with us.

The flowers though large and shewy, have a kind of dingy or lurid appearance, which greatly diminishes their beauty. We have observed the blossoms of some plants more brilliant than those of others, and we think it highly probable, that, at some future period, seminal varieties may be obtained with flowers highly improved in colour.

This species is readily raised from seeds, is of quick growth, and may be regarded as one of our more hardy green-house plants: probably it may succeed in the open air, if planted in a warm situation, and sheltered in inclement seasons.

[269]

ORNITHOGALUM NUTANS. NEAPOLITAN STAR OF BETHLEHEM.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. 6 petala, erecta, persistens, supra medium patens,
Filamenta alterna basi dilatata.

Specific Character and Synonyms.

ORNITHOGALUM *nutans* floribus secundis pendulis, nectario
stamineo campaniformi. *Linn. Syst. Vegetab. ed. 14. Murr.*
p. 328. Ait. Kew. v. i. p. 443.

ORNITHOGALUM exoticum magno flore minori innato. *Bauh.*
Pin. p. 70.

ORNITHOGALUM Neopolitanum, the Starre-flower of Naples.
Park. Parad. p. 138. p. 137. f. 8. Clus. app. alt. p. 9. fig. 7.



Nº269.

[View larger image](#)

Authors have given to this species of *Ornithogalum* the name of Neapolitan, following CLUSIUS by whom the plant is figured and described, and who so called it, merely on receiving it from Naples; it may perhaps be doubted whether it be originally a native of Italy. Prof. JACQUIN has figured it in his *Flora Austriaca*, the plant being common about Vienna, in garden-walks, under hedges, and in meadows, he does not however, from that circumstance, regard it as an original native there. CASP. BAUHIN informs us that HONORIUS BELLI sent it him from Crete under the name of *Phalangium*, leaving its true habitat to be settled more precisely hereafter, we shall observe, that it is one of those plants which soon accommodate themselves to any country; producing a numerous progeny both from roots and seeds, and by no means nice as to soil or situation; it is not long before it becomes a weed in the garden, from whence it is apt like the *Hyacinthus racemosus*, already figured, to pass into the field or meadow.

Its flowers, which if not beautiful are singular and delicate, make their appearance towards the end of April, they are of no long duration, seldom continuing above a fortnight, and are succeeded by seed-vessels which produce abundance of ripe seed, by which, as well as by its bulbs, the plant may be increased.

In the *Hortus Kewensis* it is set down as a Greenhouse plant, one of the rare errors which occur in that most useful work.

[270]

GLYCINE COCCINEA. SCARLET GLYCINE.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus. *Corollæ* carina apice vexillum reflectens.

Specific Character.

GLYCINE *coccinea* foliis ternatis, foliolis subrotundis undulatis.



Nº270.

[View larger image](#)

We here present our readers with another Glycine, very lately raised by several persons in the neighbourhood of London from Botany-Bay seeds, and which we have called *coccinea* from the colour of its blossoms.

It is a shrubby, climbing plant, which, if supported, will grow to the height of many feet, producing a great number of flowers on its pendant branches; the leaves, which grow three together, are nearly round, and, in the older ones especially, are crimped or curled at the edges; the flowers grow for the most part in pairs, are of a glowing scarlet colour, at the base of the carina somewhat inclined to purple, the bottom of the vexillum is decorated with a large yellow spot, verging to green, which adds much to the beauty of the flower.

It blossoms from April to June, and appears to be fully as much disposed to produce seed vessels, and perfect seeds, as the *rubicunda*, and by which alone it has hitherto been propagated.

We must rank it among the more tender green-house plants.

[271]

**CYRTANTHUS ANGUSTIFOLIUS. NARROW-LEAVED
CYRTANTHUS.**

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. tubulosa, clavata, curva, 6-fida, laciniae ovato-oblongae.
Filamenta tubo inserta, apice conniventia. *Linn. Fil.*

Specific Character and Synonyms.

CYRTANTHUS *angustifolius* foliis obtuse carinatis rectis,
floribus cernuis, *Linn. Fil. Ait. Kew. v. i. p. 414.*

CRINUM *angustifolium* foliis linearibus obtusis, corollis
cylindricis: laciniis alternis interglandulosis. *Linn. Suppl. 195.*



Nº271.

[View larger image](#)

CYRTANTHUS is a genus which takes its name from the curvature of its flower, was established by the younger LINNÆUS, and adopted by Mr. AITON in the *Hortus Kewensis*.

The present species is a native of the Cape, and was added to the royal collection at Kew, by Mr. MASSON, in the year 1774. The plant from whence our drawing was made flowered the preceding May with Mr. WHITLEY, Nurseryman, Old Brompton, who received it from Holland, and who has been so fortunate as to obtain young plants of it from seed.

It flowers in May and June; requires the same treatment as other Cape bulbs, and may be increased by offsets and seeds.

At the extremity of each alternate segment of the corolla there is a kind of small glandular hook, deserving of notice.

[272]

GLADIOLUS TRISTIS. SQUARE-LEAVED CORN-FLAG.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Cor. 6-partita, ringens. *Stamina* adscendentia.

Specific Character and Synonyms.

GLADIOLUS *tristis* foliis lineari-cruciatis, corollis campanulatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 86. Ait. Kew. v. 1. p. 63.*

LILIO-GLADIOLUS bifolius et biflorus, foliis quadrangulis. *Trew. Ehret. t. 39.*

GLADIOLUS *tristis* foliis linearibus sulcatis, caule bifloro, tubo longissimo, segmentis æqualibus. *Mill. Dict. ed. 6. 4to.*



Nº272.

[View larger image](#)

LINNÆUS gave to this species of *Gladiolus* the name of *tristis*, from the colour of its flowers, which however possess scarcely sufficient of the sombre to justify the appellation; still less so if they vary in the manner represented in TREW's *Ehret*, where they are painted in gay and lively colours: in the specimens we have seen, the blossoms have been of a sulphur colour, shaded in particular parts with very fine pencillings, especially on the under side: most authors describe the flowering stems as producing only two flowers, LINNÆUS has observed that they sometimes produce many, we have seen them do so where the plant has grown in perfection; in their expansion, which usually takes place in April and May, they give forth a most agreeable fragrance.

It is a native of the Cape, and other parts of Africa; was cultivated by Mr. MILLER, and flowered in the Chelsea Garden in the year 1745. *Ait. Kew.*

The leaves which so characteristically distinguish this species are highly deserving of notice, instances of such rarely occur; as the bulbs produce numerous offsets, the plant is propagated by them without difficulty, and requires the same treatment as other Cape bulbs.

[273]

DIOSMA UNIFLORA. ONE-FLOWERED DIOSMA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. 5-petala. *Nectaria* 5, supra germen. *Caps.* 3. s. 5. coalitæ.
Sem. calyptrata.

Specific Character and Synonyms.

DIOSMA *uniflora* foliis ovato oblongis, floribus solitariis terminalibus. *Linn. Sp. Pl. ed. 3. p. 287. Syst. Vegetab. ed. 14. Murr. p. 239. Ait. Kew. v. 1. p. 276.*

CISTUS *humilis æthiopicus*, inferioribus foliis rosmarini sylvestris punctatis, cæteris autem serpylli subrotundis, flore carneo. *Pluk. mant. 49. t. 342. f. 5.*



N^o273.

[View larger image](#)

The *Diosma uniflora* another native of the Cape, that never failing source of vegetable riches, was introduced to the Royal Garden at Kew by Mr. MASSON in the year 1775, it flowers in our Green-Houses from April to June, and is usually propagated by cuttings.

This plant forms a small bushy shrub, the leaves are thickly and irregularly set on the branches, quite up to the flowers, which stand singly on their summits, and are larger than those of any other known species of *Diosma*, expanding as we have found on trial beyond the size of half-a-crown, which the blossom does in our figure, though it will not appear to do so to the eye of most observers; they are without scent, the calyx is large and continuing, composed of five ovato-lanceolate leaves, reddish on the upper side, and if viewed from above visible between the petals; the petals are five in number, much larger than the calyx, and deciduous, of a white colour with a streak of red running down the middle of each, surface highly glazed, the stamina are composed of five short filaments, white and slightly hairy, broad at their

base and tapering gradually to a fine point, by which they are inserted into the hind part of the antheræ, near the bottom; the antheræ are as long as the filaments, of a brown purple colour, bending over the stigma, and opening inwardly, each carrying on the upper part of its back a gland-like substance, of a pale brown colour: besides these parts there are five filamentous bodies alternating with, and of the same length as the stamina, of a white colour, and hairy, each dilating at its extremity where it is of a reddish hue, and presenting towards the antheræ an oval somewhat concave surface, which secretes a viscous liquid; in some flowers that we have examined, and we regret seeing but few, we have observed these nectaries (for such they may be strictly called) closely adhering by their viscous summits to the glandular substances at the back of the antheræ^[2]; the germen is studded with a constellation of little glands, which pour forth, and almost deluge it with nectar; the stigma is composed of five little round knobs: seed vessels we have not seen.

[274]

BORBONIA CRENATA. HEART-LEAVED BORBONIA.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. acuminato-spinosus. *Stigma* emarginatum. *Legumen* mucronatum.

Specific Character and Synonyms.

BORBONIA *crenata* foliis cordatis multinerviis denticulatis.
Linn. Syst. Vegetab. ed. 14. Murr. p. 643. Ait. Kew. v. 2. p.
9.

FRUTEX *æthiopicus* leguminosus, foliis rusci majoribus in
ambitu spinulis fimbriatis. *Pluk. Alm. 159.*

PLANTA leguminosa æthiopica, foliis rusci. *Breyn. Cent. t. 28.*



Nº274.

[View larger image](#)

Borbonia is a genus of plants established by LINNÆUS in the 6th edition of his *Genera Plantarum*; of this genus there are six species enumerated in the 3d edition of the *Species Plant.* and two in the *Hort. Kew.* the latter of which, the *crenata*, introduced from the Cape by Mr. MASSON, in 1774, is here figured.

It is a small shrubby plant, rarely exceeding the height of three feet, producing its flowers in a small cluster on the summits of the branches; these are of a yellow colour, and have nothing about them peculiarly singular, or beautiful; it is the foliage alone which renders this plant desirable in a collection.

It flowers from June to August, and in favourable seasons ripens its seeds, by which the plant is usually propagated.

[275]

LIRIODENDRON TULIPIFERA. COMMON TULIP-TREE.

Class and Order.

POLYANDRIA POLYGYNIA.

Generic Character.

Cal. 3-phyllus. *Petala* 6. *Sem.* imbricata in strobilum.

Specific Character and Synonyms.

LIRIODENDRON *Tulipifera* foliis lobatis. *Linn. Syst. Veg. ed.*
14. Murr. p. 507. Ait. Kew. v. 2. 250.

TULIPIFERA virginiana, tripartito aceris folio: media lacinia
velut abscissa. *Pluk. Alm. 379. t. 117. f. 5. & t. 248. f. 7.*
Catesb. Carol. 1. p. 48. t. 48.

LIRIODENDRON foliis angulatis truncatis. *Trew. Ehret. t. x.*



N^o275.

[View larger image](#)

The Tulip-tree is a native of most parts of North-America, MARSHALL describes it as often growing to the size of a very large tree, 70 or 80 feet in height, and above 4 feet in diameter; he mentions two varieties, one with yellow and the other with white wood; that with yellow wood is soft and brittle, much used for boards, heels of shoes, also turned into bowls, trenchers, &c. the white is heavy, tough, and hard, and is sawed into joists, boards, &c. for building.

RAY informs us in his *Hist. Pl.* that this tree was cultivated here by Bishop COMPTON, in 1688: and from MILLER we learn, that the first tree of the kind which flowered in this country, was in the gardens of the Earl of PETERBOROUGH, at Parsons-Green, near Fulham; in Mr. ORD's garden, at Walham-Green, there is, among other choice old trees, a very fine tulip-tree, which is every year covered with blossoms, and which afforded us the specimen here figured. It flowers in June and July, rarely ripens its seeds with us, though it does readily in America.

The foliage of this plant is extremely singular, most of the leaves appearing as if truncated, or cut off at the extremity; they vary greatly in the division of their lobes, the flowers differ from those of the tulip in having a calyx, but agree as to the number of petals, which is six; and so they are described in the sixth edition of the *Gen. Pl.* of LINN. but in *Professor MURRAY'S Syst. Veg. Ait. H. K. Linn. Syst. Nat. ed. 13*, by GMELIN, 9 are given, this in the first instance must be a mere typographical error arising from the inversion of the 6.

This tree is found to flourish most in a soil moderately stiff and moist, is usually raised from seeds, the process of which is amply described by MILLER in his Dictionary.

[276]

BLITUM VIRGATUM. STRAWBERRY BLITE.

Class and Order.

MONANDRIA DIGYNIA.

Generic Character.

Col. 3-fidus. *Petala* 0. *Sem.* 1. calyce baccato.

Specific Character and Synonyms.

BLITUM *virgatum* capitellis sparsis lateralibus. *Linn. Syst. Vegetab. ed. 14. Murr. p. 53. Ait. Kew. v. 1. p. 7.*

ATRIPLEX *sylvestris* mori fructu. *Bauh. Pin. p. 519.*

ATRIPLEX *sylvestris* baccifera. *Clus. Hist. cxxxv.*



Nº276.

[View larger image](#)

This plant, not unfrequently met with in gardens, is known to most cultivators by the name of *Strawberry Spinach*; the leaves somewhat resembling those of the latter, and the fruit that of the former: C. BAUHINE likens its berries to those of the Mulberry, to which they certainly bear a greater resemblance: in most of the species of this genus the calyx exhibits a very singular phenomenon, when the flowering is over, it increases in size, becomes fleshy, and finally pulpy, containing the ripe seed, which however it does not wholly envelope; thus from each cluster of flowers growing in the alæ of the leaves are produced so many berries, of a charming red colour, to which the plant owes its beauty altogether, for the flowers are small, herbaceous, and not distinctly visible to the naked eye; they can boast however of being of the first class in the Linnean system *Monandria*, to which few belong.

Strawberry Blite is a hardy annual, growing spontaneously in some parts of France, Spain, and Tartary; is not a very old inhabitant of our gardens, Mr. AITON mentioning it as being first cultivated by Mr. MILLER in 1759. Its berries are produced from June to September; in their taste they have nothing to recommend them, though not pleasant they are harmless.

CLUSIUS we believe to be the first author who gives a figure and description of it.

It affects a dry soil, and open situation; in such there is no necessity to give any particular directions for its cultivation, as it comes up readily from seed spontaneously scattered, so much so as sometimes to prove a troublesome weed.

[277]

MAHERNIA PINNATA. WINGED MAHERNIA.

Class and Order.

PENTANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-dentatus. *Petala* 5. *Nectaria* 5 obcordata, filamentis supposita. *Caps.* 5-locularis.

Specific Character and Synonyms.

MAHERNIA *pinnata*, foliis tripartito pinnatifidis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 308. Ait. Kew. v. 1. p. 398.*

HERMANNIA foliis tripartitis, media pinnatifida. *Linn. Sp. Pl. ed. 3. p. 943.*

HERMANNIA *frutescens*, folio multifido tenui, caule rubro. *Boerh. Lugd. 1. p. 273.*



Nº277.

[View larger image](#)

LINNÆUS, in his *Spec. Pl.* regarded this plant as a species of *Hermannia*; finding afterwards that it differed materially in its fructification from that genus, he made a new one of it in his *Mantissa*, by the name of *Mahernia*; still, however, the two genera are very nearly related: one principal difference consists in the nectaria of the *Mahernia*, which are very remarkable.

This species was introduced from the Cape, where it is a native, by Mr. MASSON, in 1774, and is now very generally met with in our green-houses. It produces its little bells, of a lively red when they first open, from June to August, or September; is a small delicate plant, and easily raised from cuttings.

[278]

LILIUM CANDIDUM. WHITE LILY.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. 6-petala, campanulata: linea longitudinali nectarifera. *Caps.* valvulis pilo cancellato connexis.

Specific Character and Synonyms.

LILIUM *candidum* foliis sparsis, corollis campanulatis, intus glabris. *Linn. Sp. Pl. ed. 3. p. 433. Syst. Vegetab. ed. 14. Murr. p. 324. Ait. Kew. v. 1. p. 429.*

LILIUM album flore erecto et vulgare. *Bauh. Pin. 76.*

LILIUM album vulgare. The ordinary White Lily. *Park. Parad. p. 39. t. 37. f. 4.*



Nº278.

[View larger image](#)

We may rank the White Lily among the very oldest inhabitants of the flower-garden; in the time of GERARD it was very generally cultivated, and doubtless at a much earlier period; a plant of such stateliness, so shewy, so fragrant, and at the same time so much disposed to increase, would of course soon be found very generally in gardens, into which its introduction would be accelerated on another account; it was regarded as a plant of great efficacy; among other extraordinary powers attributed to it, we are gravely told that it taketh away the wrinkles of the face.

LINNÆUS makes it a native of Palestine and Syria; Mr. AITON of the Levant.

Its blossoms, which open early in July, continue about three weeks, and when they go off leave the flower-garden greatly thinned of its inhabitants.

Of the White Lily there are three principal varieties:

1. With double flowers.
2. With flowers blotched with purple.
3. With striped leaves, or leaves edged with yellow.

The two first of these are to be esteemed merely as curiosities; in the third the plant acquires an accession of beauty which it has not originally; though many persons object to variegated leaves, as conveying an idea of fickleness, that complaint cannot be urged against the foliage of the striped Lily, to which the borders of the flower-garden are indebted for one of their chief ornaments during the autumnal and winter months; early in September these begin to emerge, and towards spring another set rises up in their centre, of more upright growth, and which announce the rising of the flowering stem.

Besides these varieties, LINNÆUS has considered the *Lilium album floribus dependentibus s. peregrinum* of C. BAUHINE, the *Sultan Zambach* of CLUSIUS, and the *Hortus Eystettensis*, as one of its varieties also: MILLER regards this plant as a distinct species, and those who have attentively examined the figures and descriptions of CLUSIUS and the *Hort. Eyst.* will be of the same opinion.

The Lily increases most abundantly by offsets, hence it becomes necessary that the bulbs should be taken up, and reduced every second or third year; but the striped leaved variety increasing much more slowly, should remain unmolested for a greater length of time.

There is scarcely a soil or situation in which the Lily will not grow, it will thrive most in a soil moderately stiff and moist; though a native of a warm climate no severity of weather affects it with us: we may learn from this, not to regulate the culture of plants invariably by the climate in which they grow spontaneously.

The best time for removing the bulbs of this plant is about the middle of August, before they shoot forth their leaves; but they may be transplanted any time from September to spring.

PLUMERIA RUBRA. RED PLUMERIA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Contorta. *Folliculi* 2. reflexi. *Semina* membranæ propriæ inserta.

Specific Character and Synonyms.

PLUMERIA *rubra* foliis ovato-oblongis, petiolis biglandulosi.
Linn. Syst. Vegetab. ed. 14. Murr. p. 254. Ait. Kew. v. 1. p. 298.

PLUMERIA flore roseo odoratissimo. *Tourn. Inst. 659. Trew. Ehret. Tab. xli.*



Nº279.

[View larger image](#)

Plumeria is a genus of plants named by TOURNEFORT in honour of his countryman the celebrated PLUMIER, it comes near to Nerium or Oleander, and contains several species, all natives of warm climates.

The present plant is a native of Jamaica, where it is known by the name of Red Jasmine, from whence seeds and large cuttings are often sent to this country; here they require the stove to bring them to flower: seed-vessels they are never known to produce.

The flowers, which are very odoriferous, are produced in July and August in large bunches, on the summits of the branches, from whence the leaves also proceed; the stems, which grow to a considerable height as well as thickness, are naked, and the whole plant loses its foliage from the middle of winter till about the beginning of May; the branches and other parts of the plant, when broken off, give forth a milky juice, the leaves are handsome, and the veins remarkable.

Being too tender to bear the open air of this climate, it is kept in the stove even during summer, in hot weather it must have plenty of air, and in cold seasons be sparingly watered.

Is propagated by seeds, but more frequently by cuttings, which MILLER recommends to be put by for two months or ten weeks, previous to their being committed to the earth.

[280]

**APOCYNUM ANDROSÆMIFOLIUM. TUTSAN-LEAV'D, or
FLY-CATCHING DOGSBANE.**

Class and Order.

PENTANDRIA DIGYNIA.

Generic Character.

Cor. campanulata. *Glandulæ* 5 cum staminibus alternæ.

Specific Character and Synonyms.

APOCYNUM *androsæmifolium* caule rectiusculo herbaceo,
foliis ovatis utrinque glabris, cymis terminalibus. *Linn.*
Syst. Vegetab. ed. 14. Murr. p. 258. Ait. Kew. v. 1. p. 303.

APOCYNUM canadense; foliis androsæmi majoris. *Bocc. sicc.*
35. t. 16. f. 3. Moris. Hist. 3. p. 609. s. 15. t. 3. f. 16.



Nº280.

[View larger image](#)

In addition to the powerful recommendations of beauty and fragrance, the Tutsan-leav'd Dogsbane interests us on account of the curious structure of its flowers, and their singular property of catching flies.

This species is a native of different parts of North-America; Mr. W. HALE, of Alton, Hants, who resided at Halifax in Nova-Scotia several years, brought me some seeds of it gathered in that neighbourhood, which vegetated, and produced flowering plants: it is not new to this country, being known to MORISON who figures it, and to MILLER, who cultivated it in 1731.

It is a hardy perennial plant, growing to about the height of a foot and a half, or two feet, and flowering from the beginning of July, to September; it has a creeping root, thereby it increases greatly in light dry soils, and warm situations, so as even to be troublesome; it will not thrive in a wet soil; with us it produces seed-vessels but rarely; is propagated by parting its roots in

Autumn or Spring; MILLER recommends March as the most proper season, or it may be raised from seeds, which in certain situations and seasons ripen here.

The flowers of this *Apocynum* have a sweet honey-like fragrance, which perfumes the air to a considerable distance, and no doubt operates powerfully in attracting insects; when a plant of this sort is fully blown, one may always find flies caught in its blossoms, usually by the trunk, very rarely by the leg; sometimes four, or even five, which is the greatest possible number, are found in one flower, some dead, others endeavouring to disentangle themselves, in which they are now and then so fortunate as to succeed; these flies are of different species, the *musca pipiens*, a slender variegated fly with thick thighs, is a very common victim, the *musca domestica*, or house fly, we have never observed among the captives.

Previous to our explaining the manner in which it appears to us that these insects are caught, it will be necessary that we should describe, in as plain a manner as possible, those parts of the flower which more particularly constitute this fatal fly trap.

On looking into the flower we perceive five Stamina, the Antheræ of which are large, of a yellow colour, and converge into a kind of cone; each of these Antheræ is arrow-shaped, towards the top of the cone their sides touch but do not adhere, below they separate a little, so as to leave a very narrow opening or slit between each, they are placed on very short filaments, which stand so far apart that a considerable opening is left between them, which openings, however, are closed up by processes of the corolla, nicely adapted to, and projecting into them; at the bottom of, and in the very centre of the flower, we perceive two germina, or seed-buds, the rudiments of future seed-vessels, surrounded by glandular substances, secreting a sweet liquid; on the summit of these germina, and betwixt the two, stands the stigma, in the form of a little urn, the middle of which is encircled by a glandular ring, which secretes a viscid honey-like substance, to this part of the stigma the Antheræ interiorly adhere most tenaciously, so as to prevent their separation unless considerable force be applied; it is, as we apprehend, the sweet viscid substance thus secreted by the stigma, within the Antheræ, which the fly endeavours to obtain, and to this end insinuates its trunk first into the lowermost and widest part of the slit, betwixt each of the Antheræ above

described, pushing it of necessity upwards: when gratified, not having the sense to place itself in the same position as that in which it stood when it inserted its trunk, and to draw it out in the same direction downwards, unfortunately for it, it varies its position, and pulling its trunk upwards, draws it into the narrow part of the slit, where it becomes closely wedged in, and the more it pulls the more securely it is caught, and thus this heedless insect, as THOMSON calls it, terminates its existence in captivity most miserable.

In the incomparable poem of Dr. DARWIN, entitled the *Botanic Garden*, there is a figure given of this plant; and in the Supplement we have the following account written by Mr. DARWIN, of Elston.

"In the Apocynum Androsæmifolium the Anthers converge over the nectaries, which consist of five glandular oval corpuscles, surrounding the germ, and at the same time admit air to the nectaries at the interstice between each anther; but when a fly inserts its proboscis between these anthers to plunder the honey, they converge closer, and with such violence as to detain the fly, which thus generally perishes."

This explanation of a phænomenon entitled to much attention, is widely different from ours; which of the two is most consonant to truth and nature, we shall leave to the determination of future observers.

In explaining the preceding appearances, to prevent confusion we called those parts which form the cone in the middle of the flower Antheræ, but strictly speaking they are not such, the true Antheræ being situated on the inside of their summits, where they will be found to be ten in number, making in fact the Apocynum a decandrous plant.

[281]

TURNERA ANGUSTIFOLIA. NARROW-LEAV'D TURNERA.

Class and Order.

PENTANDRIA TRIGYNIA.

Generic Character.

Cal. 5-fidus, infundibuliformis, exterior 2-phyllus. *Petala* 5 calyci inserta. *Stigmata* multifida. *Caps.* 1-locularis, 3-valvis.

Specific Character and Synonyms.

TURNERA *angustifolia* floribus sessilibus petiolaribus, foliis lanceolatis rugosis acuminatis. *Mill. Dict. ed. 6. 4to.*

TURNERA frutescens folio longiore et mucronato. *Mart. Cent. 49. t. 49.*



Nº281.

[View larger image](#)

This plant here represented is generally known to the Nurserymen about London as the *Turnera ulmifolia*, or *Elm-leav'd Turnera*, its foliage however does not answer to the name, nor to the figures of the plant as given by MARTYN in his *Cent. Pl.* and LINNÆUS in his *Hortus Cliffortianus*, which figures indeed are so similar that they look like copies of each other, these represent the true elm leaf; on the same plate of *Martyn's Cent.* there is given a very excellent figure of what he considers as another species of *Turnera*, vide *Synon.* and which MILLER, who cultivated it about the year 1773, also describes as a distinct species, under the name of *angustifolia*, asserting, from the experience of thirty years, that plants raised from its seeds have constantly differed from those of the *ulmifolia*; this is our plant, which on his authority we have given as a species, though LINNÆUS regards it as a variety.

PLUMIER gave to this genus the name of *Turnera*, in honour of Dr. WILLIAM TURNER, a celebrated English Botanist and Physician, who published an

Herbal, black letter, folio, in 1568.

The present species is a native of the West-Indies, and is commonly cultivated in our stoves, where it rises with a semi-shrubby stalk, to the height of several feet, seldom continuing more than two or three years; young plants generally come up in plenty from seeds spontaneously scattered, so that a succession is easily obtained.

It flowers from June to August.

Its foliage has a disagreeable smell when bruised; its flowers are shewy, but of short duration, and are remarkable for growing out of the footstalk of the leaf.

HEDYSARUM OBSCURUM. CREEPING-ROOTED HEDYSARUM.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cor. carina transverse obtusa. *Legumen* articulis 1-spermis.

Specific Character and Synonyms.

HEDYSARUM *obscurum* foliis pinnatis, stipulis vaginalibus, caule erecto flexuoso, floribus pendulis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 676. Mant. 447. Jacq. Fl. Austr. v. 2. t. 168.*

HEDYSARUM caule recto, ramoso; foliis ovatis; siliquis pendulis, lævissimis, venosis. *Hall. Hist. Helv. n. 395.*

ONOBRYCHIS semine clypeato lævi. *Bauh. Pin. 350.*



Nº282.

[View larger image](#)

Prof. JACQUIN, in the second volume of the *Flora austriaca*, gives an excellent figure and accurate description of our plant, a native of the Alps of Germany and Switzerland, and points out the characters in which it differs from the *alpinum*, for which it has sometimes been mistaken.

It is a hardy perennial, rarely exceeding a foot in height, produces its spikes of pendulous flowers, which are of a most beautiful purple colour, in July and August; hitherto these have not been succeeded by seed-vessels with us; though we have cultivated the plant for several years.

Its size renders it a suitable plant for rock-work, on which it will grow readily, increasing by its roots, which are of the creeping kind.

HALLER mentions a variety of it with white flowers.

MIMULUS RINGENS. NARROW-LEAVED MONKEY-FLOWER.

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Cal. 4-dentatus, prismaticus. *Cor.* ringens; labio superiore lateribus replicato. *Caps.* 2-locularis, polysperma.

Specific Character and Synonyms.

MIMULUS *ringens* erectus, foliis oblongis linearibus sessilibus.
Linn. Syst. Vegetab. ed. 14. Murr. p. 575. Ait. Kew. v. 2. p. 361.

EUPHRASIA *floridana* *lysimachiae* glabræ siliquosæ foliis, quadrato caule ramosior. *Pluk. Amalth. 83. t. 393. f. 3.*

LYSIMACHIA *galericulata* s. *Gratiola elatior* non ramosa, &c.
Gron. Fl. Virg. p. 97.

DIGITALIS *perfoliata* glabra flore violaceo minore. *Moris. Hist. 2. p. 479. s. 5. t. 8. f. 6.*



Nº283.

[View larger image](#)

CLAYTON, in the *Fl. Virg.* published by GRONOVIVS, describes this plant as a native of Virginia, and says of it, "maddidis gaudet locis," it delights in wet places: LINNÆUS makes it a native of Canada also.

It is a hardy, perennial, herbaceous plant, growing with us to the height of about two feet, and producing its flowers, which are of a pale violet colour, in July and August; these are frequently succeeded by capsules containing perfect seeds, by which the plant may be propagated, as also by parting its roots in Autumn; MILLER recommends the seeds to be sown as soon as ripe.

The plant succeeds best in a moist and somewhat shady situation, with a loamy soil.

A perusal of the synonyms will shew to what a variety of genera this plant has been referred by different authors; LINNÆUS first gave to it the name of *Mimulus*, of which term we find in his *Philosophia Botanica* the following concise explanation:—"MIMULUS mimus personatus;" in plain English, a

masked mimick: MIMMULUS is a classical word for the Pedicularis, or Lousewort; the English term Monkey flower has probably been given it, from an idea that *mimulus* originated from μιμω a monkey, as in *mimusops* monkey face.

[284]

ROSA SEMPERFLORENS. EVER-BLOWING ROSE.

Class and Order.

ICOSANDRIA POLYGYNIA.

Generic Character.

Petala 5. *Cal.* urceolatus, 5-fidus, carnosus, collo coarctatus, demum baccatus, coloratus. *Antrum* duplicatum, 1-loculare, superne apertum, pericarpis osseis intus nidulantibus.

Specific Character.

ROSA semperflorens caule aculeato, foliis subternis, pedunculis subunifloris aculeato-hispidis, calycis laciniis integris.



Nº284.

[View larger image](#)

We are induced to consider the rose here represented, as one of the most desirable plants in point of ornament ever introduced to this country; its flowers, large in proportion to the plant, are semi-double, and with great richness of colour unite a most delightful fragrance; they blossom during the whole of the year, more sparingly indeed in the winter months; the shrub itself is more hardy than most greenhouse plants, and will grow in so small a compass of earth, that it may be reared almost in a coffee cup; is kept with the least possible trouble, and propagated without difficulty by cuttings or suckers.

For this invaluable acquisition, our country is indebted to the late GILBERT SLATER, Esq. of Knots-Green, near Laytonstone, whose untimely death every person must deplore, who is a friend to improvements in ornamental gardening: in procuring the rarer plants from abroad, more particularly from the East-Indies, Mr. SLATER was indefatigable, nor was he less anxious to have them in the greatest perfection this country will admit; to gain this point there was no contrivance that ingenuity could suggest, no labour, no expence withheld; such exertions must soon have insured him the first collection of the plants of India: it is now about three years since he obtained this rose from China; as he readily imparted his most valuable acquisitions to those who were most likely to increase them, this plant soon became conspicuous in the collections of the principal Nurserymen near town, and in the course of a few years will, no doubt, decorate the window of every amateur.

The largest plants we have seen have not exceeded three feet, it may no doubt be trained to a much greater height; a variety of it much more robust, having usually several flowers on a footstalk, of a pale red colour, and semi-double also, has more lately been introduced, and as far as we can learn from the same source.

[285]

JASMINUM ODORATISSIMUM. SWEETEST JASMINE.

Class and Order.

DIANDRIA MONOGYNIA.

Generic Character.

Corolla hypocrateriformis. *Bacca* dicocca. *Semina* solitaria, arillata.

Specific Character and Synonyms.

JASMINUM *odoratissimum* foliis alternis obtusiusculis ternatis pinnatisque, ramis teretibus, laciniis calycinis brevissimis. *Ait. Hort. H. v. 1. p. 10. Linn. Syst. Veget. ed. 14. Murr. p. 56.*

JASMINUM flavum odoratum. *Barr. Ic. 62.*



Nº285.

[View larger image](#)

The flowers of most of the species of Jasmine are odoriferous, trivial names therefore expressive of this quality are ineligible, as wanting character; the present name is peculiarly objectionable, inasmuch as several other species are greatly superior to this in point of fragrance; a lesson for Botanists to abstain from trivial names of the superlative degree, such as *odoratissimum*, *fœtidissimum*, *maximum*, *minimum*, &c.

The present species, according to Mr. AITON, is a native of Madeira, and was cultivated by Mr. MILLER, in 1730; it is now a plant common in most greenhouses: it will form a shrub of considerable size, which requires no support; its leaves are glossy, inclining to yellow, growing for the most part three together, sometimes pinnated; its blossoms, which are yellow, make their appearance from May to November: in point of hardiness it is superior to many greenhouse plants, and may be propagated without difficulty by cuttings.

[286]

PORTLANDIA GRANDIFLORA. GREAT-FLOWERED PORTLANDIA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. clavato-infundibuliformis. *Antheræ* 4-6. longitudinales.
Caps. 5-gona, 2-valvis, retusa, 2-locularis, polysperma,
coronata calyce 5-phyllo.

Specific Character and Synonyms.

PORTLANDIA *grandiflora* floribus pentandris. *Linn. Syst. Veg.*
ed. 14. Murr. p. 213. Ait. Kew. v. 1. p. 228. foliis ovatis.
Syst. Nat. ed. 13. Gmel. p. 360.

PORTLANDIA *grandiflora* floribus pentandris, capsulis ovatis,
foliis oblongis acuminatis. *Swartz. Obs. Bot. p. 69.*



Nº286.

[View larger image](#)

Dr. BROWN, in his Natural History of Jamaica, gives to this genus the name of *Portlandia*, in honour of the Duchess Dowager of PORTLAND, who employed many of the leisure hours of a long and happy life, in the pursuits of natural history, in which she was eminently skilled.—She was the friend and patron of Mr. LIGHTFOOT, who dedicates to her his *Flora Scotica*; the fine collection of rare and valuable trees and shrubs which enrich part of the grounds at Bulstrode, were of her planting.

Dr. SWARTZ, in his Observations on the Plants of the West-Indies, informs us, that this species grows wild in Jamaica, where (*incolit calcareosa petrosa*) it inhabits calcareous rocky places^[3], forms a small tree about the height of six feet, and flowers from the middle of Summer to Autumn; its bark, he observes, as in other plants of the same genus, is extremely bitter.

From Mr. AITON we learn, that it was introduced here by ——— ELLIS, Esq. in 1775.

It forms a very beautiful stove plant, not of difficult growth, and readily disposed to flower; we have seen blowing plants of it little more than a foot high; its blossoms are not only uncommonly large, shewy, and curious in their structure, but fragrant also, and very much so when dried.

It is usually increased by cuttings.

[287]

GOODENIA LÆVIGATA. SMOOTH GOODENIA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Flores monopetali, superi. *Caps.* bilocularis. *Cor.* supra longitudinaliter fissa, stigma urceolatum ciliatum. *Smith Trans. Linn. Soc. v. 2. p. 346.*

Specific Character.

GOODENIA lævigata foliis obovato-lanceolatis dentatis glabris.



N^o287.

[View larger image](#)

In the Autumn of 1792, SAMUEL TOLFREY Esq. most kindly invited me to inspect a vast number of the natural productions of Botany-Bay, in his possession; collected with great assiduity, and brought over in high preservation by Captain TENCH; among other curiosities, he shewed me specimens of the earths of that country, imported in very small bags. I suggested to Mr. TOLFREY, that those earths might possibly contain the seeds of some curious and unknown plants; he readily acquiesced in the idea, and permitted me to make trial of them: accordingly, in the Spring of 1793, I exposed them in shallow pans, on a gentle tan heat, keeping them duly watered; in the course of the Summer they yielded me fourteen plants, most of which were altogether new, and among others the species of *Goodenia* here figured; this we have since found to be a hardy greenhouse plant, flowering from July to October, and very readily increased by cuttings.

The oldest plant in our possession is about a foot and a half high, much branched, the stalks are round and smooth to the naked eye, green below, above purplish, the leaves are smooth, a deep bright green colour, alternate, standing on footstalks, which gradually widen into the leaves, somewhat ovate, and deeply toothed; the flowers grow in the alæ of the leaves, forming a thin spike, they are sessile, of a pale violet colour, and have a peculiar smell which is rather unpleasant; at the side of each flower are two long narrow Bracteæ; the Calyx, which is placed on the germen, is composed of five short ovate leaves, which appear edged with hairs if magnified; the Corolla is monopetalous, the lower part, which at first is tubular, splits longitudinally above, and forms a kind of half tube, the edges of which are brown, the inside yellow, the outside greenish, the mouth beset with short hairs, each of which is terminated by a small villous head; the limb is deeply divided into five linear segments, spreading out like a hand, and terminated by short points; the Filaments are five in number, of a whitish colour, somewhat broadest above, rather flat, inserted into the receptacle; Antheræ oval, flattened, yellow, bilocular, a little bent, the length of the pistillum; but this is to be understood of such flowers as are not yet fully expanded, in those that are, they are much shorter, and appear withered; the Style, in flowers about to open, the length of the filaments, upright, in those that are opened much longer, and bent somewhat downward; Stigma at first upright, in the form of a cup, having the edge curiously fringed with white hairs, afterwards it closes together, loses its

hollow, and assumes a flat appearance, and nods somewhat, the back part of it is bearded; Germen beneath the calyx, oblong, usually abortive with us.

The name of *Goodenia* has been given to this genus by Dr. SMITH, in honour of the Rev. SAMUEL GOODENOUGH, LL. D. of *Ealing*, my much-honoured friend, whose name will be ever dear to Botanists for his laborious investigation of the British Carices^[4].

[288]

PASSIFLORA CILIATA. FRINGED-LEAVED PASSION- FLOWER.

Class and Order.

GYNANDRIA HEXANDRIA.

Generic Character.

Cal. 5-phyllus. *Petala* 5. *Stamina* germini vicina. *Nectarium* multi-radiatum. *Antrum* pedicellatum duplicatum 1-loculare.

Specific Character and Synonyms.

PASSIFLORA *ciliata* foliis trilobis glabris ciliato serratis intermedio longissimo, petiolis eglandulosis. *Ait. Kew. v. 3. p. 310.*



N^o288.

[View larger image](#)

This Passion-Flower is described in the *Hort. Kew.* as a new one, under the name of *ciliata*, introduced by Mrs. NORMAN, from the West-Indies, in 1783: we saw it during the latter part of the last Summer, with great profusion of flowers, in several collections, more particularly in that of Mr. VERE, Kensington-Gore, from whence our figure and description were taken.

Its stalks are round, perfectly smooth, and run to a very great height; leaves dark green, glossy, perfectly smooth, except on the edges, where they are beset with strong glandular hairs, divided into three large and two small lobes, the middle lobe running out to a considerable length, the footstalks of the leaves are beset with a few hairs thinly scattered, at the base of each leaf is a tendril, and two finely-divided stipulæ, edged also with glandular hairs. The Involucrum is composed of three leaves, dividing into capillary segments, each of which terminates in a viscid globule, fetid when bruised; betwixt the involucrum and the blossom is a short peduncle; the pillar which supports the germen is of a bright purple colour, with spots of a

darker hue, the germen is smooth and green; Styles green; Stigmata of a dark green; Filaments six in number; Antheræ pale yellow green, the former dotted with purple; of Radii, there may be said to be four rows, variegated with white and purple, petals ten, externally greenish, internally red, deeper or paler according to circumstances.

The leaves of this plant vary greatly in form, according to the health and luxuriance of the plant; on comparing it with the *fætida*, we strongly suspect it to be a variety merely of that species: time will shew.

It is increased by cuttings, or seeds.

INDEX.

In which the Latin Names of the Plants contained in the *Eighth Volume* are alphabetically arranged.

Pl.

[280](#) *Apocynum androsæmifolium*.

[276](#) *Blitum virgatum*.

[274](#) *Borbonia crenata*.

[264](#) *Cistus formosus*.

[258](#) *Coronilla varia*.

[271](#) *Cyrtanthus angustifolius*.

[255](#) *Cytisus sessilifolius*.

[273](#) *Diosma uniflora*.

[261](#) *Erodium incarnatum*.

[272](#) *Gladiolus tristis*.

[263](#) *Glycine bimaçulata*.

[268](#) *Glycine rubicunda*.

[270](#) *Glycine coccinea*.

[287](#) *Goodenia lævigata*.

[282](#) *Hedysarum obscurum*.

[285](#) *Jasminum odoratissimum*.

[256](#) *Ixia longiflora*.

[265](#)—— *Bulbocodium*.

[253](#) *Lathyrus articulatus*.

[259](#) *Lilium Catesbæi*.

[278](#) *Lilium candidum*.

[275](#) *Liriodendron Tulipifera*.

[254](#) *Lopezia racemosa*.

[257](#) *Lychnis chalcedonica*.

[277](#) *Mahernia pinnata*.

[262](#) *Mesembryanthemum aureum*.

[260](#) *Metrosideros citrina*.

[283](#) *Mimulus ringens*.

[269](#) *Ornithogalum nutans*.

[288](#) *Passiflora ciliata*.

[279](#) *Plumeria rubra*.

[286](#) *Portlandia grandiflora*.

[267](#) *Pyrus spectabilis*.

[266](#) *Ranunculus amplexicaulis*.

[284](#) *Rosa semperflorens*.

[281](#) *Turnera angustifolia*.

INDEX.

In which the English Names of the Plants contained in the *Eighth Volume* are alphabetically arranged.

Pl.

- [267](#) Apple-tree Chinese.
- [276](#) Blite strawberry.
- [274](#) Borbonia heart-leaved.
- [264](#) Cistus beautiful.
- [272](#) Corn-flag square-leaved.
- [258](#) Coronilla purple.
- [261](#) Crane's-bill flesh-coloured.
- [266](#) Crowfoot plantain-leaved.
- [271](#) Cyrtanthus narrow-leaved.
- [255](#) Cytisus common.
- [273](#) Diosma one-flowered.
- [280](#) Dogsbane tutsan-leaved.
- [262](#) Fig-marigold golden.
- [263](#) Glycine purple.
- [268](#) Glycine dingy-flowered.
- [270](#) Glycine scarlet.
- [287](#) Goodenia smooth.
- [282](#) Hedysarum creeping-rooted.
- [285](#) Jasmine sweet.
- [256](#) Ixia long-flowered.
- [265](#) Ixia crocus-leaved.
- [253](#) Lathyrus jointed-podded.
- [259](#) Lily Catesby's.
- [278](#) Lily white.
- [254](#) Lopezia mexican.
- [257](#) Lychnis scarlet.

[277](#) Mahernia winged.
[260](#) Metrosideros harsh-leaved.
[283](#) Monkey-flower narrow-leav'd.
[288](#) Passion-flower fringed-leaved.
[279](#) Plumeria red.
[286](#) Portlandia great-flowered.
[284](#) Rose ever-blowing.
[269](#) Star of Bethlehem Neapolitan.
[275](#) Tulip-tree common.
[281](#) Turnera narrow-leaved.

[1] In honorem Licent. THOMÆ LOPEZ, Burgensis, qui aliquot annos Regii Senatoris munere functus in America, CAROLO V. imperante. In patriam reversus brevium historiæ naturalis novi orbis scripsit sub titulo de tribus elementis aëre, aqua, et terra, MS. apud eundem Muguoziū.

[2] What the use of this very extraordinary apparatus may be we can at present scarcely conjecture, future observation may perhaps enable us to speak more decisively; when we figure the *Diosma ericoides* we shall probably have more to say of this species.

[3] We wish that every person who describes foreign plants on the spot, would do thus; it would greatly facilitate their culture.

[4] Vide a Dissertation on the British species of Carex, by Dr. GOODENOUGH, in the second volume of the Transactions of the Linnean Society.

*** END OF THE PROJECT GUTENBERG EBOOK THE BOTANICAL
MAGAZINE, VOL. 08 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



Ranunculus acris L.

[back](#)



[back](#)




[back](#)



[back](#)



[back](#)



[back](#)




[back](#)



[back](#)




[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)