

THE
BOTANICAL MAGAZINE;
OR,
Flower-Garden Displayed :

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the
Open Ground, the Green-House, and the Stove, are
accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according
to the celebrated LINNÆUS; their Places of Growth,
and Times of Flowering :

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A W O R K

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as
wish to become scientifically acquainted with the Plants they cultivate.

By *WILLIAM CURTIS*,
Author of the *FLORA LONDINENSIS*.

VOL. XIII.

THE
BOTANICAL MAGAZINE;
OR,
Flower-Garden Displayed :

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the
Open Ground, the Green-House, and the Stove, are
accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according
to the celebrated LINNÆUS; their Places of Growth,
and Times of Flowering :

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A W O R K

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as
wish to become scientifically acquainted with the Plants they cultivate.

By *WILLIAM CURTIS*,
Author of the *FLORA LONDINENSIS*.

VOL. XIII.

The Project Gutenberg eBook of The Botanical Magazine, Vol.

13

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The Botanical Magazine, Vol. 13

Author: William Curtis
John Sims

Release date: October 1, 2013 [eBook #43858]
Most recently updated: October 23, 2024

Language: English

Other information and formats: www.gutenberg.org/ebooks/43858

Credits: Produced by Marc-André Seekamp, Jason Isbell and the Online Distributed Proofreading Team at <http://www.pgdp.net> (This file made using scans of public domain works at the University of Georgia.)

*** START OF THE PROJECT GUTENBERG EBOOK THE
BOTANICAL MAGAZINE, VOL. 13 ***

**THE
BOTANICAL MAGAZINE;
OR,
Flower-Garden Displayed:**

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the Open Ground, the Green-House, and the Stove, are accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according to the celebrated LINNÆUS; their Places of Growth, and Times of Flowering:

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A WORK

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as wish to become scientifically acquainted with the Plants they cultivate.

By WILLIAM CURTIS,

Author of the FLORA LONDINENSIS.

VOL. XIII.

"—— All alone, amid her Garden fair,
"From morn to noon, from noon to dewy eve,
"She spent her days, her pleasing task to tend
"The flowers; to lave them from the water-spring;
"To ope the buds with her enamoured breath,

"Rank the gay tribes, and rear them in the sun. —— ——

"Thus plied assiduous her delightful task,

"Day after day, till every herb she named

"That paints the robe of Spring."

BRUCE.

LONDON:

Printed by STEPHEN COUCHMAN, Throgmorton-Street,

For W. CURTIS, N^o 3, *St. George's-Crescent*, Black-Friars-Road;

And Sold by the principal Booksellers in Great-Britain and Ireland.

M DCC XCIX.

CONTENTS

- [433]—[AZALEA PONTICA.](#)
- [434]—[OXYBAPHUS VISCOSUS.](#)
- [435]—[GNAPHALIUM ERICOIDES.](#)
- [436]—[HIBISCUS PRÆMORSUS.](#)
- [437]—[HYDRANGEA ARBORESCENS.](#)
- [438]—[HYDRANGEA HORTENSIS.](#)
- [439]—[ILLCIUM FLORIDANUM.](#)
- [440]—[ERICA ALBENS.](#)
- [441]—[ANTHOLYZA MERIANELLA.](#)
- [442]—[GENISTA LINIFOLIA.](#)
- [443]—[ERICA PHYSODES.](#)
- [444]—[CANARINA CAMPANULA.](#)
- [445]—[CONVOLVULUS CNEORUM.](#)
- [446]—[PSORALEA BRACTEATA.](#)
- [447]—[ERICA EMPETRIFOLIA.](#)
- [448]—[MESEMBRYANTHEMUM MICANS.](#)
- [449]—[DILLENIA SPECIOSA.](#)
- [450]—[GLADIOLUS WATSONIUS.](#)
- [451]—[BLAKEA TRINERVIA.](#)
- [452]—[CARDAMINE TRIFOLIA.](#)
- [453]—[AMARYLLIS REGINÆ.](#)
- [454]—[CANNA INDICA.](#)
- [455]—[ALÖE RETUSA.](#)
- [456]—[DIOSMA SERRATIFOLIA.](#)
- [457]—[ALÖE PLICATILIS.](#)
- [458]—[ARISTEA CYANEA.](#)
- [459]—[CONVOLVULUS CNEORUM.](#)
- [460]—[MAURANDYA SEMPERFLORENS.](#)
- [461]—[JASMINUM FRUTICANS.](#)

[\[462\]—ANTHEMIS PYRETHRUM.](#)
[\[463\]—EPIDENDRUM CILIARE.](#)
[\[464\]—SISYRINCHIUM GRAMINEUM.](#)
[\[465\]—INDIGOFERA ANGUSTIFOLIA.](#)
[\[466\]—CALYCANTHUS PRÆCOX.](#)
[\[467\]—DRACOCEPHALUM](#)
[VIRGINIANUM.](#)
[\[468\]—ŒNOTHERA TETRAPTERA.](#)
[INDEX.—Latin Names](#)
[INDEX.—English Names](#)

[433]

AZALEA PONTICA. YELLOW AZALEA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. campanulata. *Stamina* receptaculo inserta. *Caps.* 2-5
locularis polysperma.

Specific Character and Synonyms.

AZALEA pontica foliis nitidis lanceolatis, utrinque glabris,
racemis terminalibus. *Linn. Sp. Pl. App. p. 1669. Pall.*
Fl. Ross. t. 69. p. 51.

CHAMÆRHODENDROS Pontica maxima Mespili folio flore
luteo. *Tourn. Cor. Hist. Rei. Herb. 42. Act. Paris 1704. p.*
348. Buxb. Cent. 5. p. 36. t. 69.



Nº433.

DESCR. Shrub from two to three feet or more in height, the thickest part of the stem not exceeding the size of the little finger, covered with a smooth brown bark, irregularly branched; Flowers appearing before the leaves are fully expanded, and produced in umbels at the extremities of the branches, from eight to twelve or more in an umbel, of a fine yellow colour and agreeable fragrance; each blossom is about the size of that of the horse-chestnut, and as some of them are produced much earlier than others, the plant of course continues a considerable time in bloom, standing on short peduncles; Calyx very short, viscous, and irregularly divided, most commonly into five ovato-lanceolate segments; Corolla, tube cylindrical, viscous, grooved, brim divided into five segments, undulated and somewhat wrinkled, ovate, pointed, three turning upwards, two downwards, of the three uppermost segments the middle one more intensely yellow than the others and inclining to orange, with which it is sometimes spotted; Stamina usually five, yellow, projecting beyond the corolla, and turning upwards near their extremities; Antheræ orange-coloured; Pollen whitish and thready; Germen somewhat conical, evidently hairy, and somewhat angular;

Style yellowish, filiform, projecting beyond the stamina, and turning upwards; Stigma forming a round green head.

The figure and description here given were taken from a plant which flowered by means of artificial heat, in the spring of 1798, at Mr. WATSON'S, Nurseryman, Islington, and which had been introduced the same year, by Mr. ANTHONY HOVE, of Warsaw.

As an hardy ornamental shrub, it bids fair to prove an acquisition truly valuable, its flowers produced in the months of June and July, being highly ornamental as well as fragrant.

We have the best authority for regarding this plant as the *Chamærhodendros Pontica* of TOURNEFORT, it agrees with his own specimens in Sir JOSEPH BANKS'S Herbarium, it accords also with his description, and figures, more especially of those flowers which are of their natural size; nor have we any doubts of its being the *Azalea Pontica* of Prof. PALLAS, figured in the *Flor. Rossica*, since it corresponds generally with his description, though not in all points with his figure, which bears evident marks of inaccuracy, the stigma, for instance, is represented as trifid.

TOURNEFORT found this plant on the eastern side of the Black Sea, Mr. HOVE on the north side near Oczakow, and elsewhere; Professor PALLAS on Mount Caucasus.

As yet there has been no opportunity of ascertaining the best means of propagating this new denizen, but there is every reason to suppose that it will succeed with the treatment bestowed on the other *Azalea*'s.

Prof. PALLAS relates that the honey of bees frequenting the flowers of this plant is supposed to be narcotic, and that goats, kine, and sheep on eating its leaves have been poisoned thereby.

By permission and with the approbation of Mr. ANTHONY HOVE, the following extracts from his journal are here inserted.

"June 9, 1796, found a few of this species of *Azalea* on the river Dnieper in swampy ground, four feet high, beginning to blow, called here the stupifying shrub, and considered by some as highly efficacious in curing the venereal disease.

"June 20, found this species on the river Dniester, on the estate of Count STANISLAUS SCZESNY POTOCKI, about sixteen English miles from the town of Mohilow, in peat earth, from four to twenty feet high; regarded by the common people as intoxicating, and used in the cure of various diseases.

"July 4, near Oczakow, found thousands of these plants fully blown, in a marsh, every spring-tide overflowed by the sea; found there also, a Tartarian farmer, who lived entirely by the profits arising from the honey which the bees extracted from the flowers of this plant, sold to Constantinople and other parts of Turkey for medicinal uses.

"July 15, arrived at Trebizond, found a valley about ten English miles from the sea covered with these plants."

On cultivation, Mr. HOVE relates that he found the *Azalea's* from Trebizond much more tender than those from the borders of the Dnieper and the Dniester, and was therefore inclined to regard them as strong varieties if not distinct species.

The leaves, when fully expanded, are in size and figure like those on the plate, hairy on both sides, and terminating in a very remarkable blunt mucro or point, which has not been noticed either by those who have described or figured the plant as it deserves, for it appears to form a very strong character.

[434]

OXYBAPHUS VISCOSUS. VISCID UMBRELLA-WORT.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Cal. 5-fidus campanulatus. *Cor.* infundibuliformis. *Nux* 5-gona
1-sperma calyce explanato persistenti circumdata.

Specific Character and Synonyms.

OXYBAPHUS *viscosus*. *L'Herit. Monograph.*

MIRABILIS *viscosa* floribus racemosis; foliis cordatis
orbiculato-acutis tomentosis. *Cav. Icon. I. n. 17. t. 19.*



Nº434.

The present is one of those plants which is entitled to our admiration, for its curious and singular structure, rather than for any extraordinary figure it makes in a collection; not that its flowers are without a certain share of beauty.

It is an annual, growing to about the height of two feet; the whole plant is viscous, and, if bruised, smells somewhat disagreeably; the flowers do not open at any particular time as those of the *Mirabilis* do, and each contains only three stamina; when they fall off, which they do soon after expanding, the calyx closes on the germen, enlarges, droops, and becomes deeply plaited; on the ripening of the seed it turns brown, expands, and is suspended like a little umbrella over the seed, which when perfectly ripe drops out on the ground; the expanded calyx in this state appears somewhat like the flower of a *Physalis*.

This plant flowers from June to October, and ripens its seeds in the open air; it is not difficult of culture, requires the same treatment as other tender annuals from Peru.

Our figure was drawn from a plant which flowered 1796, in the collection of the Marchioness of BUTE, and was raised from Peruvian seeds, sent her by Prof. ORTEGA of Madrid, under the name of *Mirabilis triandra*; Mons. CAVANILLE has figured and described it as the *Mirabilis viscosa*; Mons. L'HERITIER, from a consideration of all its characters, has been induced to make a new genus of it, which he has called *Oxybaphus*.

[435]

GNAPHALIUM ERICOIDES. HEATH-LEAVED

GNAPHALIUM, or EVERLASTING.

Class and Order.


SYNGENESIA POLYGAMIA SUPERFLUA.

Generic Character.

Recept. nudum. *Pappus* plumosus. *Cal.* imbricatus: squamis marginalibus rotundatis scariosis coloratis.

Specific Character and Synonyms.

GNAPHALIUM *ericoides* fruticosum foliis sessilibus linearibus, calycibus exterioribus rudibus, interioribus incarnatis. *Linn. Am. Acad.* v. 6. p. 99. *Syst. Vegetab. ed.* 14. *Murr.* p. 746. *Ait. Hort. Kew.* p. 174.



Nº435.

In the 6th vol. of the *Amœnitates Academicæ*, LINNÆUS describes this species of *Gnaphalium*, most probably from dried specimens collected at the Cape, where it is a native; so insignificant did the plant appear to him, that in his description, he calls it *misera*; compared with the more magnificent species, such a term might not perhaps be inapplicable: but, though small, the plant possesses much beauty when cultivated, and hence is generally kept in our green-houses.

It flowers from March to August.

Its branches, naturally weak and trailing, require to be carefully tied up; if this business be executed with taste and judgment, the natural beauty of the plant may be considerably heightened.

It is readily increased by cuttings.

Was introduced to the royal garden at Kew, by Mr. MASSON, in 1774.

[436]

HIBISCUS PRÆMORSUS. BITTEN-LEAVED HIBISCUS.

Class and Order.

MONADELPHIA POLYANDRIA.

Generic Character.

Cal. duplex: exterior polyphyllus. *Caps.* 5-locularis, polysperma.

Specific Character and Synonyms.

HIBISCUS *præmorsus* foliis subrotundis, dentato-serratis retusis pubescentibus. *Ait. Kew. v. 2. 454.*

HIBISCUS *præmorsus* hirsutus, foliis ovatis basi angustato-cordatis apice præmorsis crenatis, calycibus tomentosis, seminibus tuberculatis. *Linn. Suppl. Pl. p. 309.*

PAVONIA *cuneifolia*. *Cavan. Diff. 3. p. 139. t. 45. f. 1.*

URENA *præmorsa*. *L'Herit. Stirp. nov. t. 51.*



N^o436.

Though not so shewy a plant as many of the genus, we find this species of *Hibiscus* in most collections of green-house plants about London, flowering from June to September.

Its foliage is singular, its blossoms sulphur coloured with a tint of orange on the under side and without scent; if suffered to grow, this shrub will acquire a considerable height; it seeds freely, by which the plant is readily increased, and by these it requires to be renewed once in two or three years.

Is a native of the Cape, from whence it was introduced, by Mr. MASSON, in 1774. *Ait. Kew.*

It will be seen by the synonyms, that authors have been divided in their opinions as to the genus of this plant; LINNÆUS the younger, in his *Suppl.* makes it an *Hibiscus*; CAVANILLE, a *Pavonia*; L'HERITIER, an *Urena*; Mr. AITON, an *Hibiscus*; not partial to the multiplying of genera, unless there be an obvious necessity for it, we have in the present instance followed the first and last of these writers.

[437]

HYDRANGEA ARBORESCENS. SHRUBBY HYDRANGEA.

Class and Order.

DECANDRIA DIGYNIA.

Generic Character.

Caps. 2-locularis, 2-rostris, infera, foramine inter stylos dehiscens.

Specific Character and Synonyms.

HYDRANGEA *arborescens* caule arboreo. *Linn. Syst. Nat. ed. 13. Gmel. Syst. Veget. ed. 14. Murr. p. 410. Ait. Kew. v. 2. p. 76.*

ANONYMOS floribus albis parvis, in umbella lata magna dispositis odoratis, foliis amplis acuminatis serratis pediculis insidentibus ex adverso binis, caule fruticoso præalto non ramoso, vasculo parvo, bicapsulari, seminibus minutissimis repleto, duobus parvis filamentis seu corniculis recurvis coronato. *Clayt. n. 79. Mill. Icon. t. 251. Duham. Arb. 1. t. 3.*



Nº437.

As a shrub commonly cultivated for ornament in our gardens and plantations, the *Hydrangea arborescens* has a right to appear in this work, nevertheless it is more with a view to illustrate a peculiar character of the genus that it is here figured.

This shrub is a native of Virginia, and was introduced in 1736, by P. COLLINSON, Esq. it is of low growth, rarely exceeding four feet in height; its flowers are produced on the summits of the branches, somewhat in the manner of the *Laurustinus*; they are small, crowded, nearly white, with a tinge of red in them which is not very brilliant, and some fragrance; they are produced in July and August, and sometimes followed by ripe seeds here.

Is easily propagated by parting its roots the latter end of October, prefers a moist soil, but will grow in a dry one; if in severe frosts the stalks, which are rather soft, should be killed, new ones will be put up the following Spring.

[438]

HYDRANGEA HORTENSIS. GARDEN HYDRANGEA.

HYDRANGEA *hortensis* foliis ellipticis serratis glaberrimis
staminibus æqualibus. *Smith icon. rar. t. 12.*

HORTENSIA. *Commers. Jussieu Gen. 214.*

PRIMULA *mutabilis* caule suffruticoso multiplici, foliis ovatis
serratis, floribus nudis. *Loureir Coch. Chin. v. 1. p. 104.*

VIBURNUM *tomentosum* foliis ovatis acuminatis serratis
venosis subtus tomentosis, umbellis lateralibus. *Thunb.*
Jap. p. 123. ?

SAMBUCUS *aquatica* surculis pinguibus punctatis, &c. Sijo
vulgo Adsai et Ansai et Adsiki. *Kæmpf. Am. Exot. p. 854.*
var. fl. albo, pila florida major.



N°438.

It appears to be a point not yet fully determined, whether the present plant exhibits the appearances belonging to it in a state of nature, or those which are in a certain degree the effect of accident, or of art; in its fructification it certainly is not so completely barren as the Guilder Rose, *Viburnum Opulus*, cultivated in our gardens, since it has most of its parts perfect; yet as none of the authors who have seen it in China or Japan (where it is said not only to be much cultivated but indigenous^[1]) describe its fruit, we are inclined on that account to regard it, in a certain degree, as monstrous.

It will appear by the synonyms, that authors have entertained very different opinions as to what this plant really is; JUSSIEU following COMMERSON makes it an *Hortensia*, THUNBERG a *Viburnum*, LOUREIRO, ridiculously enough, a *Primula*, and Dr. SMITH an *Hydrangea*.

In the original description of the characters of the genus *Hydrangea* by LINNÆUS, there is no mention made of two different kinds of florets, as in the *Viburnum*, nor has any author that I am acquainted with described the *Hydrangea arbor.* as producing such; yet, to my great surprise, in a plant of

this sort which flowered in my garden at Brompton in July 1797, three of the Cymæ, and three only, threw out each of them from their circumference a very different flower from those in the centre, smaller indeed, but very similar to the flowers of the *Hydrang. hort.* see Pl. 437. In 1788, Mr. WALTER published his *Fl. Carolin.* in which he describes a second species of *Hydrangea*, which he calls *radiata*,^[2] having very distinctly, as in the *Viburnum*, two different kinds of florets in the same Cyma, this variation in the florets is added by him to the generic character: the similarity which exists between the flowers of Mr. WALTER'S *Hydrangea radiata*, and those of the present plant sufficiently justify Dr. SMITH in making it an *Hydrangea*; the appearances observed by LOUREIRO^[3] on dissecting the germen, and our discovery of the existence of two different kinds of flowers in the *Hydrangea arborescens*, tend still more to confirm its propriety; we may add, that in the very habit of these several plants there exists a considerable similarity; still, however, it is only by ripe seed-vessels of the present plant, that this doubtful matter can be satisfactorily cleared up; but it will not follow, that if it be not an *Hydrangea* it must be a *Viburnum*.

This magnificent and highly ornamental plant, according to Dr. SMITH, was introduced from China to the royal garden at Kew, by Sir JOSEPH BANKS, Bart. in 1790; it was imported by Mr. SLATER about the same time, with whom it is said to have first flowered in this country.

If room were allowed us, it would be superfluous to describe minutely a plant now so very common; suffice it to say, that from a strong perennial root, rise a number of half-shrubby, irregular, somewhat spongy stalks, strongly spotted when young with purple, from one to three feet high, terminated by large bunches of flowers, at first green, then rose-coloured, and finally green a second time; these are the most common changes to which they are liable: but it will sometimes happen that a plant which has produced red flowers one year, shall produce blue another, though growing in the same pot; this we saw happen in the year 1796 to a plant in the possession of the Countess of UPPER OSSORY, whose refined taste and superior judgment have in several instances contributed to render our works more acceptable to the public: the coloured changeable part of the flower is regarded as the calyx, in the centre of which is the corolla, containing the stamina, &c. all varying greatly in point of number; besides these, there are other flowers without any calyx, but the parts which they contain do not

seem to be more perfect than those of the others, nor more productive of ripe fruit.

Since the introduction of this plant, trials have been made in regard to its hardiness, and it is found to survive mild winters if planted in very warm sheltered situations; but in others, both stalks and leaves are liable to be killed by slight frosts, though the roots are not; if persons are anxious to have it in the open border, the best mode will be to cut down the stems at the approach of winter, and cover over the root with rotten tan, or some light substance; in the spring fresh stalks will shoot forth, but it is more common to keep this plant during winter in a green-house or well secured frame; by artificial heat it may be brought to flower in April or May, without such, it begins to blossom about June, and continues in bloom till October; when successfully treated, it will acquire the height of three feet, and produce bunches of flowers supremely magnificent: such plants in pots are admirably adapted for decorating court-yards, balconies, &c. unless carefully cut in, it is apt to grow too large for the green-house, therefore it is proper to have a succession of young plants from cuttings, which strike very freely; this plant loves water, is indeed almost an aquatic, a rich soil, and plenty of pot room.

[1] *Habitat* et ob pulchritudinem colitur Cantone Sinarum, *Lourei. Coch. Chin.* v. 1. p. 104. Crescit in sylvis variis, inter Miaco et Jedo, etiam cultum, *Thunb. Fl. Jap.* p. 123, who refers to KÆMPFER, whose plant is certainly ours; yet it must be acknowledged that THUNBERG'S description does not well accord with it.

[2] This plant, or one extremely similar to it, was introduced by Mr. WILLIAMS, Nurseryman at Paris, a few years since; we saw it in full bloom, at Mr. COLVILL'S, King's-Road, in the Summer of 1796.

[3] Pericarpium abortit, quod ex dissecto germine et per microscopium viso apparet polyspermum.

[439]

**ILLICIUM FLORIDANUM. RED-FLOWERED ILLICIUM, or
ANISEED-TREE.**

Class and Order.

POLYANDRIA POLYGYNIA.

Generic Character.

Cal. 6-phyllus. *Petala* 27. *Caps.* plures, in orbem digestæ,
bivalves, monospermæ.

Specific Character and Synonyms.

ILLICIUM *floridanum* floribus rubris. *Linn. Syst. Vegetab. ed.*
14. Murr. p. 507. Ellis Act. Angl. 1770. (v. 60.) p. 524. t.
12. Ait. Kew. v. 2. p. 250.



Nº439.

Of the genus *Illicium* there are at present only two known species, viz. the *anisatum* and *floridanum*, the former a native of China and Japan, the latter of Florida; both of them are cultivated in this country, but the latter more generally, on account of the superior beauty of its flowers, which are of a fine deep red colour, and have the appearance of being double, though the petals are not preternaturally multiplied; when the plant is in bloom the peduncles hang down, when the petals drop they become erect; the blossoms are not distinguished by their fragrance, though the seed-vessels, and seeds (which do not come to perfection with us) are said to be strongly odoriferous; the foliage of this plant is also much admired: taking it indeed altogether, there are few shrubs held in higher estimation.

According to Mr. AITON, this species was introduced by JOHN ELLIS, Esq. in 1776; but ISAAC WALKER, Esq. of Southgate, was the first who possessed it in this country, he informs me by letter, that he received plants of it from Pensacola in 1771, by the hands of Mr. JOHN BRADLEY, and that he communicated some of them to Dr. FOTHERGILL, Dr. PITCAIRN, and Mr. ELLIS. It flowers from April to July.

Cultivators differ widely as to their treatment of this plant, some keeping it in the stove, others in the green-house, while some have ventured to plant it in the open ground in warm situations; it probably is more hardy than we imagine; all agree in propagating it by layers, or by seeds if they can be procured. LINNÆUS, contrary to his usual practice, distinguishes the two species by their colour only, and THUNBERG is disposed to regard them as mere varieties.

[440]

ERICA ALBENS. PALLID HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* apice bifidæ, pertusæ. *Caps.* 4-locularis, 4-valvis, polysperma.

Specific Character and Synonyms.

ERICA *albens* antheris muticis inclusis, corollis ovatis oblongis acutis, foliis ternis, racemis secundis. *Linn. Syst. Vegetab. Murr. ed. 14. p. 367. Mant. 233.*

ERICA *albens* mutica, foliis ternis trigonis pilosis, floribus lateralibus, calyce villosa. *Thunb. Prodr. p. 70.*



N^o440.

This species, a native of the Cape, has been introduced since the publication of the *Hort. Kew.* and is now to be found in most green-house collections near town.

In its habit, its foliage, and its flowers, it is very distinct from all our other heaths; flowers from April to June, is readily increased by cuttings, and easily kept with the common treatment.

[441]

ANTHOLYZA MERIANELLA. DWARF ANTHOLYZA.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Cor. tubulosa, irregularis, recurvata. *Caps.* infera.

Specific Character and Synonyms.

ANTHOLYZA *Merianella* corollis infundibuliformibus, foliis linearibus. *Linn. Syst. Vegetab. ed. 14. Murr. p. 87. Ait. Kew. v. 1. p. 67.*

WATSONIA *humilis* foliis lineari-ensiformibus, tubo floris longissimo. *Mill. ic. 198. t. 297. f. 2.*



Nº441.

This very rare species is perfectly distinct from the *Meriana*, of more humble growth, the flowering stem seldom rising to more than a foot in height, and producing from four to six flowers, which are proportionably longer, more closed, and of a deeper red colour than those of *Meriana*.

Was introduced from the Cape by Capt. HUTCHINSON, in 1754. *Ait. Kew.*

Is readily increased by offsets, and requires the same treatment as the *Anthol. Meriana* already figured.

Flowers in May and June.

Our drawing was made from a plant which flowered with Mr. FAIRBAIRN, at the Apothecaries Garden, Chelsea, May 2, 1798.

[442]

GENISTA LINIFOLIA. FLAX-LEAVED BROOM.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus 2/3. *Vexillum* oblongum, a pistillo staminibusque deorsum reflexum.

Specific Character and Synonyms.

GENISTA *linifolia* foliis ternatis sessilibus linearibus subtus sericeis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 645. Ait. Kew. v. 3. p. 14.*

CYTISUS *argenteus* linifolius insularum stæchadum. *Tourn. Inst. 648.*



Nº442.

The *Genista linifolia* is a native of Spain, and was introduced to the royal garden at Kew, by Sir FRANCIS DRAKE, in the year 1786. *Ait. Kew.*

The bright yellow flowers, which are abundantly produced on this plant during May and June, joined to the silky appearance of its foliage, has rendered it worthy the notice of most lovers of plants, in whose collections it is now generally found.

It is most successfully propagated by seeds, which usually ripen in this country; it may also be raised from cuttings, but not readily, agreeing in this respect with leguminous plants in general.

We do not find in LINNÆUS'S works any figure of this plant referred to; but there is a minute description of it in his *Sp. Pl.*

It is usually kept in the greenhouse; being a native of Spain, it may probably be more hardy than we imagine.

[443]

ERICA PHYSODES. STICKY-FLOWERED HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* apice bifidæ pertusæ. *Caps.* 4-locularis, 4-valvis, polysperma.

Specific Character and Synonyms.

ERICA *physodes* antheris cristatis, corollis ovatis inflatis, stylo incluso, foliis quaternis, floribus subsolitariis. *Linn. Syst. Vegetab. Murr. p.* 366. *Berg. Cup.* 108.

ERICA *physodes* cristata, foliis quaternis linearibus, floribus umbellatis viscosis, calyce ovato brevi. *Thunb. Prodr. p.* 74.



N^o443.

The *Erica physodes* is a native of the Cape, and another of those species which have been introduced since the publication of the *Hort. Kew.* of Mr. AITON.

It is principally distinguished by the form, and delicate whiteness of its blossoms, which are so extremely viscid as to retain flies and other insects which settle on them.

Is scarcely to be increased by cuttings, more readily by seeds, which sometimes ripen here.

[444]

CANARINA CAMPANULA. CANARY BELL-FLOWER.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cal. 6-phyllus. *Cor.* 6-fida, campanulata. *Stigmata* 6. *Caps.* infera, 6-locularis, polysperma.

Specific Character and Synonyms.

CANARINA *Campanula* caule erecto, foliis hastatis ternis oppositisve. *Martyn Mill. Dict. Linn. Syst. Veg. ed. 14. Murr. p. 344. Ait. Kew. v. 1. p. 480. Linn. Mant. p. 225.*

CAMPANULA *canariensis* capsulis quinquelocularibus, foliis hastatis dentatis oppositis petiolatis. *Linn. Sp. Pl. ed. 3. p. 238.*

CAMPANULA *canariensis* regia s. medium radice tuberosa, foliis sinuatis cæsiis atriplicis æmulis ternis circum caulem ambientibus, flore amplo pendulo colore flammeo rutilante. *Pluk. Alm. 76. t. 276. f. 1.*



N^o444.

The flowers of this plant so strongly resemble those of the *Campanula*, that it is no wonder the older Botanists regarded it as such, LINNÆUS himself did so at first, and MILLER also; and even now it may perhaps be doubted whether it ought to be made a distinct genus of, since it is found to differ principally in the number of its parts of fructification.

It is a native of the Canary Islands, whence its name, was cultivated in the royal garden, Hampton-Court, as long since as the year 1696^[1], and is a tender herbaceous plant, to be found in most of our greenhouses; its stem rises to the height of six or more feet, its flowers produced singly from the fork of the stalk are large and shewy, they begin to open at the commencement of winter, and continue to blow till March.

"Is propagated by parting of its roots, which must be done with caution; for, as the root is fleshy, if they are broken or wounded, the milky juice will flow out plentifully; so that if these are planted before the wounds are skinned over, it occasions their rotting: the best time for transplanting and parting of their roots is in July, soon after the stalks are decayed; the soil

should be a light sandy loam, mixed with a fourth part of screened lime rubbish." MILLER.

[\[1\]](#) Ait. Kew.

[445]

CORONILLA EMERUS. SCORPION SENNA.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus: 2/3: dentibus superioribus connatis. *Vexillum* vix alis longius. *Legumen* isthmis interceptum.

Specific Character and Synonyms.

CORONILLA *Emerus* fruticosa, pedunculis subtrifloris, corollarum unguibus calyce triplo longioribus, caule angulato. *Linn. Syst. Vegetab. ed. 14. Murr. p. 669. Ait. Kew. v. 3. p. 57.*

COLUTEA siliquosa s. scorpioides major. *Bauh. Pin. p. 1046.*



N^o445.

Authors have given to this plant the name of Scorpion Senna, its seed-vessels, from their slender and jointed appearance, bearing some resemblance to the tail of a scorpion.

It is a native of France and Germany, and a very old inhabitant of our gardens, having been cultivated by GERARD in 1596; it is of low and slow growth: there is a shrub of it in the Apothecaries Garden, Chelsea, which grew there in the time of MILLER, and which now is not more than five feet high.

In the nurseries we have observed two varieties of it, one in which the flowers have been tinged with bright red inclining to orange and which is by far the most common, the other with flowers wholly yellow, scarcely worth cultivating.

Its blossoms are produced in May and June, and sometimes again in autumn; neatly trained to a wall or paling, it makes a beautiful appearance when in flower, the shortness of its shoots renders it a very proper object for this purpose.

Is propagated by seeds, layers, and cuttings; the first are not produced in any great plenty with us.

The leaves by a proper fermentation are said to produce a dye like that of Indigo.

[446]

PSORALEA BRACTEATA. OVAL-SPIKED PSORALEA.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. punctis callosis adpersus longitudine leguminis
monspermi.

Specific Character and Synonyms.

PSORALEA *bracteata* foliis ternatis obovatis recurvato-
mucronatis, mucronatis, spicis ovatis. *Linn. Mant.* 264.
Berg. Cap. 224. *Ait. Kew.* v. 3. p. 79.

TRIFOLIUM *fruticans*. *Linn. Sp. Pl. ed. 3. p.* 1085.

TRIFOLIUM *africanum fruticans flore purpurascente.* *Comm.*
Hort. 2. p. 211. t. 106.



Nº446.

The old authors, and indeed LINNÆUS himself at first, regarded this plant as a *Trifolium*; afterwards the latter changed it to *Psoralea*, and minutely described it in his *Mantissa*.

As a green-house plant, this small and delicate species has long been cultivated^[1], and still continues to hold a place in all collections of note.

Its inflorescence to us has more the appearance of a *capitulum* than a *spike*, and which when the plant is in full bloom, is by no means ovate, but rather hemispherical; the purple colour of the vexillum, contrasted with the white of the alæ, gives to the flowers a very pleasing parti-coloured appearance.

In its leaves we have a good example of the *folium mucronatum*.

It is a native of the Cape, flowers in June and July, and is usually propagated by cuttings.

^[1] By MILLER, in 1731.

[447]

ERICA EMPETRIFOLIA. CROWBERRY-LEAVED HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* apice bifidæ pertusæ. *Caps.* 4-locularis, 4-valvis, polysperma.

Specific Character and Synonyms.

ERICA *empetrifolia* foliis oblongis ciliatis, floribus verticillatis: calyce ciliato. *Thunb. Prodr.* p. 73. *Linn. Syst. Vegetab. ed. 13. Gmel. p. 627. Linn. Pflanzen Syst. 3. t. 23. f. 2.*

ERICA *empetrifolia* antheris cristatis, corollis ovatis, foliis quaternis, floribus sessilibus lateralibus. *Linn. Syst. Vegetab. ed. 14. Murr. p. 366. Ait. Kew. v. 1. p. 19.*



N^o447.

It is a most pleasing circumstance, when plants afford characters by which they may with certainty be distinguished; most of the Heaths are of this kind, and the present one in particular: exclusive of its great peculiarity of growth, so obviously expressed in our representation of it, its flowers diffuse a strong honey-like fragrance, which, if other characters were wanting, would at once discriminate it.

It is one of those Heaths which are enumerated in the *Hort. Kew.* of Mr. AITON, and was introduced to the royal garden by Mr. MASSON, in 1774; is now to be met with in most of the collections of green-house plants about town, and flowers in May and June.

Is usually propagated by cuttings.

[448]

**MESEMBRYANTHEMUM MICANS. GLITTERING FIG-
MARYGOLD.**

Class and Order.

ICOSANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-fidus. *Petala* numerosa, linearia. *Caps.* carnosae, inferae, polyspermae.

Specific Character and Synonyms.

MESEMBRYANTHEMUM *micans* foliis subcylindricis papulosis distinctis, caule scabro. *Linn. Syst. Veg. ed. 14. Murr. p. 470. Ait. Kew. v. 2. p. 190.*

MESEMBRYANTHEMUM *micans* flore phoeniceo: filamentis atris. *Dill. Elth. 292. t. 215. f. 1282.*

FICOIDES *capensis*, tereti folio, flore croceo. *Pet. gaz. t. 7. f. 9.*

FICOIDES *capensis*, folio tereti argenteo, petalis perplurimis aurantiacis. *Bradl. Succ. 1. p. 9. t. 8.*



N°448.

The *Mesembryanthemum micans*, so called from the glittering particles which are conspicuous on its stalks and leaves, is a species which has long been introduced to our gardens (having been cultivated by Prof. BRADLEY in 1716) for the beauty of its flowers, which in richness of colour are indeed surpassed by few; they are produced in the months of July and August, but do not expand fully, unless the sun shines powerfully on them; nor do they long retain that regular expansion observable in some species, but quickly assume a somewhat ragged appearance; nevertheless, upon the whole, it is one of those species which is highly deserving of culture, by those who are partial to this tribe of plants.

It is a native of the Cape, and readily propagated by cuttings.

Varies with flowers of a paler hue.

[449]

DILLENIA SPECIOSA. SHEWY DILLENIA.

Class and Order.

POLYANDRIA POLYGYNIA.

Generic Character.

Cal. 5-phyllus. *Petala* 5. *Caps.* polyspermæ, connatæ, pulpa repletæ.

Specific Character and Synonyms.

DILLENIA *speciosa* foliis oblongis rotundato-acutis denticulatis, pedunculis unifloris. *Thunb. in Linn. Trans.* 1. p. 200.

DILLENIA *indica*. *Linn. Syst Vegetab. ed. 14. Murr. p.* 507.

SYALITA Malabaris. *Rheede Hort. Malab. tom. iii. p.* 39. t. 38. 39.



Nº449.

The name of *Dillenia* was given by LINNÆUS to this genus of plants, in honour of JOHN JAMES DILLENIIUS, Professor of Botany at Oxford, and the celebrated author of the *Hist. Muscorum*, *Hort. Elthamensis*, &c.

Until the publication of the first volume of the *Linnean Transactions*, only one species of *Dillenia* was generally known; in that work Prof. THUNBERG minutely describes five others, three of which are there figured; all these, and one more described by Dr. ROXBURGH in his work on the Coromandel plants, are inserted by Prof. MARTYN in his new and highly improved edition of MILLER'S *Dictionary*.

The present species, which now loses its name of *indica* in that of *speciosa*, and which though not enumerated in the *Hort. Kew.* has many years been cultivated at Kew, and in the stoves of the curious near town, is a native of Malabar and Java; in its native soil it becomes a vast tree, here we rarely see it more than two or three feet high; its flowers are large and shewy, but quickly deciduous, and remarkable for the unpleasantness of their scent,

which is like that of the *Lycium japonicum*, but not being readily exhaled does not infect the air of the stove.

This species is increased without difficulty by cuttings, which quickly produce flowering plants.

[450]

GLADIOLUS WATSONIUS. WATSON'S CORN-FLAG.

Class and Order.

TRIANDRIA MONOGYNIA.


Generic Character.

Cor. 6-partita, tubulosa, ringens. *Stamina* adscendentia.

Specific Character and Synonyms.

GLADIOLUS *Watsonius* corollis infundibuliformibus, limbi laciniis subæqualibus, foliis linearibus ad oras revolutis. *Jacq. Icon. rar.* 2. t. 233. *Collect.* 3. p. 257. *Linn. Sp. Pl.* ed. *Wilden.* p. 214.

GLADIOLUS *Watsonius* monostachyus corollis cernuis, tubo duplici, limbi laciniis oblongis, foliis linearibus glabris. *Thunb. Prod.* p. 8.



Nº450.

Professors JACQUIN and THUNBERG have both described this rare bulbous plant, a native of the Cape, and newly introduced to this country from Holland among a great variety of others: it seems highly probable that the descriptions of both these authors were taken from dried specimens, since they accord so little with the living plant as it flowers with us.

At first sight, one would be led to regard this plant as an *Antholyza* rather than a *Gladiolus*, its flowers bearing a great affinity to those of the *Antholyza Meriana*, which differs widely from those of the *Gladiolus communis*: Prof. THUNBERG having thought fit to make a *Gladiolus* of that plant, he could do no less than regard this as a *Gladiolus* also; we regret that the infinite variety to which all the productions of nature are subject should give occasion to versatile minds perpetually to alter genera, often without due consideration. This species flowers in February and March, requires the same treatment as other Cape bulbs, and is propagated in the same manner.

DESCR. Stalk from twelve to eighteen inches high, upright, smooth; Leaves about three or four, the lowermost a sheath merely, the second leaf springs

from a long sheath, is puckered at its base, three inches in length, upright, rigid, flat, linear-lanceolate, having three strong ribs, one in the middle, two at the margin, which, projecting on each side, give to the edge of the leaf a thick appearance, the leaves as they ascend, gradually differ from this, and finally become hollow bracteæ, which at first envelope the flowers, and afterwards contribute to support them; Flowers from two to three, each standing on a peduncle-like tube, enclosed by a bifid spatha, contained within and about half the length of the bracteæ; Corolla bright red, funnel-shaped, tube bent somewhat downwards, nearly cylindrical, a little flattened, and glossy, limb divided into six ovato-lanceolate segments, spreading outwards, the uppermost segment incumbent, the three lowermost smaller than the others; Filaments three, whitish, nearly straight; Antheræ oblong, straight, purple; Style red; Stigma trifid, each segment dividing into two villous lips.

[451]

BLAKEA TRINERVIA. THREE-RIBBED BLAKEA.

Class and Order.

DODECANDRIA MONOGYNIA.

Generic Character.

Cal. inferus, 6-phyllus, superus, integer. *Petala* 6. *Capsula* 6-ocularis, polysperma.

Specific Character and Synonyms.

BLAKEA *trinervia* bicalyculata, foliis enerviis transversim subtilissime striatis. *Linn. Suppl. p. 246. Syst. Veg. ed. 14. Murr. p. 442.*

BLAKEA foliis ellipticis trinerviis nitidis, floribus lateralibus. *Brown. Jam. 323. t. 35. The wild Rose.*



Nº451.

"This vegetable is certainly one of the most beautiful productions of *America*. It is but a weakly plant at first, and supports itself for a time by the help of some neighbouring shrub or tree; but it grows gradually more robust, and at length acquires a pretty moderate stem, which divides into a thousand weakly declining branches, well supplied with beautiful rosy blossoms on all sides that give it a most pleasing appearance in the season.

"It is chiefly found in cool, moist, and shady places, and grows generally to the height of ten or fourteen feet; but rises always higher when it remains a climber, in which state it continues sometimes. It thrives best on the sides of ponds or rivulets, and those that would choose to have it flourish in their gardens, where it must naturally make a very elegant appearance, ought to supply it with some support while it continues young and weakly.

"It is called *Blakea*, after Mr. MARTIN BLAKE, of Antigua, a great promoter of every sort of useful knowledge, and a gentleman to whose friendship the Natural History of Jamaica chiefly owes its early appearance." *Brown's Jamaica*.

Our figure was drawn from a very fine healthy plant which flowered in the collection of Lady DOWNE, at her villa of Bookham-Grove, near Leatherhead, in April 1799; though not enumerated in the *Hortus Kewensis*, it had produced blossoms in several other collections near town long before this period; those when they once expand are of short duration, but the foliage when healthy is always handsome.

It is usually kept in the stove with other Jamaica plants, and propagated by layers.

We cannot see the propriety of applying *foliis enerviis* to the description of this species, since LINNÆUS himself, in his *Sp. Pl.* describes the leaves as *trinervia*; three strong ribs they always have, and usually two others near the margin which are finer.

[452]

**CARDAMINE TRIFOLIA. THREE-LEAVED CUCKOW-
FLOWER.**

Class and Order.

TETRADYNAMIA SILIQUOSA.

Generic Character.

Siliqua elasticæ dissiliens valvulis revolutis. *Stigma* integrum.
Cal. subhians.

Specific Character and Synonyms.

CARDAMINE trifolia foliis ternatis obtusis, caule subnudo.
*Linn. Syst. Vegetab. ed. 14. Murr. p. 593. Ait. Kew. v. 2.
p. 387.*

NASTURTIUM alpinum trifolium. *Bauh. Pin. 104.*

CARDAMINE trifolia. Trefoile Ladies smockes. *Parkins.
Parad. p. 389.*



Nº452.

Such as are attached to the smaller alpine plants, will regard this species of *Cardamine* as worthy a place in their collections; one would scarcely have expected to find it in PARKINSON'S *Parad.* yet there it is described, and the following account given of its introduction: "It was sent me by my especial good friend JOHN TRADESCANTE, who brought it among other dainty plants from beyond the seas, and imparted thereof a root to me."

This species is perennial, hardy, and of very humble growth; the leaves grow thickly together, forming a kind of tuft; the flowering stems rarely rise above the height of six inches, and produce on their summits numerous flowers, waved on their edges; all those which we have had an opportunity of seeing have been perfectly white, PARKINSON and HALLER describe them as being sometimes tinged with red or purple; they begin to appear towards the end of March and continue through April, the shelter of a hand-glass open at top is often necessary to protect and improve the flowering of this and other such early-blowing plants.

It grows readily either in a pot or in the open border, succeeds best when planted in bog earth in a situation moderately moist and shady, and is readily increased by parting its roots, which are somewhat of the creeping kind.

Grows spontaneously in most of the northern parts of Europe, especially Lapland, Switzerland, and Austria.

[453]

AMARYLLIS REGINÆ. MEXICAN LILY.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. hexapetaloidea irregularis. *Filamenta* fauci tubi inserta, declinata, inæqualia proportione vel directione. *Linn. fil.*

Specific Character and Synonyms.

AMARYLLIS *reginæ* spatha subbiflora, pedicellis divaricatis, corollis campanulatis breve tubulosis nutantibus, fauce tubi hirsuta, foliis lanceolatis patulis. *Linn. fil. Ait. Kew. 416. Mill. ic. p. 16. t. 24.*

LILIUM americanum puniceo flore Belladonna dictum. *Herm. par. 194. t. 194.*



Nº453.

DESCR. "Bulb green, scape round, somewhat flattened: Corolla scarlet, with a bottom of a whitish green, the three outer petals round at the tip, the three inner fringed at the base; the style red, the flower stems seldom rise more than one foot high; each stem supports two, three, or four flowers, rarely more; they are large, and of a bright copper colour, inclining to red; the spathe which cover the buds before they open, divides into two parts to the bottom, standing on each side the umbel of flowers joined to the peduncles.

"It flowered in Mr. FAIRCHILD's garden at Hoxton, in 1728, when the late Dr. JAMES DOUGLASS caused a figure of it to be drawn, and wrote a folio pamphlet on it. He gave it the title of *Lilium Reginae*, because it was in full beauty on the first of March, which was the late queen's birth-day. Mr. FAIRCHILD told me the roots were brought from Mexico; so he gave it the name of Mexican Lily, which is still continued to it by the English gardeners. It flowers constantly in the spring, when it is placed in a very warm stove. It is in beauty in February, and those which are in a moderate temperature of air will flower in March or April.

"Not being so hardy as some others, it must be placed in a warm stove, and if the pots are plunged into a hot-bed of tanner's bark, the roots will thrive better, and the flowers will be strong.

"It is increased by offsets." *Mill. Dict. ed. Mart.*

[454]

CANNA INDICA. COMMON INDIAN REED or SHOT.

Class and Order.

MONANDRIA MONOGYNIA.

Generic Character.

Corolla 6-partita erecta: labio bipartito, revoluta. *Stylus* lanceolatus, corollæ adnatus. *Calyx* 3-phyllus.

Specific Character and Synonyms.

CANNA *Indica* foliis ovatis utrinque acuminatis nervosis. *Sp. Pl. 1. Ait. Kew. vol. 1. p. 1.*

CANNACORUS. *Rumph. amb. 5. p. 177. t. 71. f. 2.*

ARUNDO indica latifolia. *Bauh. Pin. 19.*

CANNA *Indica* flore rubro. Red flowred Indian Reede. *Park. Parad. p. 376.*



Nº454.

The *Canna indica*, a native of both the Indies, is a plant greatly admired for the beauty of its foliage and flowers, and on that account generally cultivated; it has been called by some *Indian Shot*, from the roundness and hardness of its seeds.

We find it to have existed in our gardens in the time of GERARD, 1596. PARKINSON was acquainted with that variety of it which has yellow spotted flowers: Prof. MARTYN, in his edition of *Miller's Dict.* has quoted the chief of what these authors say of it, which as a matter of curiosity we shall here transcribe: "GERARD informs us, that in his time it was in the garden at Padua, that he had planted it in his garden divers times, but it never came to flowering; and that it must be set or sown in a pot, with fine earth, or in a bed made of horse-dung, in such manner as Cucumbers and Musk-Melons are: PARKINSON says, in some kindly years this beautiful plant has borne its brave flowers, but never any ripe seed, and that it will not abide the extremities of our winters, unless it meet with a stove, or hot-house, such as are used in Germany; for neither house nor cellar will preserve it: CLUSIUS

saw it flowering by house-sides in Spain and Portugal, and says, that the inhabitants there use the seeds for making their rosaries."

Mr. AITON enumerates four varieties of it, viz. *rubra*, *lutea*, *coccinea*, and *patens*.

"Being a native of the warmest parts of America, it requires to be placed in a moderate stove in winter, where they always flower in that season, at which time they make a fine appearance, and in the summer place them abroad in a sheltered situation with other tender exotic plants, where they generally flower again, and produce ripe seeds annually." *Mill. Dict.*

"These plants will continue many years with proper management, but as young plants always flower better than the old root, so it is scarce worth while to continue them after they have borne good seeds, which should be sown on a hot-bed in the spring." *Mill. Dict.*

[455]

ALOE RETUSA. CUSHION ALOE.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. erecta, ore patulo fundo nectarifero. *Filam.* receptaculo inserta.

Specific Character and Synonyms.

ALOE retusa acaulis foliis quinquefariis deltoideis. *Thunb.*
Aloe, n. 15. *Ait. Kew.* 471.

ALOE retusa floribus sessilibus triquetris bilabiatis labio inferiore revoluta. *Linn. Sp. Pl.* 459.

ALOE africana brevissimo crassissimoque folio, flore viridi.
Comm. hort. 2. p. 11. t. 6. *Till. pis.* 6. t. 5.



N^o455.

Though the flowers of this Aloe have little to recommend them, there is much to admire in the form and structure of its leaves; and this pleasing circumstance attends it, it is perfectly distinct from all the other species: when first introduced, it was no doubt an object of great admiration; FAIRCHILD, the celebrated Gardener of Hoxton, who preceded MILLER, had it engraved, with several other succulents, on a plate which is prefixed to Dr. BLAIR'S *Botanic Essays*, and which he described to the Doctor, betwixt whom and Mr. FAIRCHILD there appears to have subsisted a great degree of intimacy: the Essays were printed in 1720.

This species is a native of the Cape, and flowers in June, but not regularly so, increases very fast by offsets: Mr. AITON makes it as he does all the Aloes indiscriminately, Dry Stove Plants, but it may be kept in a good green-house, taking care to place it in the driest and most airy part, and to guard it at all times from much wet, but more especially in the winter season.

[456]

DIOSMA SERRATIFOLIA. SERRATED or SAW-LEAVED

DIOSMA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. 5-petala. *Nectaria* 5 supra germen. *Caps.* 3. s. 5. coalitæ.
Sem. calyptrata.

Specific Character.

DIOSMA *serratifolia* foliis lanceolatis glanduloso-serrulatis,
pedunculis axillaribus oppositis subunifloris.



Nº456.

It was in the green-house of Mr. WHITLEY, Nurseryman at Old Brampton, that we first saw this plant in flower, on the 25th of March 1799; previous to this period it had flowered in the Royal Garden at Kew, and specimens of it had been deposited in the Herbarium of Sir JOSEPH BANKS, under the name of *Diosma serratifolia*.

It forms a neat pretty shrub, which is rendered more desirable by its early flowering; its blossoms are pure white, set off by Antheræ of a lively purple colour; the whole plant has a strong scent very like that of Pennyroyal, but more powerful.

This shrub is one of the many which have been raised within these few years from Botany-Bay seeds, is a greenhouse plant, of easy culture, blows freely, and is readily increased by cuttings.

DESCR. Twigs somewhat angular, reddish purple; Leaves opposite, narrow, on very short footstalks, spreading, slightly truncated at the extremity, finely toothed, a transparent gland in the angle of each tooth, beset on both sides with numerous glands, which project and give to the upper surface a

manifest roughness; Flowers from the alæ of the leaves, opposite, on peduncles about one-third of an inch long, usually supporting one, sometimes two flowers; Calyx composed of five leaves which are tinged with red and permanent; Corolla five petals, white, ovato-lanceolate; Stamina filaments five, white, hairy, at first upright, afterwards bending back betwixt the petals; Antheræ before they open purple; Pollen yellow; Nectaries of two kinds, five white, springing up between each filament, and which may perhaps be considered as so many imperfect stamina, they are broader, much smoother, and about one-third of the length of the filaments, terminating in a transparent gland instead of an anthera; five green, forming a kind of calyx to the germen, from the top of which they spring, they consist of five roundish, fleshy, spreading, green leaves, edged with hairs; besides these, there is a glandular ring at the base of the germen; Style white, tapering, very hairy; Stigma simple; Flowers usually dropping off without ripening the seeds.

[457]

ALOE PLICATILIS. FAN ALOE.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. erecta, ore patulo, fundo nectarifero. *Filam.* receptaculo inserta.

Specific Character and Synonyms.

ALOE *plicatilis* subacaulis, foliis linguæformibus lævibus distichis, floribus racemosis pendulis cylindricis. *Ait. Kew. v. 1. p. 470.*

ALOE *disticha* var. ϵ . *Linn. Sp. Pl. ed. 3. p. 459.*

ALOE *plicatilis* foliis ensiformibus inermis ancipitibus, floribus laxè spicatis, caule fruticoso. *Mill. Dict. ed. 6. 4to.*

ALOE *africana* arborescens montana non spinosa, folio longissimo plicatili, flore rubro. *Comm. Hort. 2. p. 5. t. 3.*



Nº457.

"The Fan Aloe grows to the height of six or seven feet, with a strong stem, towards the upper part of which are produced two, three, or four heads, composed of long, compressed, pliable leaves, of a sea-green colour, and ending obtusely; these are placed in a double row, lying over each other, with their edges the same way; the flowers are produced in short loose spikes, are of a red colour, and appear at different times of the year." *Mill. Dict.*

LINNÆUS originally made this plant a variety of his *Aloe disticha*, the leaves in their mode of growth are indeed truly distichous, few plants afford a better example of such, but they differ materially from those of the real *disticha* both in form and colour: Mr. MILLER, with great propriety, made a distinct species of it, by the name of *plicatilis*, or *Fan Aloe*, which Mr. AITON has continued; and by the name of *Fan Aloe* it is very generally known: we may remark, however, that though this term may be justified by the form into which the leaves expand, the *folium plicatile* of LINNÆUS is a very different kind of a leaf.

Both the foliage and flowers of this plant are very handsome, in the course of many years it grows to a great size; in the Chelsea Garden there are some fine plants of it, which grew there in the time of MILLER, by whom it was cultivated in 1731.

It is native of Africa, requires the same treatment as the other Aloes, and is propagated by dividing and planting its heads.

[458]

ARISTEA CYANEA. GRASS-LEAVED ARISTEA.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Petala 6. *Stylus* declinatus. *Stigma* infundibuliforme hians.
Caps. infera polysperma.

Specific Character and Synonyms.

ARISTEA *cyanea*. Ait. *Kew.* v. 1. p. 67.

IXIA *africana* floribus capitatis, spathis laceris. Linn. *Sp. Pl.*
ed. 3. p. 51.

MORÆA *africana* floribus capitatis spathis laceris. Murr. *Syst.*
Vegetab. ed. 14. p. 93.

IXIA foliis ad radicem nervosis gramineis, floribus ac fructu
convolutis. *Burm. Afric.* 191. t. 70. f. 2.

BERMUDIANA capensis, capitulis lanuginosis. *Pet. sicc.* 242.

GRAMEN eriophorum africanum flore lanato. *Pluk. Mant.* 98.



Nº458.

It will be seen, on consulting the synonyms, that this native of the Cape, though introduced to the Kew Garden by Mr. MASSON in 1774, was long before known to a considerable number of Botanists, and it is curious to see the different opinions which they entertained of it; we abide by that of Mr. AITON, who has called it *Aristea*, from the bearded appearance, we apprehend, of the Spathæ.

It is a small fibrous-rooted plant, rarely exceeding when in bloom the height of six or eight inches, and would be too insignificant for a green-house collection, were not its flowers of a very brilliant blue colour; indeed MILLER, who appears evidently to have cultivated it, says, the flowers make little appearance, and so the plant is only kept for the sake of variety. *Dict. 4to. ed. 6. Ixia africana.*

Mr. AITON tells us, that it flowers from April to June, yet Mr. ANDREWS, intent on giving to Messrs. LEE and KENNEDY the credit of flowering it first, disregards this information, and is pleased to conjecture that the plant never flowered at Kew, because Mr. AITON, as he alleges, has not given to it any

specific character; not aware that, as a new genus, its parts of fructification are described at the end of the *Hort. Kew.* and that no specific character is ever given to a plant, where there is only one of a genus, and that for the most obvious reason.

The *Aristea* is a plant easily propagated by parting its roots, as well as by seeds, will succeed in a small pot, and though a green-house plant, will not be hurt by the moderate heat of the stove, but flower the better for it.

The blossoms do not expand fully unless the sun shines hot on them.

[459]

**CONVOLVULUS CNEORUM. SILVERY-LEAVED BIND-
WEED.**

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. campanulata, plicata. *Stigmata* 2. *Caps.* 2-locularis
loculis dispermis.

Specific Character and Synonyms.

CONVOLVULUS *Cneorum* foliis lanceolatis tomentosis,
floribus umbellatis, calycibus hirsutis, caule erecto. *Linn.*
Syst. Veg. 203. *Ait. Kew. vol. 1. p.* 213.

CONVOLVULUS *argenteus umbellatus erectus.* *Tournef. Inst.*
84.

CONVOLVULUS *major erectus creticus argenteus.* *Moris.*
Hist. 2. p. 11 s. 1. t. 3. f. 1.

CONVOLVULUS *saxatilis erectus villosus perennis.* *Barr.*
rar. 4. t. 470. Bocc. Mus. 2. p. 79. t. 70.

CNEORUM *album folio argenteo molli.* *Bauh. Pin.* 463.



Nº459.

The *Convolvulus Cneorum* is a native of Spain and the Levant, was cultivated in the Botanic Garden at Chelsea in 1739, and flowers from May to September. *Ait. Kew.*

In size, habit, &c. this species has some affinity to the *Convolvulis linearis*, figured pl. 289, but differs from it, and other species usually cultivated with us, in the silky appearance of its foliage, which it is not in the artist's power to imitate, and for the beauty of which, more than that of its flowers, it is very generally kept in collections of green-house plants; its blossoms are nearly white and rarely or never productive of seeds in this country, hence it is increased by cuttings.

It is a hardy green-house plant, requiring a dry rather than a moist regimen.

[460]

**MAURANDYA SEMPERFLORENS. CLIMBING MAURANDYA,
or BASTARD FOXGLOVE.**

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Caps. bilocularis, truncata, bisulca, apice inæqualiter
dehiscens. *Cor.* ringens, tubulosa campanulata,
ventricosa, bisulcato-plicata.

Specific Character and Synonyms.

MAURANDYA *semperflorens* caule fruticoso scandente,
foliis hastatis nervosis. C. G. Orteg. Nov. Pol. Dec. 2. p.
21.

USTERIA. Cavanill. Icon. vol. 2. p. 5. num. 126. t. 116.



Nº460.

The plant whose elegant form is here so happily delineated by the masterly pencil of Mr. EDWARDS, according to Dr. ORTEGA, is an inhabitant of Mexico, where its seeds originally were collected by Dr. MARTIN SESSE, and sent to the royal garden at Madrid, in which the plant produced flowers and seeds in abundance; from thence seeds were obtained by the Marchioness of BUTE, about the year 1786, who most kindly communicated them to different persons in the neighbourhood of London, and among others to my most generous benefactor, JAMES VERE, Esq. in whose collection at his villa, Kensington-Gore, by the careful management of his Gardener, WILLIAM ANDERSON, it was first brought to flower in this country, anno 1797.

Mons. CAVANILLE, who resides at Madrid, where he seizes every opportunity of publishing whatever new plants appear there, has figured and described the present one, under the name of *Usteria*, not aware that WILDENOU had previously bestowed that name on a different plant; for this and other reasons^[1], Dr. ORTEGA, in a new work of his above referred to, has changed the term *Usteria* to that of *Maurandya*; and, though we cannot cordially

coincide with the Doctor in the propriety either of his generic or trivial name, we have adopted them.

This climber rises with a shrubby stalk to the height of many feet, is very prolific in branches, and produces flowers abundantly from July to September, which are succeeded by ripe seed-vessels and seeds.—As the plant is easily propagated by cuttings, as well as by seeds, it will soon become common to our greenhouses, though it is rather better suited to the conservatory; if its blossoms, which have a great affinity to those of the Foxglove, had more colour in them, the plant would be more desirable: at some future period such may probably be obtained from seeds.

[1] OBSERV. Quandoquidem USTERIÆ nomen huicGeneri à clar. CAVANILLES, qui primus id descripsit, impositum permanere non potest ut pote antea ab illust. WILDENOU aliiGeneri inditum; propterea illud D. CATHARINÆ PANCRATIÆ MAURANDY, lectissimæ feminæ, D. AUGUSTINI JUAN, Reg. Botanic. Horti Carthaginensis Professoris uxori, et Botanicorum laborum sociæ, nuncupandum duxi; oblataque opportunitate usus plantam denuo recognovi, et tum Characterem genericum, tum etiam descriptionem specificam ad incudem revocans reformavi.

[461]

JASMINUM FRUTICANS. YELLOW JASMINE.

Class and Order.

DIANDRIA MONOGYNIA.

Generic Character.

Cor. 5-fida. *Bacca* dicocca. *Sem.* arillata. *Antheræ* intra tubum.

Specific Character and Synonyms.

JASMINUM *fruticans* foliis alternis ternatis simplicibusque ramis angulatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 56.*

JASMINUM *fruticans* foliis alternis ternatis foliolis obovatis cuneiformibusque obtusis ramis angulatis laciniis calycinis subulatis. *Ait. Kew. v. 1. p. 9.*

JASMINUM *luteum*, vulgo dictum bacciferum. *Bauh. Pin. 298.*

POLEMONIUM sieu Trifolium fruticans. Shrubby Trefoil. *Ger. Herb. p. 1129. f.*



N°461.

The Yellow Jasmine is often planted against walls, pales, &c. as the branches are weak and slender and it will grow to be ten or twelve feet high if thus supported; it may however be planted in shrubbery quarters, to which it is better suited than the White Jasmine. The young shoots are of a fine strong green colour, angular, and a little hairy. The leaves are trifoliate, though sometimes they grow singly. They are placed alternately on the branches, are of a thick consistence, smooth, and of a fine deep green colour. These leaves in well-sheltered places remain until the spring before they fall off, so that this plant may not improperly be planted among the Evergreens, especially as the young shoots are always of a strong green. The flowers are yellow, and do not possess the fragrance of the common Jasmine. They are produced chiefly in June, and the blow is soon over. They are succeeded by berries, which when ripe are black, whence its name of Berry-bearing Jasmine. Although this shrub possesses a certain stiffness, which gives it somewhat the appearance of an artificial flower, yet the fine yellow colour of its blossoms contrasts so well with the rich green of the foliage, that in the flower-pot or bouquet it never fails to have a conspicuous and pleasing effect.

Is a native of the South of Europe and the Levant, was cultivated by Mr. JOHN GERARD, in 1597, and flowers from May to October. *Ait. Kew.*

Is easily propagated by suckers or layers; as the flowers have no scent, is not so much cultivated as formerly. *Miller's Dict.*

[462]

ANTHEMIS PYRETHRUM. PELLITORY OF SPAIN.

Class and Order.

SYNGENESIA POLYGAMIA SUPERFLUA.

Generic Character.

Recept. paleaceum. *Pappus* nullus. *Cal.* hemisphæricus, subæqualis. *Flosculi* radii plures quam 5.

Specific Character and Synonyms.

ANTHEMIS *Pyrethrum* caulibus simplicibus unifloris decumbentibus foliis pinnato multifidis. *Linn. Syst. Veget. ed. 14. Murr. p. 776. Woodville's Medical Botany, p. 286.*

CHAMÆMELUM specioso flore, radice longa fervida. *Shaw. Afr. 138.*

PYRETHRUM flore bellidis. *Bauh. Pin. 148.*



Nº462.

The Pellitory of Spain is more celebrated as an useful than an ornamental plant, the root which is of a very hot and biting nature being a common application for the cure of the tooth-ach; but, divested of its utility as a medicinal plant, it merits a place in collections on account of the beauty both of its foliage and flowers, the latter are more handsome when in bud than when fully expanded, the underside of the florets being of a fine purple colour, the upper pure white; it is moreover a very rare plant in this country, notwithstanding it was cultivated here so long since as 1570: PARKINSON evidently grew it, as he observes that the roots of the cultivated plant, were much larger than those of the wild one; he tells us also, that it was too tender to endure our winters: to the latter cause, as well as to the difficulty of propagating it, for it does not ripen its seeds in this country, we may attribute its present scarcity: Mr. MILLER raised this plant in 1732, in a very curious way, from seeds picked out from among raisins.

In its place of growth it is not confined to Spain, but is found in the Levant, Syria, Arabia, and elsewhere; flowers with us from May to July, and may be increased by cuttings of the roots, a mode of propagating by which we

sometimes happily succeed with rare and valuable plants when all others fail.

It is a plant not very nice as to the quality of the soil in which it grows, but must have a warm dry situation, will succeed very well in a pot, or it may be planted in the open border; but especial care must be taken to secure it against frost in the winter.

[463]

EPIDENDRUM CILIARE. FRINGED EPIDENDRUM.

Class and Order.

GYNANDRIA DIANDRIA.

Generic Character.

Cor. 6-petala. *Nectarium* turbinatum, obliquum reflexum.
Caps. infera 1-locularis, 3-valvis, fenestrata.

Specific Character and Synonyms.

EPIDENDRUM *ciliare* foliis oblongis aveniis, nectarii labio tripartito ciliato: intermedia lineari, cauli bifolio. *Linn. Syst. Vegetab. ed. 14. Murr. p. 818. Jacq. Amer. pict. p. 110. t. 209.*



Nº463.

Of this parasitical tribe the number of species described in the third edition of the *Sp. Pl. Linn. 1764*, amounts to thirty; in GMELIN'S thirteenth edition of the *Syst. Nat. Linn. 1791*, no less than seventy-five are enumerated, which is a vast accession in so short a period; most of these are natives of hot climates, and on that account require a stove heat in this country; from their nature also they require a situation not to be exactly imitated, they are therefore to be regarded as plants very difficult of culture, and we are not to be surprised that three species only are enumerated in the *Hort. Kew.*—but, since the publication of that work, many others have been added to the royal collection, and this among the rest.

The rare and singular species here represented, a native of the warmer parts of America, and the West-Indies, flowered with Mr. WHITLEY, Nurseryman, Old-Brompton, in Feb. 1799, and at irregular periods before that time; he informs me that it is not constant as to the time of its blowing, and that though the plant flowers with him, it never assumes a fine healthy green appearance, he propagates it by dividing its limbs or branches, which often

put forth small roots; the plant grows in a pot, in a mixture of loam and peat or bog-earth, and is kept constantly plunged in the tan-pit of the stove.



[464]

SISYRINCHIUM GRAMINEUM. GRASS-LEAVED

SISYRINCHIUM.

Class and Order.

GYNANDRIA TRIANDRIA.

Generic Character.

Spatha 2-phylla. *Petala* 6-plana. *Caps.* 3-locularis infera 3-valvis.

Specific Character and Synonyms.

SISYRINCHIUM *gramineum* caule ancipiti lato, germinibus glabris.

SISYRINCHIUM *angustifolium* foliis lineari-gladiolatis, pedunculis longioribus. *Mill. Dict.*

SISYRINCHIUM *cæruleum* parvum, gladiato caule, virginianum. *Pluk. alm. 348. t. 61. f. 1.*

BERMUDIANA *graminea*, flore minore cæruleo. *Dill. Elth. 49. t. 41. f. 49.*

GLADIOLUS cæruleus hexapetalus caule etiam gladiato. *Banist. virg. 1926.*



Nº464.

In a former Number of this Work, we gave a figure of the large variety of *Sisyrinch. Bermud.* of Linn. regarding it as a distinct species, and naming it *iridioides*, conformably to DILLENIUS's specific description; we regret now that we did not continue to it the name of *Bermudiana* (it being the true Bermudas plant) and which cannot with propriety be applied to the present species, a native of Virginia, far more diminutive, with flowers much smaller, of a paler blue colour, a much hardier plant also, and of more ready growth; it is indeed a truly hardy perennial, adapted to the open border, in which it will grow readily, and produce abundance of flowering stems in June and July; the flowers expand to the sun, and are followed by numerous seed-vessels which ripen their seeds, by which the plant may be increased, or by parting its roots in the autumn. Its stalk affords an excellent example of the *caulis anceps*. We readily embrace the opportunity here afforded us of correcting an error in regard to the hardiness of the *Sisyrinch. iridioides*, which we were led into partly by MILLER's account, and partly from observing the plant to survive a mild winter in the open ground; we have since found that it is a tender plant, and to be ranked with those of the

greenhouse; an additional proof of its being a distinct species from the one here figured, which has long^[1] been cultivated in our gardens.

[1] By Mr. JACOB BOBART, in 1693. *Ait. Kew.*

[465]

INDIGOFERA ANGUSTIFOLIA. NARROW-LEAVED INDIGO.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. patens. *Cor.* carina, utrinque calcari subulato patulo.
Legumen lineare.

Specific Character and Synonyms.

INDIGOFERA *angustifolia* foliis pinnatis linearibus, racemis
elongatis, caule fruticoso. *Linn. Syst. Veget. ed. 14. Murr.*
p. 678. Mant. 272. Ait. Kew. v. 2. p. 69.



Nº465.

This small, delicate, and rather elegant species of Indigo, to be found in most collections of greenhouse plants near town, rises with an upright, shrubby stem, to the height of several feet; its leaves, of a lively green colour, are furnished with pinnæ, which are numerous and unusually narrow, whence its name; its flowers, produced on long racemi springing from the sides of the stem or branches, are of a singular dull red colour, and rarely followed by seeds with us.

It is a native of the Cape^[1], from whence it was introduced by Mr. MASSON, in 1774; flowers from June to October, and is usually increased by cuttings.

There is a description of this species in the *Mantiss. Pl. Linn.* but we know of no figure of it that has yet been published.

^[1] Ait. Kew.

[466]

CALYCANTHUS PRÆCOX. JAPAN ALL-SPICE.

Class and Order.

ICOSANDRIA POLYGYNIA.

Generic Character.

Cal. 1-phyllus, urceolatus, squarrosus: foliolis coloratis. *Cor.* calycina. *Styli* plurimi, stigmati glanduloso. *Sem.* plurima, caudata, intra calycem succulentum.

Specific Character and Synonyms.

CALYCANTHUS *præcox* petalis interioribus minutis. *Linn. Sp. Pl. ed. 3. p. 718. Ait. Kew. v. 2. p. 220. tab. x.*

OBAI s. *Robai*. Jasminum flore pleno suavi fœtido, fructu turbinato, semine phascoli. *Kæmpf. Amæn. exot. p. 878.*



N^o466.

The learned and instructive KÆMPFER in his *Amæn. Exot.* that vast fund of most useful information, gives a figure of this plant, in which it is represented both with flowers and seed-vessels, accompanied with a description and short account of it; from which we learn that it is cultivated in Japan as an ornamental plant, that the flowers are produced in February, before the leaves, that they have the scent of the violet, but become unpleasant on being long smelt to.

Hearing that Lord COVENTRY was the first who possessed this plant in England, I took the liberty of writing to his Lordship in January 1799, to request some information on this point, as well as some others relative to its culture, &c. On the 13th of the same month, his Lordship had the goodness to send me a beautiful specimen of the plant in bloom, a seedling plant one year old, together with a seed-vessel of the year 1798, and some seeds; in the Earl's letter is the following passage:—"the beauty of the *Calycanthus præcox* at this moment surpasses all description, it is covered with blossoms from top to bottom, and the fragrance of it may be perceived at the distance of fifty yards from the conservatory."

By his Lordship's direction, I received at the same time from his Gardener, Mr. WILLIAM DEAN, the following information, in answer to my queries:—"My Lord received the plant from China in 1766:—it was planted in a conservatory, is now sixteen feet high, and expands ten feet wide:—bears a succession of flowers from September to March:—the time of its first blowing I cannot precisely ascertain, but believe it to be nearly twenty years back:—it is propagated by layers, cuttings, and seeds, the latter it produces most years at Croome, but I believe at no other place in England:—there are plants of it at Croome six feet high, in a warm situation in the open border, which have stood out several years by being covered with a single mat in severe weather."

Not expecting to receive a plant from Lord COVENTRY in bloom, our drawing was made from one which flowered with Mr. WHITLEY, Nurseryman, Old-Brompton, December 22, 1798, and which came originally from Croome, his Lordship having presented most of the Nurserymen about town with plants of it; the blossoms of that from Croome were somewhat larger than those here represented, and the petals were less striped, indeed almost wholly tinged with purple, the leaves also proceeded more from the summit of the stalks and were of a much greener hue, owing no doubt to its being kept in the conservatory, while Mr. WHITLEY'S plant was tacked to the outside of the bottom of the greenhouse.

In the number of its stamina, which is rarely more than five, it does not accord with the character of the class icosandria, nor do the seeds agree with the generic character as described by LINNÆUS.

[467]

**DRACOCEPHALUM VIRGINIANUM. VIRGINIAN
DRAGON'S-HEAD.**

Class and Order.

DIDYNAMIA GYMNOSPERMIA.

Generic Character.

Cor. faux inflata: labium superius fornicatum.

Specific Character and Synonyms.

DRACOCEPHALUM *virginianum* foliis lineari-lanceolatis serratis, floribus confertis. *Linn. Syst. Nat. ed. 13. Gmel. p. 915.*

DRACOCEPHALUM *virginianum* floribus spicatis, foliis lanceolatis serratis. *Linn. Syst. Veg. ed. 14. Murr. p. 543.*

DRACOCEPHALUM *virginianum* floribus spicatis confertis, foliis lineari-lanceolatis serratis. *Ait. Kew. v. 2. p. 317.*



Nº467.

This elegant species of *Dracocephalum*, a native of Virginia, and other parts of North-America, is a hardy herbaceous plant, rising to the height of about two feet, and producing numerous flowers in long spikes, usually arranged on each side of the stalk.

It comes near to the *denticulatum* already figured, but differs in its superior height, the form of its leaves, the number of its flowers, and many other particulars.

It flowers from July to September, and with me has generally ripened much of its seed, from which it may be easily raised, as also by parting of its roots in spring or autumn; it succeeds best in a moist situation, and its stalks require to be carefully and timely sticked.

Was cultivated as long since as 1683, by Mr. JAMES SUTHERLAND.

[468]

ŒNOTHERA TETRAPTERA. WHITE-FLOWERED

ŒNOTHERA.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-fidus. *Petala* 4. *Caps.* cylindrica infera. *Sem.* nuda.

Specific Character and Synonyms.

ŒNOTHERA *tetraptera* foliis lanceolatis, basi pinnatifido-dentatis, capsulis pedicellatis obovatis quadrialatis. *Linn. Sp. Pl. edit. Willdenow.*

ŒNOTHERA foliis alternis subpinnatis crispis, flore mutabili fructû tetraptero. *Cavanill. Icon. t. 3. p. 40. tab. 279.*



Nº468.

Of this genus we have already figured six different species; this is another newly discovered one, which, like most of its kind, displays its beauties chiefly in the night. It is the only one, as far as we yet know, that has white blossoms; these, when first expanded, are beautifully so, but in the morning they change to a purple colour, fade, and their place is supplied by a fresh succession. In this remarkable change of colour, it bears some affinity to the *Ænothera anomala*, which may be considered as strengthening our opinion that the latter plant belongs to this genus rather than to that of *Gaura*. The *Ænothera tetraptera* is a native of Mexico, its duration as yet not certainly ascertained, but may be treated as a tender annual; and such plants as do not flower the first year, may be preserved under glasses through the winter. It was raised from seeds sent by Mr. DONN, from Cambridge; but was probably first introduced into this country from seeds sent to the Marchioness of BUTE, by Prof. ORTEGA, of Madrid.

INDEX.

In which the Latin Names of the Plants contained in the *Thirteenth Volume* are alphabetically arranged.

Pl.

- [457](#) Alöe plicatilis.
- [455](#) Alöe retusa.
- [453](#) Amaryllis Reginae.
- [462](#) Anthemis Pyrethrum.
- [441](#) Antholyza Merianella.
- [458](#) Aristea cyanea.
- [433](#) Azalea pontica.
- [451](#) Blakea trinervia.
- [466](#) Calycanthus præcox.
- [444](#) Canarina Campanula.
- [454](#) Canna indica.
- [452](#) Cardamine trifolia.
- [459](#) Convolvulus Cneorum.
- [445](#) Coronilla Emerus.
- [449](#) Dillenia speciosa.
- [456](#) Diosma serratifolia.
- [467](#) Dracocephalum virginianum.
- [463](#) Epidendrum ciliare.
- [440](#) Erica albens.
- [447](#) Erica empetrifolia.
- [443](#) Erica physodes.

INDEX.

In which the English Names of the Plants contained in the *Thirteenth Volume* are alphabetically arranged.

Pl.

- [466](#) Allspice, Japan.
- [455](#) Aloe, cushion.
- [457](#) Aloe, fan.
- [441](#) Antholyza, dwarf.
- [458](#) Aristea, grass-leaved.
- [433](#) Azalea, yellow.
- [444](#) Bell-flower, canary.
- [459](#) Bindweed, silvery-leaved.
- [451](#) Blakea, three-ribbed.
- [442](#) Broom, flax-leaved.
- [450](#) Corn-Flag, Watson's.
- [452](#) Cuckow-Flower, three-leav'd.
- [449](#) Dillenia, shewy.
- [456](#) Diosma, serrated or saw-leaved.
- [467](#) Dragon's Head, virginian.
- [463](#) Epidendrum, fringed.
- [448](#) Fig-Marygold, glittering.
- [435](#) Gnaphalium or Everlasting, heath-leaved.
- [447](#) Heath, crowberry-leaved.
- [443](#) Heath, sticky-flowered.

- [442](#) *Genista linifolia*.
[450](#) *Gladiolus Watsonius*.
[435](#) *Gnaphalium ericoides*.
[436](#) *Hibiscus præmorsus*.
[437](#) *Hydrangea Arborescens*.
[438](#) *Hydrangea Hortensis*.
[461](#) *Jasminum fruticans*.
[439](#) *Illicium floridanum*.
[465](#) *Indigofera angustifolia*.
[460](#) *Maurandya semperflorens*.
[448](#) *Mesembryanthemum micans*.
[468](#) *Ænothera tetraptera*.
[434](#) *Oxybaphus viscosus*.
[446](#) *Psoralea bracteata*.
[464](#) *Sisyrinchium gramineum*.
- [440](#) Heath, pallid.
[436](#) Hibiscus, bitten-leaved.
[438](#) Hydrangea, garden.
[437](#) Hydrangea, shrubby.
[461](#) Jasmine, yellow.
[439](#) Illicium, red-flowered or
 Aniseed tree.
[454](#) Indian Reed or Shot,
 common.
[465](#) Indigo, narrow-leaved.
[453](#) Lily, Mexican.
[460](#) Maurandya or Bastard
 Foxglove, climbing.
[468](#) Ænothera, white-flowered.
[462](#) Pellitory of Spain.
[446](#) Psoralea, oval-spiked.
[445](#) Senna, scorpion.
[464](#) Sisyrinchium, grass-leaved.
[434](#) Umbrella Wort, viscid.

London: Printed by STEPHEN COUCHMAN, Throgmorton-Street.

*** END OF THE PROJECT GUTENBERG EBOOK THE BOTANICAL
MAGAZINE, VOL. 13 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual

work is unprotected by copyright law in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)




[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)