


Catalogue of British Columbia Birds


Francis Kermode



Project Gutenberg

Catalogue of British Columbia Birds

Francis Kermode

The lower half of the cover features a vibrant pink background with a complex, abstract pattern of blue geometric shapes. This pattern includes various line segments, right angles, and triangles, some of which are solid blue while others are outlines. The shapes are arranged in a way that suggests a stylized, modern architectural or landscape design.

Project Gutenberg

The Project Gutenberg eBook of Catalogue of British Columbia Birds

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: Catalogue of British Columbia Birds

Author: Francis Kermode

Release date: August 1, 2004 [eBook #6329]

Most recently updated: December 29, 2020

Language: English

Other information and formats: www.gutenberg.org/ebooks/6329

Credits: Produced by Karen Fabrizius, David Starner and the Online
Distributed Proofreading Team

*** START OF THE PROJECT GUTENBERG EBOOK CATALOGUE
OF BRITISH COLUMBIA BIRDS ***

Produced by Karen Fabrizius, David Starner

and the Online Distributed Proofreading Team

Catalogue of British Columbia Birds.

PROVINCIAL MUSEUM, VICTORIA B. C.

PREFACE.

The present list is intended to include all birds which have been so far proven to occur in British Columbia.

I am fully aware of its incompleteness, and very much regret much of its lack of more definite information regarding the distribution of certain species, and with regard to that portion of the Province in north and northeastern interior, which, I have no doubt, is a valuable field for work.

Since the publication of the list of British Columbia birds in 1898 by the late John Fannin, which contained 339 species and sub-species, and with the information to hand, enables me to add to this 24.

I wish to extend my thanks especially to the following constant observers for more or less extended local lists, notes and specimens:—A. C. Brooks, Chilliwack, Okanagan and Cariboo; Rev. J. H. Keen, Queen Charlotte Islands and Metlakatla; Thos. Kermode, William Head Quarantine Station; E. P. VENABLE, Yernon; Chas. de Blois Green, Fairview and Keremeos.

I have also obtained valuable information from the check-list published by the late John Fannin.

FRANCIS KERMODE,
Curator Provincial Museum.

Victoria, B. C., August, 1904.

Order PYGOPODES. Diving birds.

Family PODICIPEDIDÆ. Grebes.

ÆCHMOPHORUS Coues.

1. Western Grebe.

Æchmophorus occidentalis (Lawr.).

A common winter resident along the coast of Vancouver Island and Mainland. Mr. Brooks says a few remain all winter at Okanagan Lake.

COLYMBUS Linnæus.

2. American Red-necked Grebe. "Holboell's Grebe."

Colymbus holboelli (Reinh.).

A common winter resident along the coast of Vancouver Island and Mainland. Winter resident on Okanagan Lake. (*Brooks.*) Not common at Metlakatla. (*Rev. J. H. Keen.*) Breeds on many of the lakes along the Cariboo Road above Clinton. (*Fannin.*)

3. Horned Grebe.

Colymbus auritus (Linn.).

A common winter resident on the coast, and very common in the straits near Victoria, in April. Brooks reports it at Okanagan Lake all winter.

Breeds on the lakes from Okanagan to Dense Lake. (*Fannin.*) Rare at Metlakatla. (*Rev. J. H. Keen.*)

4. American Eared Grebe.

Colymbus nigricollis californicus (Heerm.).

Taken at Kamloops, June, 1889. (*Spreadborough.*) Found breeding on lakes at Kamloops; Catalogue of Canadian Birds. (*Macoun.*)

PODILYMBUS Lesson.

5. Pied-billed Grebe.

Podilymbus podiceps (Linn.).

Common resident on Vancouver Island; it breeds on lakes close to Victoria. Common resident on Island and Mainland; breeds throughout its range. (*Fannin.*)

Family GAVIIDÆ Loons.

GAVIA Forster.

6. Great Northern Diver. Loon.

Gavia imber (Gunn.).

Common resident throughout the Province; breeds on Vancouver Island and Mainland.

7. Black-throated Loon.

Gavia arcticus (Linn.).

Taken at Burrard Inlet (*Fannin*), and at Dease Lake, Cassiar, by James Porter.

8. Pacific Loon.

Gavia pacificus (Lawr.).

Not common; one specimen taken at Comox by W. B. Anderson, and at Chilliwhack by Brooks; Victoria, May 13th, 1904, by Fred. Foster.

9. Red-throated Loon.

Gavia lumme (Gunn.).

I found this species quite common on Barkley Sound, V. I., April 19th, 1894, and fairly common near Esquimalt. Mr. Brooks reports it at Chilliwhack.

Family ALCIDÆ. Auks, Murres and Puffins.

LUNDA Pallas.

10. Tufted Puffin.

Lunda cirrhata (Pall.).

Common along the coast of Vancouver Island, Queen Charlotte Islands and Mainland; breeds on Bare Island, near Sidney, B. C.

FRATERCULA Brisson.

11. Horned Puffin.

Fratercula corniculata (Naum.).

Rare at Massett, Queen Charlotte Islands. (*Rev. J. H. Keen.*)

CERORHINCA Bonaparte.

12. Rhinoceros Auklet.

Cerorhinca monocerata (Pall.).

Coasts of Vancouver Island and Mainland. Breeds on islands in Gulf of Georgia.

PTYCHORAMPUS Brandt.

13. Cassin's Auklet.

Ptychorampus aleuticus (Pall.).

The entire coast line of the Province and West Coast of Vancouver Island. (*Fannin.*) This species was seen in the Gulf of Georgia, between Salt Spring Island and Nanaimo, May 7th, 1887. (*Macoun.*)

SYNTHLIBORAMPUS Brandt.

14. Ancient Murrelet.

Synthliborampus antiquus (Gmel.).

Not common. West Coast of Vancouver Island and taken in the Straits, near Victoria. Rev. J. H. Keen reports it rare on Queen Charlotte Islands.

BRACHYRAMPHUS Brandt.

15. Marbled Murrelet.

Brachyramphus marmoratus (Gmel.).

An abundant resident along the coast of British Columbia; have taken it at Skidegate, Queen Charlotte Islands, in August, 1895. Rare at Metlakatla. (Rev. J. H. Keen.) Breeds on Vancouver Island, and on some of the smaller islands in the Gulf of Georgia, and along many of the inlets of the Mainland. (Fannin.)

CEPPHUS Pallas.

16. Pigeon Guillemot.

Cephus columba (Pall.).

An abundant resident along the coast from Race Rocks to Alaska, and quite common in Skidegate Inlet, Q. C. I., August, 1895. Breeds throughout its range.

URIA Brisson.

17. California Murre, Guillemot.

Uria troile californica (Bryant).

A resident along the coast of Vancouver Island and Mainland. Breeds on West Coast of Vancouver Island.

Order LONGIPENNES. Long-winged Swimmers.

Family STERCORARIIDÆ. Skuas and Jaegers.

STERCORARIUS Brisson.

18. Pomarine Jaeger.

Stercorarius pomarius (Temm.).

One specimen taken near Victoria, October 22nd, 1898.

19. Parasitic Jaeger.

Stercorarius parasiticus (Linn.).

One specimen taken at Victoria by Wm. L. Gilchrist, November, 1897. Another specimen was taken at Comox by A. C. Brooks, September 12th, 1903.

20. Long-tailed Jaeger.

Stercorarius longicaudus (Vieill.).

Once shot and several times seen at Sumas Lake, September, 1890. (*Brooks*).

Family LARIDÆ. The Gulls and Terns.

PAGOPHILA Kaup.

21. Ivory Gull. Snow Gull.

Pagophila alba (Gunn.).

One specimen shot at Dease Lake, Cassiar, by James Porter, September, 1889. Another specimen was shot at Okanagan Lake in November, 1897, by Mr. J. T. Studley, and presented to the Museum. Mr. A. C. Brooks also reports it from Okanagan.

RISSA Stephens.

22. Pacific Kittiwake.

Rissa tridactyla pollicaris (Ridgw.).

Taken near Discovery Island, January, 1896, by W. Lindley, and at Queen Charlotte Islands by Dr. C. F. Newcombe, September, 1895.

LARUS Linnæus.

23. Glaucous-winged Gull.

Larus glaucesceus (Naum.).

An abundant resident on the coast; breeds on some of the islands in the Gulf of Georgia. Mr. Brooks reports it from Okanagan Lake.

24. Western Gull.

Larus occidentalis (Aud.).

An abundant resident on the coast during the winter months. Taken at Chilliwack. (A. C. Brooks.) Breeds in Similkameen Valley. (Fannin.)

25. American Herring Gull.

Larus argentatus smithsonianus (Coues.).

An abundant resident on the coast. It breeds on the coast and in the interior of Mainland. Common at Metlakatla. (Rev. J. H. Keen.) Chilliwack and Okanagan. (Brooks.)

26. California Gull.

Larus californicus (Lawr.).

A common winter resident on the coast. Common in the Lower Fraser Valley and on the Okanagan Lake in winter. (*Brooks.*)

27. Ring-billed Gull.

Larus delawarensis (Ord.).

Common along the coast of Vancouver Island and Mainland. Common in winter at Okanagan Lake and Chilliwack. (*Brooks.*) Breeds in the Interior, especially to the northward. (*Fannin.*)

28. Short-billed Gull.

Larus brachyrhynchus (Rich.).

Common winter resident along the coast of Vancouver Island and Mainland. Common in Lower Fraser Valley. (*Brooks.*) During the early part of May, 1891, I saw quite a number on the lakes of the Cariboo District, where it probably breeds. (*Fannin.*)

29. Heermann's Gull. White-headed Gull.

Larus heermanni (Cass.).

Common along the southern coast of Vancouver Island during July and August. Taken at William Head in full plumage, June 28th, 1904, by Thos. Kermode.

30. Bonaparte's Gull.

Larus philadelphia (Ord.).

An abundant resident throughout the Province, and found in great numbers about the inlets and rivers in March and April; it breeds in the interior of the Province. I have also seen it on the coast in June and July.

XEMA Leach.

31. Sabine's Gull.

Xema sabinii (Sab.).

Taken at Okanagan Lake, September 9th, 1897. (*Brooks.*)

STERNA Linnæus.

32. Common Tern.

Sterna hirundo (Linn.).

One specimen shot at Cowichan Gap by R. D. McClure, September 15th, 1896, and presented to the Museum.

33. Arctic Tern.

Sterna paradisæa (Brunn.).

From Dease Lake south, through the interior of the Mainland to Okanagan Lake. Two specimens—one taken at Dease Lake by James Porter, and another shot at Okanagan Lake by A. C. Brooks, September, 1897.

HYDROCHELIDON Boie.

34. Black Tern.

Hydrochelidon nigra surinamensis (Gmel.).

Not common. I observed it at Okanagan, June, 1895. Taken at Sumas and Okanagan Lake by A. C. Brooks, Burrard Inlet, Eraser River and interior of Mainland. (*Fannin.*)

Order TUBINARES. Tube-nosed Swimmers.

Family DIOMEDEIDÆ. Albatrosses.

DIOMEDEA Linnæus.

35. Black-footed Albatross.

Diomedea nigripes (Aud.).

West Coast of Vancouver Island. (*Fannin.*) Rare at Queen Charlotte Islands. (*Rev. J. H. Keen.*) One specimen taken near Nanaimo, June 3th, 1904.

36. Short-tailed Albatross.

Diomedea albatrus (Pall.)

In April, 1894, I found this species quite common in the Pacific Ocean, near Cape Beale.

Tolerably common on both coasts of Vancouver Island, but more abundant on the West Coast; a few have been taken in the straits off Victoria Harbour. (*Fannin.*)

Family PROCELLARIIDÆ. Fulmars & Shearwaters.

FULMARS Stephens.

37. Pacific Fulmar.

Fulmarus glacialis glupischa (Stejn.).

One specimen taken at Chemainus, Vancouver Island, November, 1895.
(Fannin.)

PUFFINUS Brisson.

38. Black-vented Shearwater.

Puffinus opisthomelas (Coues).

Not common; four specimens taken off Albert Head, October 24th, 1891.

39. Dark-bodied Shearwater.

Puffinus griseus (Gmel.).

During the fall of 1895 Dr. C. F. Newcombe found this species common on the West Coast of Queen Charlotte Islands. Since then one was picked up dead on the beach near Victoria.

40. Slender-billed Shearwater.

Puffinus tenuirostris (Temm.).

Not common; one specimen shot at William Head and presented to the Museum by T. Kermode, February 23rd, 1904.

OCEANODROMA Reichenbach.

41. Gray Fork-tailed Petrel.

Oceanodroma furcata (Gmel.).

Found along both coasts of Vancouver Island, but more common on the Pacific Coast; a few have been taken in the Straits near Victoria.

Very rare at Metlakatla; one picked up exhausted.

(Rev. J. H. Keen.)

42. Leach's Fork-tailed Petrel. White-rumped Petrel.

Oceanodroma leucorhoa (Vieill.) Coues.

West Coast of Vancouver Island. Four specimens taken in Hecate Strait, July, 1904. One specimen found dead on beach off Beacon Hill, Victoria, November, 1893.

Order STEGANOPODES. Totipalmate Swimmers.

Family PHALACROCORACIDÆ. Cormorants.

PHALACROCORAX Brisson.

43. White-crested Cormorant.

Phalacrocorax dilophus cinnatus (Brandt).

Common on both coasts of Vancouver Island and coast of Mainland, from Race Rocks to Alaska.

44. Brandt's Cormorant. Pencilled Cormorant.

Phalacrocorax penicillatus (Brandt).

Two specimens were killed in the Straits, near Victoria, by D. E. Campbell, April 19th, 1897. Later in the month another specimen was found on the beach at Beacon Hill.

45. Violet-green Cormorant.

Phalacrocorax pelagicus robustus (Ridgw.).

This is the most abundant Cormorant in the Province, and is found along both coasts of the Island, and has been taken as far north as Port Simpson. Breeds on islands close to Sidney Island. Common at Metlakatla. (*Rev. J. H. Keen.*)

Family PELECANIDÆ. Pelicans.

PELECANUS Linnæus.

46. American White Pelican.

Pelecanus erythrorhynchos (Gmel.).

Not common. Four specimens taken—Sicamous, October, 1890; Alexandria Bridge, September, 1895; Tranquille, August, 1897; Comox, September 8th, 1900. Mr. Brooks reports it from Chilliwhack and Okanagan. Said to breed in the Chilcotin country. (*Fannin.*)

47. California Brown Pelican.

Pelecanus californicus (Ridgw.).

Not common. One specimen in the Museum killed at Race Rocks, January, 1897. Seen at Sumas Lake. (*Brooks.*)

Order ANSERES. Lamellirostral Swimmers.

Family ANATIDÆ. Mergansers, Ducks, Geese and Swans.

MERGANSER Brisson.

48. American Merganser.

Merganser americanus (Cass.).

Found throughout the Province, have taken it on West Coast of Vancouver Island in May; also on Queen Charlotte Islands in July, 1895, I took four young birds. Brooks reports it found on Okanagan Lake all winter.

49. Red-breasted Merganser.

Menganser serrator (Linn.).

Found distributed throughout the Province. I saw them at Sumas Lake May 17th, 1904. Rev. J. H. Keen says common at Metlakahtla, Chilliwhack and Okanagan. (*Brooks.*)

LOPHODYTES Reichenbach.

50. Hooded Merganser.

Lophodytes cucullatus (Linn.).

This species is common throughout the Province and on the Pacific Coast. Brooks has taken it at Chilliwhack and Okanagan.

ANAS Linnæus.

51. Mallard.

Anas boschas (Linn.).

An abundant resident throughout the Province; breeds in suitable localities throughout its range.

CHAULELASMUS Bonaparte.

52. Gadwall. Gray Duck.

Chaulelasmus strepera (Linn.).

Not common; a few have been taken near Victoria. Mr. Brooks reports it from Chilliwhack and Okanagan.

MARECA Stephens.

53. European Widgeon.

Mareca penelope (Linn.).

Rare. Two specimens, one taken near Victoria February 9th, 1899, and another near Saanich.

54. American Widgeon. Baldpate.

Mareca americana (Gmel.).

A common winter resident on the coast. Brooks reports it from Chilliwhack and Okanagan. Breeds on lakes on Cariboo Road above Clinton. (*Fannin.*)

NETTION Kaup.

55. Green-winged Teal.

Nettion carolinensis (Gmel.).

An abundant resident, Breeds in the interior of the Mainland. Common on the coast throughout the winter. Rare on Queen Charlotte Islands. (*Rev. J. H. Keen.*)

QUERQUEDULA Stephens.

56. Blue-winged Teal.

Querquedula discors (Linn.).

Not common on the coast; a few are taken every year. Taken at Chilliwhack. (*Brooks.*)

57. Cinnamon Teal.

Querquedula cyanoptera (Vieill.).

A summer visitor in the interior of the Province. Brooks has taken it at Chilliwhack and Okanagan.

SPATULA Boie.

58. Shoveller. Spoon-bill.

Spatula clypeata (Linn.).

A common resident on the Mainland, rarely met with on Vancouver Island. Brooks says common resident in Lower Fraser Valley about Sumas Lake, has also taken it on Okanagan.

DAFILA Stephens.

59. Pintail. Sprigtail.

Dafila acuta (Linn.).

An abundant winter resident on the coast. Breeds in the interior of the Mainland.

AIX Boie.

60. Wood Duck.

Aix sponsa (Linn.).

Not common. A summer resident on Island and Mainland. Brooks has taken it at Chilliwack and Okanagan.

AYTHYA Boie.

61. Red-head. Pochard.

Aythya americana (Eyt.).

A winter resident on the coast, but nowhere common; a few are taken every season. Fannin found it breeding on the lakes along the Cariboo Road.

62. Canvas-back Duck.

Aythya vallisneria (Wils.).

A winter resident on the coast. Brooks has taken it on Okanagan Lake. I found them in pairs on the lakes near 108-Mile House, Cariboo Road, in April and May, 1891, where I was told they breed. (*Fannin*).

63. American Scaup Duck. Blue-bill.

Aythya marila (Linn.).

An abundant winter resident on the coast. Breeds in the interior of Mainland.

64. Lesser Scaup Duck.

Aythya affinis (Eyt.).

Not common on the coast. Fannin has taken it in May near Ashcroft, and Brooks has seen it all winter at Okanagan.

65. Ring-necked Duck.

Aythya collaris (Donov.).

Not common. A few have been taken on Vancouver Island. Mr. Brooks says common in Lower Fraser Valley, and is found all winter on Okanagan Lake.

CLANGULA Leach.

66. American Golden-eye. Whistler.

Clangula clangula americana (Faxon.)

A common winter resident on the coast. Fannin says breeds in interior. Brooks has taken it at Chilliwack and Okanagan.

67. Barrow's Golden-eye.

Clangula islandica (Gmel.).

Not common. It winters on the coast; a few have been taken on Vancouver Island. Brooks has taken it at Okanagan. Fannin found it plentiful on the lakes along the Cariboo Road, May, 1891.

CHARITONETTA Stejneger.

68. Buffle-head. Butter-ball.

Charitonetta albeola (Linn.).

An abundant winter resident on the coast. Breeds in the interior of the Mainland. Common at Metlakatla. (Rev. J. H. Keen.)

HARELDA Stephens.

69. Long-tailed Duck. Old Squaw.

Harelda hyemalis (Linn.).

Common; the coasts of Vancouver Island and Mainland. Brooks has taken it at Chilliwack. Rev. J. H. Keen reports it common at Metlakatla.

HISTRIONICUS Lesson.

70. Harlequin Duck.

Histrionicus histrionicus (Linn.).

A common resident on the coast. I found it breeding at Bear River, Vancouver Island, May, 1894. Brooks has taken it at Chilliwhack. Rare at Queen Charlotte Islands. (Rev. J. H. Keen.)

OIDEMIA Fleming.

71. American Black Scoter.

Oidemia americana (Swains.).

Not common; has been taken at Victoria and Port Simpson. Fannin saw a small flock on a lake near 108-Mile House, Cariboo Road, 10th May, 1891. Fairly common at Matlakatla. (Rev. J. H. Keen.)

72. White-winged Scoter.

Oidemia deglandi (Bonap.).

An abundant resident, and is found on the coast throughout the year. Brooks reports it from Chilliwhack and Okanagan. Rev. J. H. Keen says is common at Metlakatla.

73. Surf Scoter. Sea Coot.

Oidemia perspicillata (Linn.).

Abundant resident along the coasts of Vancouver Island and Mainland. Found at Chilliwhack and Okanagan. (Brooks.) Common at Metlakatla. (Rev. J. H. Keen.)

ERISMATURA Bonaparte.

74. Ruddy Duck.

Erismatura jamaicensis (Gmel.).

Not common on the Coast; a few are taken in the winter. Fannin found it breeding in the lakes along the Cariboo Road, May, 1891. Brooks has seen it at Chilliwhack and Okanagan.

CHEN Boie.

75. Lesser Snow Goose.

Chen hyperborea (Pall.).

A winter resident on the Coast. Brooks has seen it at Chilliwhack. During some winters large numbers congregate off the mouth of Fraser River. (*Fannin.*)

76. Ross's Snow Goose.

Chen rossii (Cassin.).

This is a rare bird in B. C. One specimen taken at Comox, January, 1894; another was taken at Kuper Island, April, 1895.

ANSER Brisson.

77. American White-fronted Goose.

Anser albifrons gambeli (Hartl.).

Abundant on the Coast in winter. Breeds both on Island and Mainland. The young have been taken on Cowichan Lake. (*Fannin.*) Brooks has taken it at Sumas and Okanagan Lakes. Common at Metlakatla. (*Rev. J. H. Keen.*)

BRANTA Scopoli.

78. Canada Goose.

Branta canadensis (Linn.).

An abundant winter resident on the Coast. Breeds in the interior of the Mainland; the eggs have been taken at Penticton by C. DeBlois Green.

79. Hutchin's Goose.

Branta canadensis hutchinsii (Rich.).

Abundant in the spring and fall migrations, and winters on the Coast.

80. White-cheeked Goose.

Branta canadensis occidentalis (Baird).

Taken at Chilliwhack by A. C. Brooks.

81. Cackling Goose.

Branta canadensis minima (Ridgw.).

Winter resident on the Coast; has been taken at Elk Lake, near Victoria. Brooks reports it from Chilliwhack and Okanagan.

82. Brant.

Branta bernicla glaucogastra (Brehm.).

Two specimens shot at Comox by A. C. Brooks, January 10th, 1904.

83. Black Brant.

Branta nigricans (Lawr.).

An abundant winter resident on the Coast. In April, 1894. I saw them in great numbers off Barkley Sound migrating north.

PHILACTE.

84. Emperor Goose.

Philacate canagica (Sevast.).

One specimen shot at Chemainus, December, 1894.

OLOR Wagler.

85. Whistling Swan.

Olor columbianus (Ord.).

A winter resident on Vancouver Island and southern Mainland. Mr. Brooks records it from Chilliwhack and Okanagan.

86. Trumpeter Swan.

Olor buccinator (Rich.).

I have only seen this bird at Deaso Lake, Cassiar. (*Fannin.*)
Mr. Brooks has found it at Sumas and Okanagan.

Order HERODIONES. Herons, Ibises, etc.

Family IBIDIDÆ. Ibises.

PLEGADIS Kaup.

87. White-faced Glossy Ibis.

Plegadis guarauna (Linn.).

Only two specimens known to be taken in the Province, one on Salt Spring Island, the other at mouth of Fraser River. Check List of B. C. Birds, 1891. (*Fannin.*)

Family ARDEIDÆ. Herons and Bitterns.

BOTAURUS Hermann.

88. American Bittern.

Botaurus lentiginosus (Montag.).

Common throughout the greater portion of the Province. Breeds both east and west of Cascades.

ARDEA Linnæus.

89. Great Blue Heron.

Ardea herodias (Linn.).

A common resident on the coast, and is quite common at Sumas Lake and Okanagan. Breeds throughout its range.

90. Northwest Coast Heron.

Ardea herodias fannini (Chapman).

In August, 1895, I found this form, now named after the late John Fannin by Frank M. Chapman, quite common at Skidegate, Queen Charlotte Islands.

EGRETTA Forester.

91. Snowy Heron.

Egretta candidissima (Gmel.).

Rare in British Columbia. Two specimens collected at Burrard Inlet, May, 1879. (*Fannin.*) One specimen is in the Museum.

Order PALUDICOLÆ. Cranes, Rails, etc.

Family GRUIDÆ. Cranes.

GRUS Pallas.

92. Little Brown Crane.

Grus canadensis (Linn.).

Common, during migrations, throughout the Province.

93. Sandhill Crane.

Grus mexicana (Mull.).

Common throughout the Province; it breeds in the interior of Mainland; numbers pass over Victoria in the spring and autumn migrations.

Family RALLIDÆ. Rails and Coots.

RALLUS Linnæus.

94. Virginia Rail.

Rallus virginianus (Linn.).

Tolerably common on Island and Mainland; breeds close to Victoria. Brooks has taken it at Chilliwhack and Okanagan.

PORZANA Vieillot.

95. Carolina Rail. Sora.

Porzana Carolina (Linn.).

Found on Vancouver Island and Mainland. Rev. J. H. Keen reports it very rare on Queen Charlotte Islands. Common east of Cascades. (*Fannin.*)

FULICA Linnæus.

96. American Coot. Mud-hen.

Fulica americana (Gmel.).

A common resident on Island and Mainland; breeds throughout its range.

Order LIMICOLÆ. Shore Birds.

Family PHALAROPODIDÆ. Phalaropes.

CRYMOPHILUS Vieillot.

97. Red Phalarope.

Crymophilus fulicarius (Linn.).

Taken at Clover Point, near Victoria, during migration.

PHALAROPUS Brisson.

98. Northern Phalarope.

Phalaropus lobatus (Linn.).

Abundant along the coast of Island and Mainland in the spring and autumn. Brooks has taken it at Chilliwack and Okanagan. Fannin has taken it at Burrard Inlet in July.

STEGANOPUS. Vieillot.

99. Wilson's Phalarope.

Steganopus tricolor (Vieill.).

Taken at Chilliwack by A. C. Brooks.

Family SCOLOPACIDÆ. Snipes, Sand-pipers, Etc.

GALLINAGO Leach.

100. Wilson's Snipe.

Gallinago delicata (Ord.)

Common throughout the Province on Island and Mainland; breeds in the interior. Rev. J. H. Keen reports it common at Metlakatla.

MACRORHAMPHUS Leach.

101. Long-billed Dowitcher. Red-breasted Snipe.

Macrorhamphus scolopaceus (Say.).

Tolerably abundant throughout the Province. I found it quite common at Clayoquot, West Coast Vancouver Island, in May, 1894. Fannin says breeds in the interior. Brooks has taken it at Chilliwack. Fairly common at Metlakatla. (*Rev. J. H. Keen.*)

MICROPALAMA Baird.

102. STILT SANDPIPER.

Micropalama himantopus (Bonap.).

Taken at Chilliwack, August 19th, 1899, by A. C. Brooks.

TRINGA Linnæus.

103. KNOT. ROBIN SNIPE.

Tringa canutus (Linn.).

Abundant during migrations, chiefly along the coast. (*Fannin.*)
Taken at Chilliwack. (*Brooks.*)

ACTODROMAS Kaup.

104. SHARP-TAILED SANDPIPER.

Actodromas acuminata (Horsf.).

Two specimens taken at Massett, Queen Charlotte Islands, December 27th, 1897, by Rev. J. H. Keen.

105. PECTORAL SANDPIPER.

Actodromas maculata (Vieill.).

Not common; taken along the coast during migrations. Brooks has taken it at Chillihwack and Okanagan.

106. BAIRD'S SANDPIPER.

Actodromas bairdii (Coues.).

Distributed along the coast of Island and Mainland, Rev. J. H. Keen reports it rare at Metlakatla. Taken at Chillihwack and Okanagan (*Brooks*); and at Clover Point, Victoria (*E. M. Anderson*).

107. LEAST SANDPIPER.

Actodromas minutilla (Vieill.).

Common along the coast of Island and Mainland. I found it common at Clayoquot in May, 1894; I have also taken it near Victoria in July. Brooks has taken it at Chillihwack and Okanagan. Fannin has taken it at 108-Mile House, Cariboo Road, in May.

PELIDNA Cuvier.

108. RED-BACKED SANDPIPER.

Pelidna alpina sakhalina (Vieillot).

Common in the spring and autumn migration along the coast of Island and Mainland. Brooks has taken it at Chilliwack.

EREUNETES Illiger.

109. SEMIPALMATED SANDPIPER.

Ereunetes pusillus (Linn.).

Not uncommon in migration along the Coast. Chilliwack and Okanagan. (Brooks.) Common at Metlakatla. (Rev. J. H. Keen.)

110. WESTERN SEMIPALMATED SANDPIPER.

Ereunetes pusillus occidentalis (Lawr.).

Abundant in the fall along the Coast. I also found it very common at Clayoquot, May, 1894, and have taken it near Victoria in July. Brooks has taken it at Chilliwack.

CALIDRIS Cuvier.

111. SANDERLING.

Calidris arenaria (Linn.).

Not common; several have been taken near Victoria by A. H. Maynard. Brooks has taken it at Chilliwack. Fairly common at Metlakatla. (Rev J. H. Keen.)

LIMOSA Brisson.

112. MARBLED GODWIT.

Limosa fedoa (Linn.).

The whole of British Columbia; breeds chiefly east of Cascades.
(*Fannin.*) Port Simpson. (*W. B. Anderson.*)

TOTANUS Bechstein.

113. GREATER YELLOW-LEGS.

Totanus melanoleucus (Gmel.).

Common along the coast in winter; have taken it at Clayoquot in May. Brooks has taken it at Chilliwack and Okanagan. I found it breeding around the lakes at Clinton, May, 1891. (*Fannin.*)

114. LESSER YELLOW-LEGS.

Totanus flavipes (Gmel.).

Tolerably common through the Province; winters on the coast. Brooks has taken it at Chilliwack and Okanagan.

HELODROMAS Kaup.

115. SOLITARY SANDPIPER.

Helodromas solitarius (Wils.).

Found throughout the Province. Taken at Chilliwack and Okanagan.
(*Brooks.*)

116. WESTERN SOLITARY SANDPIPER.

Helodromas solitarius cinnamomeus (Brewster).

Taken at Chilliwhack and Okanagan. (*Brooks*).

SYMPHEMIA Rafinesque.

117. WESTERN WILLET.

Symphemia semipalmata inornata (Brewster).

One specimen shot at Clover Point, near Victoria, August 18th, 1898, by Mr. J. Henley.

HETERACTITIS Stejneger.

118. WANDERING TATLER.

Heteractitis incanus (Gmel.).

Tolerably common along the coast of Island and Mainland. I took it at Clayoquot, May, 1894, and at Skidegate, Queen Charlotte Islands, August 5th, 1895.

BARTRAMIA Lesson.

119. BARTRAMIAN SANDPIPER.

Bartramia longicauda (Bechst.).

One specimen shot at Comox, August 20th, 1895, by W. B. Anderson. Another was taken at 150-Mile House, Cariboo Road, May 16th, 1901, by A. C. Brooks.

TRYNGITES Cabanis.

120. BUFF-BREASTED SANDPIPER.

Tryngites subruficollis (Vieill.).

Not common. Taken at Chilliwack by A. C. Brooks.

ACTITIS Illiger.

121. SPOTTED SANDPIPER.

Actitis macularia (Linn.).

This bird is found along the entire coast; I have taken it at Clayoquot Vancouver Island, and at Skidegate, Queen Charlotte Islands, July 26th, 1895. Brooks has taken it at Chilliwack and Okanagan.

NUMENIUS Brisson.

122. LONG-BILLED CURLEW.

Numenius longirostris (Wils.).

A few specimens have been taken on the coast, but is to be more frequently met with in the interior of the Mainland, south through the Okanagan.

123. HUDSONIAN CURLEW.

Numenius hudsonicus (Lath.).

Not common, but distributed along the coast of Island and Mainland. I have taken it at Clayoquot in May, 1894. W. B. Andersen has taken it at Port Simpson, and J. Maynard at Cadboro Bay, near Victoria.

Family CHARADRIIDÆ. Plovers, Etc.

CHARADRIUS Linnæus.

124. BLACK-BELLIED PLOVER.

Charadrius squatarola (Linn.).

Abundant during migrations along the coast. Brooks has taken it at Chilliwhack and Okanagan.

125. AMERICAN GOLDEN PLOVER.

Charadrius dominicus (Mull.).

Common on the coast during migrations. Brooks has taken it at Chilliwhack and Okanagan. Rev. J. H. Keen reports it common at Metlakatla. W. B. Anderson has taken it at Port Simpson.

ÆGIALITIS Boie.

126. KILLDEER PLOVER.

Ægialitis vocifera (Linn.).

Found throughout the Province; occasionally on the coast in winter. Brooks has taken it at Chilliwhack and Okanagan.

127. SEMIPALMATED PLOVER.

Ægialitis semipalmata (Bonap.).

Not common. I took two specimens on the West Coast, Vancouver Island—one at Ucluelet, April 28th, 1894, the other at Clayoquot, May, 1894. A. H. Maynard shot one at Cadboro Bay, Victoria.

Family APHRIZIDÆ. Surf-birds and Turnstones.

APHRIZA Audubon.

128. SURF-BIRD.

Aphriza virgata (Gmel.).

Not uncommon along the entire coast line of the Province. I found it quite common at Clayoquot, May, 1894. It has been taken at Port Simpson by W. B. Anderson.

ARENARIA Brisson.

129. TURNSTONE.

Arenaria interpres (Linn.).

Along the entire coast line, but not common. It has been taken at Port Simpson and Sidney Island. Brooks has taken it at Sumas Lake.

130. BLACK TURNSTONE.

Arenaria melanocephala (Vig.).

Common along the entire coast of the Province. I have taken it at Clayquot in May, 1894, and I found it very common at Skidegate, Queen Charlotte Islands, in July, 1895, and it is found near Victoria throughout the winter.

Family HÆMATOPODIDÆ. Oyster-Catchers.

HÆMATOPUS Linnæus.

131. BLACK OYSTER-CATCHER.

Hæmatopus bachmani (Aud.).

Abundant along the entire coast of Island and Mainland. Breeds throughout its range. Common on Queen Charlotte Islands, at Skidegate.

Order GALLINÆ. Gallinaceous Birds.

Family TETRAONIDÆ. Grouse, Partridges, Etc.

OREORTYX Baird.

132. MOUNTAIN PARTRIDGE.

Oreortyx pictus (Dougl.).

Common on Vancouver Island. Introduced from California.

LOPHORTYX Bonaparte.

133. CALIFORNIA PARTRIDGE.

Lophortyx californicus (Shaw.).

Common on Vancouver Island. Introduced from California.

DENDRAGAPUS Elliot.

134. SOOTY GROUSE.

Dendragapus obscurus fuliginosus (Ridgw.).

Abundant west of Cascade Mountains, including Vancouver Island, Queen Charlotte Islands, and all the larger islands along the coast.

135. RICHARDSON'S GROUSE.

Dendragapus obscurus richardsonii (Dougl.).

An abundant resident east of Cascade Mountains to Rocky Mountains.

CANACHITES Stejneger.

136. FRANKLIN'S GROUSE.

Canachites franklinii (Dougl.).

An abundant resident throughout the wooded portion of the interior east of the Cascade Mountains, from Okanagan to Cassiar.

BONASA Stephens.

137. CANADIAN RUFFED GROUSE.

Bonasa umbellus togata (Linn.).

An abundant resident east of and including the Cascade Mountains. Taken at Chilliwhack by A. C. Brooks.

138. GRAY RUFFED GROUSE.

Bonasa umbellus umbelloides (Dougl.).

Rocky Mountain District, Soda Creek and Beaver Pass. Check List of B. C. Birds, 1898. (*Fannin.*) Okanagan, A. C. Brooks.

139. OREGON RUFFED GROUSE.

Bonasa umbellus sabini (Dougl.).

An abundant resident on Vancouver Island and all the larger islands of the coast, and on the Mainland west of Cascade Mountains.

LAGOPUS Brisson.

140. WILLOW PTARMIGAN.

Lagopus lagopus (Linn.).

Northern portion of British Columbia, Dease Lake, Cassiar. (*James Porter.*) Log Cabin, Atlin. (*Fletcher and Englehardt.*)

141. ROCK PTARMIGAN.

Lagopus rupestris (Gmel.).

Common on the summit of most of the mountains on the Mainland and Vancouver Island. (*Fannin.*) Quite common at Atlin.

142. WHITE-TAILED PTARMIGAN.

Lagopus leucurus (Swains. & Rich.).

Found on the summit of most mountains on the Mainland except the Coast Range. Taken at Okanagan. (*Brooks.*) Cassiar. (*James Porter.*)

PEDIOECETES Baird.

143. COLUMBIAN SHARP-TAILED GROUSE. PRAIRIE CHICKEN.

Pedioecetes phasianellus columbianus (Ord.).

An abundant resident east of Cascade Range through the southern portions of the Province. E. P. VENABLE reports it common near Vernon.

CENTROCERCUS Swainson.

144. SAGE GROUSE.

Centrocercus urophasianus (Bonap.).

Three specimens taken by Mr. G. B. Martin at Osoyoos Lake in October, 1864. (Check List of B. C. Birds, 1891. *Fannin*.)

Family PHASIANIDÆ. Pheasants.

PHASIANUS Linnæus.

145. RING-NECKED PHEASANT.

Phasianus torquatus (Linn.).

Abundant on Vancouver Island and in the Lower Fraser River Valley and other portions of the Mainland. Introduced from China; now thoroughly naturalised.

Order COLUMBÆ. Pigeons.

Family COLUMBIDÆ. Pigeons.

COLUMBA Linnæus.

146. BAND-TAILED PIGEON.

Columba fasciata (Say.).

A common summer resident in the south-western portions of the Province including Vancouver Island.

ECTOPISTES Swainson.

147. PASSENGER PIGEON.

Ectopistes migratorius (Linn.).

Mentioned in John Keast Lord's "Naturalist in British Columbia" (1866). If it ever did occur here, it is now, without doubt, extinct. (*Fannin.*)

ZENAIDURA Bonaparte.

148. MOURNING DOVE.

Zenaidura macroura (Linn.).

Not common on Vancouver Island. I found it quite common in the Okanagan in June, 1895. Brooks has taken it at Chilliwhack. Rev. J. H. Keen reports it rare at Metlakatla.

Order RAPTORES. Birds of Prey.

Family CATHARTIDÆ. American Vultures.

GYMNOGYPS Lesson.

149. CALIFORNIA VULTURE.

Gymnogyps californianus (Shaw.).

In September, 1880, I saw two of these birds at Burrard Inlet. It is more than probable they are accidental visitants here. Lord says: Mouth of Fraser River; seldom visits the Interior. (*Fannin.*)

CATHARTES Illeger.

150. TURKEY VULTURE.

Cathartes aura (Linn.).

Tolerably common on Vancouver Island and portions of the Mainland. Brooks has taken it at Chilliwhack and Okanagan.

Family FALCONIDÆ. Falcons, Hawks, Eagles, etc.

CIRCUS Lacépède.

151. MARSH HAWK.

Circus hudsonius (Linn.).

Common on Vancouver Island and Mainland. Abundant on the Mainland. Partially migratory; a few are found throughout the winter at Ladners. (*Fannin.*) Chilliwhack and Okanagan. (*Brooks.*)

ACCIPITER Brisson.

152. SHARP-SHINNED HAWK.

Accipiter velox (Wils.).

Abundant on Vancouver Island and portions of the Mainland. Brooks has taken it at Chilliwack and Okanagan.

153. COOPER'S HAWK.

Accipiter cooperii (Bonap.).

Found on Vancouver Island and Mainland; have taken it at Sicamous. Brooks has taken it at Chilliwack and Okanagan.

154. AMERICAN GOSHAWK.

Accipiter atricapillus (Wils.).

Rather common in interior of Province. A few have been taken on Vancouver Island. Brooks has taken it at Chilliwack and Okanagan.

155. WESTERN GOSHAWK.

Accipiter atricapillus striatulus (Ridgw.).

Common throughout the Province. I found it very common at Skidegate, Queen Charlotte Islands, in August, 1895.

BUTEO Cuvier.

156. WESTERN RED-TAILED HAWK.

Buteo borealis calurus (Cass.).

An abundant resident on Vancouver Island and coast of Mainland. Taken at Chilliwack and Okanagan. (*Brooks.*) Rare at Massett, Queen Charlotte Islands. (*Rev. J. H. Keen.*)

157. RED-BELLIED HAWK.

Buteo lineatus elegans (Cass.).

Not common. I have no record of it on Vancouver Island. It has been taken at Burrard Inlet by J. Fannin, and at Chilliwack by A. C. Brooks.

158. SWAINSON'S HAWK.

Buteo swainsoni (Bonap.).

Not common on Vancouver Island; it has been taken at Victoria and Union. Brooks has taken it at Chilliwack and Okanagan.

ARCHIBUTEO Brehm.

159. AMERICAN ROUGH-LEGGED HAWK.

Archibuteo lagopus sancti-johannis (Gmel.).

Not common. Found on both Vancouver Island and Mainland. Brooks has taken it at Chilliwack and Okanagan.

AQUILA Brisson.

160. GOLDEN EAGLE.

Aquila chrysaetos (Linn.).

Found throughout the entire Province. Rare on Vancouver Island. A resident.

HALIÆTUS Savigny.

161. BALD EAGLE.

Haliaeetus leucocephalus alascanus (C. H. Townsend).

An abundant resident throughout the entire Province, on Island and Mainland. Breeds throughout its range. Common on Queen Charlotte Islands.

FALCO Linnæus.

162. GYRFALCON.

Falco rusticolus gyrfalco (Linn.).

One specimen shot at Comox, October, 1894, by W. Harvey. Brooks has taken it at Chilliwack.

163. PRAIRIE FALCON.

Falco mexicanus (Schleg.).

Taken at Chilliwack by A. C. Brooks, and at Kamloops, November, 1894, by W. Fortune.

164. DUCK HAWK.

Falco peregrinus anatum (Bonap.).

Apparently more common on the Mainland than Vancouver Island. Several have been taken in the Okanagan Country.

165. PEALE'S FALCON.

Falco peregrinus pealei (Ridgw.).

Found chiefly along the coasts of Mainland and Vancouver Island. Several specimens have been taken at Cadboro Bay, near

Victoria. Mr. Brooks has taken it at Chilliwhack. Rare at Metlakatla. (*Rev. J. H. Keen.*)

166. PIGEON HAWK.

Falco columbarius (Linn.).

Apparently more common east of the Coast Range. A few have been taken on the Coast. Brooks has taken it at Chilliwhack and Okanagan.

167. BLACK MERLIN.

Falco columbarius suckleyi (Ridgw.).

Abundant; summer resident on Vancouver Island and portions of Mainland. Mr. Brooks has taken it at Chilliwhack and Okanagan.

168. RICHARDSON'S MERLIN.

Falco richardsonii (Ridgw.).

Not common on Vancouver Island; a few specimens have been taken near Victoria. Brooks has taken it at Chilliwhack and Okanagan.

169. DESERT SPARROW HAWK.

Falco spaverius phaloena (Lesson).

Abundant throughout the southern portion of Mainland, including Vancouver Island, from the coast to Rocky Mountains.

PANDION Savigny.

170. AMERICAN OSPREY. FISH HAWK.

Pandion halioetus carolinensis (Gmel.).

Common throughout the Province, on Mainland and Island, including Queen Charlotte Islands. Their nest is a huge heap of sticks, usually on the broken top of a tree.

Family BUBONIDÆ. Horned Owls, etc.

ASIO Brisson.

171. AMERICAN LONG-EARED OWL.

Asio wilsonianus (Less.).

Not common; a few have been taken on Vancouver Island. Mr. Brooks has taken it at Chilliwack and Okanagan.

172. SHORT-EARED OWL.

Asio accipitrinus (Pall.).

Common throughout the Province, on Island and Mainland; rather common near Victoria throughout the winter.

SYRNIUM Sanigny.

173. NORTHERN SPOTTED OWL.

Syrnium occidentale caurinum (Merriam).

Rare resident at Chilliwack and Mount Lehman (*Brooks*.) The late John Fannin had this bird on his first list of B. C. birds, he having taken it at

Burrard Inlet some years ago, but in his lists of 1891 and 1898 he dropped it.

SCOTIAPTEX Swainson.

174. GREAT GREY OWL.

Scotiaptex nebulosa (Forster).

A rare bird in this Province; it has been taken at Vernon and Stewart's Lake. Brooks has taken it at Chilliwack.

CRYPTOGLAUX Richmond.

175. RICHARDSON'S OWL.

Cryptoglaux tengmalmi richardsoni (Bonaparte).

Taken at Okanagan by A. C. Brooks.

176. SAW-WHET OWL.

Cryptoglaux acadica (Gmel.).

Not common on Vancouver Island; several have been taken near this city. Tolerably at Chilliwack and common in winter at Okanagan. (*Brooks.*)

176A. NORTH-WEST SAW-WHET OWL.

Cryptoglaux acadica scotæa (Osgood).

Queen Charlotte Islands.

MEGASCOPS Kaup.

177. KENNICOTT'S SCREECH OWL.

Megascops asio kennicottii (Elliot).

An abundant resident throughout the Province. Breeds in the neighbourhood of Victoria.

178. MACFARLANE'S SCREECH OWL.

Megascops asio macfarlanei (Brewst.).

Southern portion of the Province, east of Cascade Mountains.

BUBO Dumeril.

179. GREAT HORNED OWL.

Bubo virginianus (Gmel.).

Taken at Chilliwhack by A. C. Brooks, and near Victoria by John Hall, November, 1896.

180. WESTERN HORNED OWL.

Bubo virginianus pallescens (Stone).

Common on Vancouver Island and Mainland. Taken at Chilliwhack and Okanagan. (*Brooks.*)

181. ARCTIC HORNED OWL.

Bubo virginianus arcticus (Swains.).

One specimen taken near Victoria, November, 1896, by A. H. Maynard, and at Chilliwhack by A. C. Brooks.

182. DUSKY HORNED OWL.

Bubo virginianus saturatus (Ridgw.).

An abundant resident in this Province; common on Vancouver Island. Brooks has taken it at Chilliwack and Okanagan.

NYCTEA Stephens.

183. SNOWY OWL.

Nyctea nyctea (Linn.).

A resident in the northern portions of the Province, south in winter to mouth of Fraser River, Chilliwack and Okanagan, occasionally to Vancouver Island. Rev. J. H. Keen reports it rare at Metlakatla.

SURNIA Duméril.

184. American Hawk Owl.

Surnia ulula caparoch (Mull.).

Not common, but found throughout the southern part of the Province from the Coast to Okanagan. Rare on Vancouver Island. It has been taken at Victoria, Sardis, Grand Prairie and Vernon.

SPEOTYTO Gloger.

185. Burrowing Owl.

Speotyto cunicularia hypogaea (Bonap.).

Not common, but is found in suitable localities in the southern portions of the Province from Chilliwack to Okanagan. Mr. Brooks has taken three

specimens at Chilliwhack. Rare at Vernon. (*E. P. Venables.*)

GLAUCIDIUM Boie.

186. Pygmy Owl.

Glaucidium gnoma (Wagl.).

Common throughout the Province. (*Fannin.*)

187. California Pygmy Owl.

Glaucidium gnoma californicum (Scl.).

Common throughout the southern portions of the Province, including Vancouver Island.

Order COCCYGES. Cuckoos and Kingfishers.

Family CUCULIDÆ. Cuckoos.

COCCYZUS Vieillot.

188. Western Yellow-billed Cuckoo. California Cuckoo.

Coccyzus americanus occidentalis (Ridgw.).

A regular summer resident in the southwestern portions of the Province. It breeds on Vancouver Island near Victoria. Mr. Brooks has taken it at Chilliwhack.

Family ALCEDINIDÆ. Kingfishers.

CERYLE Boie.

189. Belted Kingfisher.

Ceryle alcyon (LINN.).

An abundant resident throughout the Province, breeds throughout its range.

Order PICI. Woodpeckers, Wrynecks, etc.

Family PICIDÆ. Woodpeckers.

DRYOBATES Boie.

190. Northern Hairy Woodpecker.

Dryobates villosus leucomelas (Bodd.).

Common resident on the Mainland east of Cascade Mountains. I found it quite common at Sicamous, May, 1895. Mr. Brooks reports having taken it several times at Chilliwhack.

191. Harris's Woodpecker.

Dryobates villosus harrisii (Aud.).

A common resident west of Coast Range, including Vancouver Island.

192. Cabanis's Woodpecker.

Dryobates villosus hyloscopus (Cab.).

Rocky Mountain district. Taken at Okanagan by A. C. Brooks.

193. Queen Charlotte Woodpecker.

Dryobates villosus picoideus (Osgood).

Queen Charlotte Islands.

194. Gairdner's Woodpecker.

Dryobates pubescens gairdneri (Aud.).

A very common resident on Vancouver Island, Coast of Mainland and Lower Fraser Valley.

195. Batchelder's Woodpecker.

Dryobates pubescens homorus (Cab.).

Common throughout the interior of the Province, along the C. P. R., at Ducks, and at Vernon. Mr. Brooks has taken it at Chilliwhack and Okanagan.

XENOPICUS Baird.

196. White-headed Woodpecker.

Xenopicus albolarvatus (Cass.).

A rare bird in this Province, found east of the Coast Range. Museum specimen from Similkameen Valley, collected by R. V. Griffin.

PICOIDES Lacépède.

197. Black-backed Three-toed Woodpecker.

Picoides arcticus (Swains.).

Found only east of Coast Range, where in some localities it is quite common. Mr. Brooks reports it from Okanagan.

198. Alaskan Three-toed Woodpecker.

Picoides americanus fasciatus (Bird).

West of Cascade Mountains common. Rare on Vancouver Island. Brooks says common resident in the spruce zone on the Coast Range.

199. Alpine Three-toed Woodpecker.

Picoides americanus dorsalis (Baird).

Mountains east of Cascade Mountains, north to Cassiar. (*Fannin.*)

SPHYRAPICUS Baird.

200. Red-naped Sapsucker.

Sphyrapicus varius nuchalis (Baird).

Generally distributed east of Cascade Mountains throughout the Province. It was quite common at Duck's, May, 1895.

201. Red-breasted Sapsucker.

Sphyrapicus ruber notkensis (Suckow.).

Generally distributed along the Coast. Not uncommon on Vancouver Island. Mr. Brooks has taken it at Chilliwhack.

202. Williamson's Sapsucker.

Sphyrapicus thyroideus (Cass.).

Taken at Similkameen, B. C., by R. V. Griffin, June, 1882, (Check List of B. C. Birds, 1891—*Fannin*.)

CEOPHLOEUS Cabanis.

203. Northern Pileated Woodpecker.

Ceophloeus pileatus abieticola (Bangs.).

Common resident throughout the Province, on Vancouver Island and Mainland through the interior to Okanagan. Mr. Brooks found it scarce in the Cariboo District.

ASYNDESMUS (Wils.) Coues.

204. Lewis's Woodpecker.

Asyndesmus torquatus (Wils.).

Abundant through the interior of the Province, at Vernon and Grand Prairie. I found it quite common at Chilliwack this year. A summer resident on Vancouver Island.

COLAPTES Swainson.

205. Northern Flicker.

Colaptes auratus luteus (Bangs.).

A rare bird of this Province, but is found on Vancouver Island and east to the Rocky Mountains.

206. Red-shafted Flicker.

Colaptes cafer collaris (Vigors).

Common east of Cascade Mountains through the interior, at Duck's, Sicamous and Okanagan.

207. North-western Flicker.

Colaptes cafer saturation (Ridgew.).

Abundant resident west of Cascade Mountains; very common on Vancouver Island.

Order MACROCHIRES. Goatsuckers, Swifts, etc.

Family CAPRIMULGIDÆ. Goatsuckers.

PHALÆNOPTILUS Ridgway.

208. Poor-will.

Phalænoptilis nuttallii (Aud.).

Southern interior portions of Province from Kamloops through the Okanagan. Summer resident.

CHORDEILES Swainson.

209. Nighthawk.

Chordeiles virginianus (Gmel.).

An abundant summer resident east of Cascade Mountains. Mr. Brooks has taken it at Chilliwhack.

210. Western Nighthawk.

Chordeiles virginianus henryi (Cass.).

A summer resident west of Cascade Mountains. Abundant on Vancouver Island.

Family MICROPODIDÆ. Swifts.

CYPSELOIDES Streubel.

211. Black Swift.

Cypseloides niger borealis (Ridgw.).

Common near Victoria during migration. Rev. J. H. Keen says only pass over at Metlakatla. Said to breed in mountains back of Chilliwhack and in the neighbourhood of Comox. (*Fannin.*) Taken at Chilliwhack and Okanagan. (*Brooks.*)

CHÆTURA Stephens.

212. Vaux's Swift.

Chætura vauxii (Townsend.).

A summer resident on Vancouver Island and Mainland. I found it very common near Sicamous, May 28th, 1895.

Family TROCHILIDÆ. Hummingbirds.

TROCHILUS Linnæus.

213. Black-chinned Hummingbird.

Trochilus alexandri (Bourc. & Muls.).

Confined to the Mainland. I have never seen it on Vancouver Island. Mr. Brooks has taken it at Chilliwhack and Okanagan.

SELASPHORUS Swainson.

214. Red-backed Rufous Hummingbird.

Selasphorus rufus (Gmel.).

An abundant summer resident west of Cascade Mountains, including Vancouver Island. Mr. Brooks has taken it at Okanagan. Common at Metlakatla (*Rev. J. H. Keen.*)

215. Green-backed Rufous Hummingbird. Allen's Hummingbird.

Selasphorus alleni (Hensh.).

Eastern Cascade and Rocky Mountain districts. (*Fannin.*)

STELLULA Gould.

216. Calliope Hummingbird.

Stellula calliope (Gould).

East and west of Cascade Mountains. (*Fannin.*)

Order PASSERES. Perching Birds.

Family TYRANNIDÆ. Tyrant Flycatchers.

TYRANNUS Cuvier.

217. Kingbird.

Tyrannus tyrannus (Linn.).

A common summer resident on the Mainland at Chilliwack, Sardis, Ducks and Vernon, and has been taken as far north as Port Simpson. Rare on Vancouver Island.

218. Gray Kingbird.

Tyrannus dominicensis (Gmel.).

Accidentally in B. C. One specimen taken at Cape Beale, October 9, 1889, by Miss Cox, and presented to the Museum.

219. Arkansas Kingbird.

Tyrannus verticalis (Say).

A common summer resident on the Mainland. I have found it quite common at Chilliwack, Ducks and Okanagan, rarely west to Vancouver Island.

SAYORNIS Bonaparte.

220. Say's Phoebe.

Sayornis saya (Bonap.).

A summer resident. Not common on Vancouver Island, chiefly on Mainland in the interior. Mr. Brooks has taken it at Chilliwhack and Okanagan.

CONTOPUS Cabanis.

221. Olive-sided Flycatcher.

Contopus borealis (Swains.).

A summer resident on Vancouver Island and Mainland. Mr. Brooks has taken it at Chilliwhack and Okanagan.

222. Western Wood Pewee.

Contopus richardsonii (Swains.).

A common summer resident on Vancouver Island and Mainland. Very common at Sicamous, May, 1895.

EMPIDONAX Cabanis.

223. Western Flycatcher.

Empidonax difficilis (Baird).

A common summer resident west of Cascade Mountains on Mainland and Vancouver Island. I also found it quite common at Skidegate, Queen Charlotte Islands, July, 1895.

224. Traill's Flycatcher.

Empidonax trailli (Aud.).

A summer resident. I have found it quite common on Vancouver Island and the Mainland at Sicamous and Vernon. Mr. Brooks has taken it at Chilliwhack and Okanagan.

225. Hammond's Flycatcher.

Empidonax hammondi (Xantus.).

Summer resident. Chiefly on the Mainland, east and west of Cascade Mountains. A few have been taken on Vancouver Island.

226. Wright's Flycatcher.

Empidonax wrightii (Baird).

Summer resident. Taken at Chilliwhack and Okanagan. (*Brooks.*)

Family ALAUDIDÆ. Larks.

OTOCORIS Bonaparte.

227. Pallid Horned Lark.

Otocoris alpestris leucoloema (Coues.).

Common in spring and autumn on Vancouver Island. Have taken it at Clover Point near Victoria. Mr. Brooks has taken it at Chilliwhack and Okanagan.

228. Streaked Horned Lark.

Otocoris alpestris strigata (Hensh.).

Spring and autumn migrant on Vancouver Island. Have taken it near Victoria in September. Mr. Brooks has taken it at Chilliwhack. West of Cascades; taken at Burrard Inlet and Vancouver Island (*Fannin*). Fort Simpson by W. B. Anderson.

229. Dusky Horned Lark.

Otocoris alpestris merrilli (Dwight).

Chiefly east of Cascades. I have never known it to be taken on Vancouver Island. Mr. Brooks has taken it at Chilliwhack and Okanagan. Chas. de B. Green has taken it at Osoyoos.

Family CORVIDÆ. Crows, Jays, Magpies.

PICA Brisson.

230. American Magpie.

Pica pica hudsonica (Sab.).

A common resident on the Mainland, rarely west to Vancouver Island. Breeds in the interior of the Mainland, east of Cascade Mountains.

CYANOCITTA Strickland.

231. Steller's Jay.

Cyanocitta stelleri (Gmel.).

An abundant resident west of Coast Range on Mainland and Vancouver Island throughout the year.

232. Black-headed Jay.

Cyanocitta stelleri annectens (Baird).

This form is quite common in the interior from the Cascades east down through the Okanagan. Mr. Brooks has taken it at Chilliwhack.

233. Queen Charlotte Jay.

Cyanocitta stelleri carlottæ (Osgood).

Queen Charlotte Islands; not uncommon at Skidegate, August, 1895.

PERISOREUS Bonaparte.

234. Rocky Mountain Jay.

Perisoreus canadensis capitalis (Ridgw.).

East of Cascade Mountains. Taken at Okanagan and in the Caribou District by A. C. Brooks.

235. Oregon Jay.

Perisoreus obscurus (Ridgw.).

An abundant resident west of Cascades, on the Mainland, but not so numerous on Vancouver Island.

236. Gray Jay.

Perisoreus obscurus griseus (Ridgw.).

California to British Columbia, east of the Coast and Cascade Ranges. (Ridgway.)

CORVUS Linnæus.

237. Northern Raven.

Corvus corax principalis (Ridgw.).

A resident throughout the Province; more common on the Coast. I have seen it quite common at Clayoquot, V. I., and in August, 1895, it was very common at Skidegate, Queen Charlotte Islands. Mr. Brooks has taken it at Chilliwhack and Okanagan. Common at Metlakatla. (*Rev. J. H. Keen.*)

238. American Crow.

Corvus brachyrhynchos (Brehm.)

Common east of Coast Range; common at Sicamous, May, 1895. Mr. Brooks has taken it at Chilliwhack and Okanagan.

239. Northwest Crow.

Corvus caurinus (Baird).

An abundant resident west of Cascade Mountains, on Mainland and Vancouver Island, and fairly common at Skidegate, Queen Charlotte Islands, July, 1895.

NUCIFRAGA Brisson.

240. Clarke's Nutcracker.

Nucifraga columbiana (Wils.).

A common resident on the Mainland, chiefly east of the Coast Range. Rare on Vancouver Island.

CYANOCEPHALUS Bonaparte.

241. Blue Crow. Piñon Jay.

Cyanocephalus cyanocephalus (Wied.).

Rocky Mountain region to the Pacific Coast Ranges, but rather more southerly; north to British Columbia; south to Lower California.—"Key to N. A. Birds (Coues), Fifth Edition."

Family ICTERIDÆ. Blackbirds, Orioles, Etc.

DOLICHONYX Swainson.

242. Bobolink.

Dolichonyx oryzivorus (Linn.)

East and west of Cascade Mountains. Taken at Chilliwhack by A. C. Brooks.

MOLOTHRUS Swainson.

243. Cowbird.

Molothrus ater (Bodd.).

A summer resident. I have found it fairly common in the neighbourhood of Ducks, on C. P. R., and Okanagan. It is a rare straggler on Vancouver Island; it has been taken near Victoria. Mr. Brooks has taken it at Chilliwhack. Rev. J. H. Keen reports it rare at Metlakatla.

XANTHOCEPHALUS Bonaparte.

244. Yellow-headed Blackbird.

Xanthocephalus xanthocephalus (Bonap.).

A rare summer resident on the Mainland. I have heard of it at Vernon. Fannin has found it above Clinton, on the Cariboo Road. Brooks has taken it at Chilliwhack.

AGELAIUS Vieillot.

245. San Diego Redwing.

Agelaius phoeniceus neutralis (Ridgw.).

Breeding range north, to eastern British Columbia. (*Ridgway.*) East of Cascades I found this form quite common near Vernon, June, 1895.

246. Northwestern Red-wing.

Agelaius phoeniceus caurinus (Ridgw.).

Common west of Cascades, on Mainland and Vancouver Island; it is found throughout the year near Victoria.

STURNELLA Vieillot.

247. Western Meadowlark.

Sturnella magna neglecta (Aud.).

Abundant resident on Mainland and Vancouver Island, east and west of Cascades.

ICTERUS Brisson.

248. Bullock's Oriole.

Icterus bullocki (Swains.).

A fairly common summer resident, chiefly east of Cascades. I found it breeding along the valley of the Thompson and in the Okanagan, near Vernon, May and June, 1895. Mr. Brooks has found it breeding at Chilliwhack.

EUPHAGUS Cassin.

249. Rusty Blackbird.

Euphagus carolinus (Muller).

Rare. One specimen shot at Metlakatla by Rev. J. H. Keen, November, 26th, 1901, and presented by him to the Museum.

250. Brewer's Blackbird.

Euphagus cyanocephalus (Wagler.).

Common, and generally distributed on the Mainland; quite common in the Lower Fraser Valley, Grand Prairie, south side of Thompson River, down into the Okanagan country. Breeding in suitable localities. Not common on Vancouver Island; a few have been taken near Victoria.

Family FRINGILLIDÆ. Finches, Sparrows, Etc.

HESPERIPHONA Bonaparte.

251. Western Evening Grosbeak.

Hesperiphona vespertina montana (Ridgw.).

Chiefly on the Mainland. West in winter to Vancouver Island and Lower Fraser Valley. Taken at Chilliwhack and Okaiiagan by A. C. B.

PINICOLA Vieillot.

252. Alaskan Pine Grosbeak.

Pinicola enucleator alascensis (Ridgw.).

Mainland east and west of Cascade Mountains. Common in winter in the neighbourhood of Clinton. Mr. Brooks reports it from Chilliwhack and Okanagan. Rare at Metlakatla. (Rev. J. H. Keen.)

CARPODACUS Kaup.

253. California Purple Finch.

Carpodacus purpureus californicus (Baird).

An abundant summer resident west of Cascades. Quite a number winter on Vancouver Island.

254. Cassin's Purple Finch.

Carpodacus cassinii (Baird).

East of Cascades to Rocky Mountains. Tolerably common.

LOXIA Linnæus.

255. American Crossbill.

Loxia curvirostra minor (Brehm.).

An abundant resident throughout the Province, Island and Mainland.

256. White-winged Crossbill.

Loxia leucoptera (Gmel.).

An abundant resident chiefly on the Mainland, occasionally west to Vancouver Island.

LEUCOSTICTE Swainson.

257. Gray-crowned Leucosticte.

Leucosticte tephrocotis (Swains.).

Rocky Mountain district. In winter, occasionally west to Cascades. Mr. Brooks has taken it as far west as Chilliwhack.

258. Hepburn's Leucosticte.

Leucosticte tephrocotis littoralis (Baird).

Common on the Mainland from the Coast to Rocky Mountains. Breeds above timber line in the interior.

ACANTHIS Bechstein.

259. Hoary Redpoll.

Acanthis hornemannii exilipes (Coues.).

Mouth of Quesnelle, S. Williams. Chilliwhack, A. C. Brooks.

260. Redpoll.

Acanthis linaria (Linn.).

The Province at large, but more abundant in the Rocky Mountain district. West in winter to Vancouver Island.

ASTRAGALINUS Cabanis.

261. American Goldfinch.

Astragalinus tristis (Linn.).

Both slopes of Cascades to Rocky Mountains on the Mainland. Taken at Okanagan by A. C. Brooks.

262. Willow Goldfinch.

Astragalinus tristis salicamans (Grinnell).

Taken at Okanagan by A. C. Brooks.

SPINUS Koch.

263. Pine Siskin.

Spinus pinus (Wils.).

An abundant resident throughout the Province. Breeds throughout its range.

PASSER Brisson.

264. House Sparrow.

Passer domesticus (Linn.).

This bird has now become quite common in the cities along the Coast.

PASSERINA (Vieill).

265. Snowflake.

Passerina nivalis (Linn.).

Not common on Vancouver Island. More abundant east of Cascades on the Mainland.

CALCARIUS Bechstein.

266. Lapland Longspur.

Calcarius lapponicus (Linn.).

The Province at large, but nowhere common. It has been taken at Victoria, Chilliwhack, Burrard Inlet, Okanagan, Fort Simpson and Metlakatla.

RHYNCHOPHANES Baird.

267. McCown's Longspur.

Rhynchophanes mccownii (Lawr.).

Taken at Chilliwhack by A. C. Brooks.

POOECETES Baird.

268. Western Vesper Sparrow.

Pooecetes gramineus confinis (Baird).

Interior portions of the Mainland east of Cascades, and in the Okanagan.

269. Oregon Vesper Sparrow.

Pooecetes gramineus affinis (Miller).

West of Cascades, including Vancouver Island.

PASSERCULUS Bonaparte.

270. Sandwich Sparrow.

Passerculus sandwichensis (Gmel.).

An abundant summer resident west of Cascades on the Mainland, including Vancouver Island. Mr. Brooks has taken it at Okanagan.

271. Western Savanna Sparrow.

Passerculus sandwichensis alaudinus (Bonaparte).

A common summer resident on the Coast, including Vancouver Island.

COTURNICULUS Bonaparte.

272. Western Grasshopper Sparrow.

Coturniculus savannarum bimaculatus (Swainson).

A summer resident in the Okanagan, near Vernon. (*Brooks.*)

CHONDESTES Swainson.

273. Western Lark Sparrow.

Chondestes grammacus strigatus (Swains.).

Interior southern portions of the Mainland, from Chilliwack east through the Okanagan.

ZONOTRICHIA Swainson.

274. Harris's Sparrow.

Zonotrichia querula (Nutt.).

Cadbora Bay, near Victoria, October, 1894, A. H. Maynard; Comox, November, 1894, W. B. Anderson; Sumas (Lower Fraser), 10th January, 1895, A. C. Brooks.

275. Gambel's Sparrow.

Zonotrichia leucophrys gambelii (Nutt.).

Mainland, east of Coast Range, east to Okanagan. Common.

276. Nuttall's Sparrow.

Zonotrichia leucophrys nuttalli (Ridgw.).

An abundant summer resident on Vancouver Island and Mainland, west of Cascades.

277. Golden-crowned Sparrow.

Zonotrichia coronata (Pallas.).

Abundant during migrations on the Coast; have also taken it at Chilliwack and Okanagan.

SPIZELLA Bonaparte.

278. Western Tree Sparrow.

Spizella monticola ochracea (Brewst.).

Chilliwack and Okanagan, A. C. Brooks; Victoria, A. H. Maynard.

279. Western Chipping Sparrow.

Spizella socialis arizonæ (Coues.).

An abundant summer resident east and west of Cascades. Breeds in the neighbourhood of this City. Also a number remain throughout the winter.

280. Clay-coloured Sparrow.

Spizella pallida (Swains.).

Carpenter's Mountain, 150-Mile House, Cariboo, July 3rd, 1901.
(Brooks.)

281. Brewer's Sparrow.

Spizella breweri (Cass.).

Eastern Cascades and Rocky Mountains districts. Similkameen, R. V. Griffin; Okanagan, A. C. Brooks.

JUNCO Wagler.

282. Slate-coloured Junco.

Junco hyemalis (Linn.).

Taken at Chilliwhack by A. C. Brooks.

283. Oregon Junco.

Junco hyemalis oregonus (Townsend).

Abundant resident west of Cascades, including Vancouver Island.

284. Shufeldt's Junco.

Junco hyemalis shufeldti (Coale).

Breeding from interior British Columbia (Stuarts Lake) east to Rocky Mountains, south to Vancouver Island.

MELOSIPZA Baird.

285. Rusty Song Sparrow.

Melosipza cinerea morphna (Oberh.).

An abundant resident throughout the Province. Breeds on Vancouver Island and Mainland.

286. Sooty Song Sparrow.

Melosipza cinerea rufina (Bonap.).

An abundant resident, chiefly along the coast of Mainland and Vancouver Island. I found it quite common at Skidegate, Queen Charlotte Islands, August, 1895.

287. Lincoln's Sparrow.

Melosipza lincolnii (Aud.).

Not common. East and west of Cascades; a few have been taken near Victoria. Mr. Brooks has taken it at Okanagan.

288. Forbush's Sparrow.

Melosipza lincolnii striata (Brewst.).

Collected at Comox, September, 1888, by E. H. Forbush. Taken at Chilliwhack by A. C. Brooks.

PASSERELLA Swainson.

289. Townsend's Sparrow.

Passerella iliaca unalaschcensis (Gmel.).

A common summer resident west of Cascades. A few remain throughout the winter on Vancouver Island; I have taken it in November and January.

290. Slate-coloured Sparrow.

Passerella iliaca schistacea (Baird).

Southern portions of Mainland, east of Cascades. Sicamous, J. Fannin, Okanagan, A. C. Brooks.

PIPILO Vieillot.

291. Spurred Towhee.

Pipilo maculatus megalonyx (Baird).

East of Cascades, south through the Okanagan.

292. Oregon Towhee.

Pipilo maculatus oregonus (Bell.).

An abundant resident west of Cascades. Breeds on Vancouver Island and Mainland.

ZAMELODIA Coues.

293. Black-headed Grosbeak.

Zamelodia melanocephala (Swains).

A summer resident on Vancouver Island. Taken at Chilliwhack. (*Brooks.*)
A summer resident east and west of Cascades. (*Fannin.*)

CYANOSPIZA Baird.

294. Lazuli Bunting.

Cyanospiza amoena (Say.).

A summer resident chiefly east of Cascades. I have seen it at Chilliwhack and near Vernon. Rare on Vancouver Island.

Family TANAGRIDÆ. Tanagers.

PIRANGA Vieillot.

295. Louisiana Tanager.

Piranga ludoviciana (Wils.).

An abundant summer resident throughout the Province. Vancouver Island and Mainland.

Family HIRUNDINIDÆ. Swallows.

PROGNE Boie.

296. Purple Martin.

Progne subis (Linn.).

A common summer resident chiefly in the cities along the Coast. West of Cascades.

PETROCHELIDON Cabanis.

297. Cliff Swallow.

Petrochelidon lunifrons (Say.).

Common summer resident on the Mainland east and west of Cascades, Chilliwhack, Okanagan and Cariboo Road.

HIRUNDO Linn.

298. Barn Swallow.

Hirundo erythrogastra (Bodd).

An abundant summer resident throughout the Province, including Vancouver Island and Queen Charlotte Islands.

TACHYCINETA Cabanis.

299. White-bellied Swallow.

Tachycineta bicolor (Vieill.).

An abundant summer resident east and west of Cascades.

300. Northern Violet-green Swallow.

Tachycineta thalassina lepidus (Mearns).

An abundant summer resident throughout the Province, east and west of Cascades.

RIPARIA Forester.

301. Bank Swallow.

Riparia riparia (Linn.).

An abundant summer resident, chiefly east of Cascades, north to Cariboo District. Very common from Sicamous down through the Okanagan.

Taken at Chilliwhack by A. C. Brooks.

STELGIDOPTERYX Baird.

302. Rough-winged Swallow.

Stelgidopteryx serripennis (Aud.).

A common summer resident throughout the Province. Vancouver Island and Mainland.

Family AMPELIDÆ. Waxwings.

AMPELIS Linnæus.

303. Bohemian Waxwing.

Ampelis garrulus (Linn.).

An abundant resident chiefly east of Cascades. A winter visitant to the Coast and Vancouver Island.

304. Cedar Bird.

Ampelis cedrorum (Vieill.).

A common summer resident throughout the Province. Vancouver Island and Mainland.

Family LANIIDÆ. Shrikes.

LANIUS Linnæus.

305. Northern Shrike. Butcher-bird.

Lanius borealis (Vieill.).

Not common, but found distributed throughout the Province. Found throughout the winter on the Coast.

306. White-rumped Shrike.

Lanis ludovicianus excubitorides (Swains.).

Chilliwhack, A. C. Brooks. Victoria, A. H. Maynard.

Family VIREONIDÆ. Vireos.

VIREO Vieillot.

307. Red-eyed Vireo.

Vireo olivaceus (Linn.).

Distributed over the southern portions of the Province, including Vancouver Island. A summer resident. Chilliwhack and Okanagan, A. C. Brooks.

308. Western Warbling Vireo.

Vireo gilvus swainsonii (Baird).

Common summer resident, west of Rocky Mountains to Pacific. Mainland and Vancouver Island.

309. Cassin's Vireo.

Vireo solitarius cassinii (Xantus.).

A summer resident east and west of Cascades, including Vancouver Island.

310. Anthony's Vireo.

Vireo huttoni obscurus (Anthony).

A summer resident on Vancouver Island.

Family MNIOTILTIDÆ. Wood Warblers.

HELMINTHOPHILA Ridgway.

311. Calaveras Warbler.

Helminthophila ruficapilla gutturalis (Ridgw.).

Rocky Mountains to Pacific. A summer resident. Common in the Okanagan. (*Brooks.*)

312. Orange-crowned Warbler.

Helminthophila celata (Say.).

A common summer resident east and west of Cascades.

313. Lutescent Warbler.

Helminthophila celata lutescens (Ridgw.).

An abundant summer resident, chiefly west of Cascades, along the coast of Mainland and Vancouver Island.

314. Tennessee Warbler.

Helminthophila peregrina (Wils.).

Taken at 150-Mile House, Cariboo, by A. C. Brooks.

DENDROICA Gray.

315. Alaskan Yellow Warbler.

Dendroica æstiva rubiginosa (Pallas).

An abundant summer resident throughout the Province, from Rocky Mountains to the Pacific.

316. Myrtle Warbler.

Dendroica coronata (Linn.).

An abundant summer resident on Vancouver Island and Mainland, chiefly west of Cascades. Rare at Metlakatla. (Rev. J. H. Keen.) Okanagan. (Brooks).

317. Audubon's Warbler.

Dendroica auduboni (Townsend.).

An abundant summer resident throughout the Province, Vancouver Island and Mainland east to Rocky Mountains.

318. Magnolia Warbler.

Dendroica maculosa (Gmel.).

Okanagan, 1898 (Rhoads).

319. Black-throated Gray Warbler.

Dendroica nigrescens (Townsend.).

Not common. A summer resident west of Cascades, but chiefly along the Coast. Taken at Chilliwhack (Brooks).

320. Townsend's Warbler.

Dendroica townsendi (Townsend.).

Not common, but found distributed throughout the Province, chiefly west of Cascades, including Vancouver Island and Queen Charlotte Islands.

321. Hermit Warbler. Western Warbler.

Dendroica occidentalis (Townsend).

A summer resident, chiefly west of Cascade Range.

SEIURUS Swainson.

322. Grinnell's Water-thrush.

Seiurus noveboracensis notabilis (Ridgw.).

Taken at Ducks, C. P. R., by Clark P. Streater, August 7th, 1889.

Taken at Quesnel, August 30th, 1900, by A. C. Brooks.

GEOTHTYPIS Cabanis.

323. Macgillivray's Warbler.

Geothlypis tolmiei (Townsend).

A common summer resident throughout the Province, Vancouver Island and Mainland.

324. Western Yellow-throat.

Geothlypis trichas occidentalis (Brewst.).

A common summer resident throughout the Province.

325. Pacific Yellow-throat.

Geothlypis trichas arizela (Oberh.)

Summer resident in the southern portion of Mainland and Vancouver Island.

ICTERIA Vieillot.

326. Long-tailed Chat.

Icteria virens longicauda (Lawr.).

Summer resident in the southern portions of the Province, from Sumas east to Okanagan, south of Ashcroft.

WILSONIA Bonaparte.

327. Wilson's Warbler.

Wilsonia pusilla (Wils.).

Vancouver Island and Mainland, east and west of Cascades. S. N. Rhoads, 1892 (*Fannin*.)

328. Pileolated Warbler.

Wilsonia pusilla pileolata (Pall.).

A common summer resident on Vancouver Island and Mainland.

329. Golden Pileolated Warbler.

Wilsonia pusilla chryseola (Ridgw.).

A summer resident, New Westminster and Mount Lehman.

SETOPHAGA Swainson.

330. American Redstart.

Setophaga ruticilla (Linn.).

Found through the interior of the Province, from Barkerville south to Okanagan, chiefly east of, and accidentally west of, Cascades. Taken at Chilliwhack by A. C. Brooks.

Family MOTACILLIDÆ. Wagtails and Pipits.

ANTHUS Bechstein.

331. American Pipit.

Anthus pensilvanicus (Lath.).

Very abundant during migration, east and west of Cascades. Found throughout some winters on Vancouver Island.

Family CINCLIDÆ. Dippers.

CINCLUS Bechstein.

332. American Dipper.

Cinclus mexicanus (Swains.).

Common throughout the Province in suitable localities.

Family TROGLODYTIDÆ. Wrens.

GALEOSOPTES Cabanis.

333. Catbird.

Galeoscoptes carolinensis (Linn.).

A common summer resident east and west of Cascades. Rare on Vancouver Island.

SALPINCTES Cabanis.

334. Rock Wren.

Salpinctes obsoletus (Say.).

Common east of Cascades, rarely west to the Coast. Chilliwack. (*Brooks.*) Burrard Inlet. (*Fannin.*)

THRYOMANES Sclater.

335. Vigor's Wren.

Thryomanes bewickii spilurus (Vig.).

A common summer resident west of Cascades. Found throughout the winter on Vancouver Island.

TROGLODYTES Vieillot.

336. Parkman's Wren.

Troglodytes aëdon parkmanii (Aud.).

A common summer resident east and west of Cascades, including Vancouver Island.

OLBIORCHILUS Oberholser.

337. Western Winter Wren.

Olbiorchilus hiemalis pacificus (Baird).

An abundant resident east and west of Cascades, including Vancouver Island.

TELMATODYTES Cabanis.

338. Tule Wren.

Telmatodytes palustris paludicola (Baird).

Abundant east of Cascades, from the lakes along the Cariboo Road, south through the Okanagan. Taken at Chilliwhack by A. C. Brooks.

Family CERTHIIDÆ. Creepers.

CERTHIA Linnæus.

339. Rocky Mountain Creeper.

Certhia familiaris montana (Ridgw.).

Southern portion of east of Cascades. Okanagan.

340. California Creeper.

Certhia familiaris occidentalis (Ridgw.).

Common on Vancouver Island. East and west of Cascades on the Mainland.

Family SITTIDÆ. Nuthatches.

SITTA Linnæus.

341. Slender-billed Nuthatch.

Sitta carolinensis aculeata (Cass.).

Common east of Cascades, from Ashcroft south through the Okanagan. I found it quite common near Shuswap, May, 1895.

342. Red-breasted Nuthatch.

Sitta canadensis (Linn.).

Common east and west of Cascades. Found throughout the winter on Vancouver Island.

343. Pygmy Nuthatch.

Sitta pygmæa (Vig.).

East of Cascades, in the south-eastern portions of the Province. I found it fairly common at Grand Prairie, May, 1895.

Family PARIDÆ. Titmice or Chickadees.

PARUS Linnæus.

344. Long-tailed Chickadee.

Parus atricapillus septentrionalis (Harris.).

Common east of Cascades, from Cornwall's south to Okanagan.

345. Oregon Chickadee.

Parus atricapillus occidentalis (Baird.).

A common resident west of Cascades.

346. Mountain Chickadee.

Parus gambeli (Ridgw.).

East of Cascades to Rocky Mountains, and south through Okanagan.

347. Columbia Chickadee.

Parus hudsonicus columbianus (Rhoads).

East of Cascades to Rocky Mountains, from Cassiar District south to Okanagan.

348. Chestnut-backed Chickadee.

Parus rufescens (Townsend.).

An abundant resident chiefly west of Cascades. Vancouver Island and Queen Charlotte Islands. Taken at Arrow Lake by A. C. Brooks.

Family CHAMÆIDÆ. Wren, Tits and Bush-Tits.

PSALTRIPARUS Bonaparte.

349. Least Bush-tit.

Psaltiriparus minimus (Towns.).

Rare. Taken at Sumas, November 25, 1899, by A. C. Brooks.

Family SYLVIIDÆ. Warblers. Kinglets.

REGULUS Cuvier.

350. Western Golden-crowned Kinglet.

Regulus satrapa olivaceus (Baird).

Abundant throughout the Province.

351. Ruby-crowned Kinglet.

Regulus calendula (Linn.).

Abundant throughout the Province.

Family TURDIDÆ. Thrushes, Solitaires and Bluebirds.

MYADESTES Swainson.

352. Townsend's Solitaire.

Myadestes townsendii (Aud.).

Not common. Found east and west of Cascades, including Vancouver Island.

HYLOCICHLA Baird.

353. Willow Thrush.

Hylocichla fuscescens salicicola (Ridgw.).

A common summer resident east of Cascades.

354. Russet-backed Thrush.

Hylocichla ustulata (Nutt.).

A common summer resident west of Cascades. Very common on Queen Charlotte Islands, July, 1895.

355. Olive-backed Thrush.

Hylocichla ustulata swainsoni (Cab.).

Southern portion of Mainland east of Cascades.

356. Alaska Hermit Thrush.

Hylocichla guttata (Pallas).

West of Cascades, chiefly coastwise. Common at Skidegate, Queen Charlotte Islands, August, 1895.

357. Audubon's Hermit Thrush.

Hylocichla guttata auduboni (Baird).

Southern portions of the Province, east of Cascades to Rocky Mountains.

MERULA Leach.

358. American Robin.

Merula migratoria (Linn.).

Both forms are to be found on Vancouver Island. Taken at Chilliwhack by A. C. Brooks.

359. Western Robin.

Merula migratoria propinqua (Ridgw.).

An abundant resident throughout the Province.

IXOREUS Bonaparte.

360. Varied Thrush.

Ixoreus naevius (Gmel.).

Common throughout the Province, east and west of Cascades.

SIALIA Swainson.

361. Western Bluebird.

Sialia mexicana occidentalis (Townsend.).

A common summer resident east and west of Cascades. A few winter on Vancouver Island.

362. Mountain Bluebird.

Sialia arctica (Swains.).

A summer resident east and west of Cascades, including Vancouver Island.

INDEX.

A.

No.

Albatross, Black-footed 35

Albatross, Short-tailed 36

Auklet, Rhinoceros 12

Auklet, Cassin's 13

B.

Bittern, American 88

Blackbird, Yellow-headed 244

Blackbird, San Diego Redwing 245

Blackbird, North-western 246

Blackbird, Rusty 249

Blackbird, Brewer's 250

Bluebird, Western 361

Bluebird, Mountain 362

Bobolink 242

Brant 82

Brant, Black 83

Bunting, Lazuli 294

Bush-tit, Least 349

C.

Catbird 333
Chat, Long-tailed 326
Chickadee, Long-tailed 344
Chickadee, Oregon 345
Chickadee, Mountain 346
Chickadee, Columbia 347
Chickadee, Chestnut-backed 348
Cormorant, White-crested 43
Cormorant, Brandt's 44
Cormorant, Violet-green 45
Cowbird 243
Coot, American 96
Crane, Little Brown 92
Crane, Sandhill 93
Creeper, Rocky Mountain 339
Creeper, California 340
Crow, American 238
Crow, Northwest 239
Crossbill, American 255
Crossbill, White-winged 256
Cuckoo, Western Yellow-billed 188
Curlew, Long-billed 122
Curlew, Hudsonian 123

D.

Dipper, American 332
Dove, Mourning 148
Dowitcher, Long-billed 101

Duck, Mallard 51
Duck, Gadwall 52
Duck, European Widgeon 53
Duck, American Widgeon 54
Duck, Green-winged Teal 55
Duck, Blue-winged Teal 56
Duck, Cinnamon Teal 57
Duck, Shoveller 58
Duck, Pintail 59
Duck, Wood 60
Duck, Red-head 61
Duck, Canvas-back 62
Duck, American Scaup 63
Duck, Lesser Scaup 64
Duck, Ring-necked 65
Duck, American Golden-eye 66
Duck, Barrow's Golden-rye 67
Duck, Buffle-head 68
Duck, Long-tailed 69
Duck, Harlequin 70
Duck, American Black Scoter 71
Duck, White-winged Scoter 72
Duck, Surf Scoter 73
Duck, Ruddy 74

E.

Eagle, Golden 160
Eagle, Bald 161

F.

Falcon, Prairie 163
Falcon, Peale's 165
Finch, California Purple 253
Finch, Cassin's Purple 254
Flycatcher, Olive-sided 221
Flycatcher, Western 223
Flycatcher, Traill's 224
Flycatcher, Hammond's 225
Flycatcher, Wright's 226
Flicker, Northern 205
Flicker, Red-shafted 206
Flicker, North-western 207
Fulmar, Pacific 37

G.

Godwit, Marbled 112
Goose, Lesser Snow 75
Goose, Ross's Snow 76
Goose, American White-fronted 77
Goose, Canada 78
Goose, Hutchin's 79
Goose, White-cheeked 80
Goose, Cackling 81
Goose, Emperor 84
Goldfinch, American 261
Goldfinch, Willow 262
Grebe, Western 1
Grebe, American Red-necked 2
Grebe, Horned 3
Grebe, American Eared 4

Grebe, Pied-billed 5
Grouse, Sooty 134
Grouse, Richardson's 135
Grouse, Franklin's 136
Grouse, Canadian Ruffed 137
Grouse, Gray Ruffed 138
Grouse, Oregon Ruffed 139
Grouse, Columbian Sharp-tailed 143
Grouse, Sage 144
Grosbeak, Western Evening 251
Grosbeak, Alaskan Pine 252
Grosbeak, Black-headed 293
Guillemot, Pigeon 16
Gull, Ivory 21
Gull, Glaucous-winged 23
Gull, Western 24
Gull, American 25
Gull, California 26
Gull, Ring-billed 27
Gull, Short-billed 28
Gull, Heermann's 29
Gull, Bonaparte's 30
Gull, Sabine's 31
Gyr Falcon 162

H.

Hawk, Marsh 151
Hawk, Sharp-shinned 152
Hawk, Cooper's 153
Hawk, American Goshawk 154

Hawk, Western Goshawk 155
Hawk, Western Red-tailed 156
Hawk, Red-bellied 157
Hawk, Swainson's 158
Hawk, American Rough-legged 159
Hawk, Duck 164
Hawk, Pigeon 166
Hawk, Desert Sparrow 169
Heron, Great Blue 89
Heron, North-west Coast 90
Heron, Snowy 91
Hummingbird, Black-chinned 213
Hummingbird, Red-backed Rufous 214
Hummingbird, Green-backed Rufous 215
Hummingbird, Calliope 216

I.

Ibis, White-faced Glossy 87

J.

Jaeger, Parasitic 19
Jaeger, Pomarine 18
Jaeger, Long-tailed 20
Jay, Steller's 231
Jay, Black-headed 232
Jay, Queen Charlotte 233
Jay, Rocky Mountain 234
Jay, Oregon 235
Jay, Gray 236
Jay, Piñon 241

Junco, Slate-colored 282

Junco, Oregon 283

Junco, Shufeldt's 284

K.

Kittiwake, Pacific 22

Kingfisher, Belted 189

Kingbird 217

Kingbird, Gray 218

Kingbird, Arkansas 219

Kinglet, Western Golden-crowned 350

Kinglet, Ruby-crowned 351

Knot 103

L.

Lark, Pallid Horned 227

Lark, Streaked Horned 228

Lark, Dusky Horned 229

Lencosticte, Gray-crowned 257

Lencosticte, Hepburn's 258

Longspur, Lapland 266

Longspur, McCown's 267

Loon, Great Northern Diver 6

Loon, Black-throated 7

Loon, Pacific 8

Loon, Red-throated 9

M.

Martin, Purple 296
Magpie, American 230
Meadowlark, Western 247
Merlin, Black 167
Merlin, Richardson's 168
Merganser, American 48
Merganser, Red-breasted 49
Merganser, Hooded 50
Murrelet, Ancient 14
Murrelet, Marbled 15
Murre, California 17

N.

Nighthawk 207
Nighthawk, Western 208
Nutcracker, Clarke's 240
Nuthatch, Slender-billed 341
Nuthatch, Red-breasted 342
Nuthatch, Pygmy 343

O.

Oriele, Bullock's 248
Osprey, American 170
Owl, American Long-eared 171
Owl, Short-eared 172
Owl, Northern Spotted 173
Owl, Great Gray 174
Owl, Richardson's 175
Owl, Saw-whet 176
Owl, North-West Saw-whet 176A

Owl, Kennicott's Screech 177
Owl, Macfarlane's Screech 178
Owl, Great Horned 179
Owl, Western Horned 180
Owl, Arctic Horned 181
Owl, Dusky Horned 182
Owl, Snowy 183
Owl, American Hawk 184
Owl, Burrowing 185
Owl, Pygmy 186
Owl, California Pygmy 187
Oyster-catcher, Black 131

P.

Partridge, Mountain 132
Partridge, California 133
Pelican, American White 46
Pelican, California Brown 47
Petrel, Gray Fork-tailed 41
Petrel, Leach's 42
Pewee, Western Wood 222
Phalarope, Red 97
Phalarope, Northern 98
Phalarope, Wilson's 99
Pheasant, Ring-necked 145
Phoebe, Say's 220
Pigeon, Band-tailed 146
Pigeon, Passenger 147
Pipit, American 331
Plover, American Golden 125

Plover, Black-bellied 124
Plover, Killdeer 126
Plover, Semipalmated 127
Poor-Will 208
Ptarmigan, Willow 140
Ptarmigan, Rock 141
Ptarmigan, White-tailed 142
Puffin, Tufted 10
Puffin, Horned 11

R.

Rail, Virginia 94
Rail, Carolina 95
Raven, Northern 237
Redpoll, Hoary 259
Redpoll 260
Redstart, American 330
Robin, American 358
Robin, Western 359

S.

Sanderling 111
Sandpiper, Stilt 102
Sandpiper, Sharp-tailed 104
Sandpiper, Pectoral 105
Sandpiper, Baird's 106
Sandpiper, Least 107
Sandpiper, Red-backed 108
Sandpiper, Semipal mated 109
Sandpiper, Western Semipalmated 110

Sandpiper, Solitary 115
Sandpiper, Western Solitary 116
Sandpiper, Bartramian 119
Sandpiper, Buff-breasted 120
Sandpiper, Spotted 121
Sapsucker, Red-naped 200
Sapsucker, Red-breasted 201
Sapsucker, Williamson's 202
Shrike, Northern 305
Shrike, White-rumped 306
Shearwater, Black-vented 38
Shearwater, Dark-bodied 39
Shearwater, Slender-billed 40
Siskin, Pine 263
Snipe, Wilson's 100
Snowflake 265
Solitaire, Townsend's 352
Sparrow, House 264
Sparrow, Western Vesper 268
Sparrow, Oregon Vesper 269
Sparrow, Sandwich 270
Sparrow, Western Savanna 271
Sparrow, Western Grasshopper 272
Sparrow, Western Lark 273
Sparrow, Harris's 274
Sparrow, Gambel's 275
Sparrow, Nuttall's 276
Sparrow, Golden-crowned 277
Sparrow, Western Tree 278
Sparrow, Western Chipping 279

Sparrow, Clay-colored 280
Sparrow, Brewer's 281
Sparrow, Rusty Song 285
Sparrow, Sooty Song 286
Sparrow, Lincoln's 287
Sparrow, Forbush's 288
Sparrow, Townsend's 289
Sparrow, Slate-coloured 290
Surf-bird 128
Swan, Whistling 85
Swan, Trumpeter 86
Swallow, Cliff 297
Swallow, Barn 298
Swallow, White-bellied 299
Swallow, Northern violet-green 300
Swallow, Bank 301
Swallow, Rough-winged 302
Swift, Black 211
Swift, Vaux 212

T.

Tanager, Louisaria 295
Tatler, Wandering 118
Tern, Common 32
Tern, Arctic 33
Tern, Black 34
Thrush, Willow 353
Thrush, Russet-backed 354
Thrush, Olive-backed 355
Thrush, Alaska Hermit 356

Thrush, Audubon's 357

Thrush, Varied 360

Towhee, Spurred 291

Towhee, Oregon 292

Turnstone 129

Turnstone, Black 130

v.

Vireo, Red-eyed 307

Vireo, Western Warbling 308

Vireo, Cassin's 309

Vireo, Anthony's 310

Vulture, California 149

Vulture, Turkey 150

w.

Warbler, Calaver's 311

Warbler, Orange-crowned 312

Warbler, Lutescent 313

Warbler, Tennessee 314

Warbler, Alaskan Yellow 315

Warbler, Myrtle 316

Warbler, Audubon's 317

Warbler, Magnolia 318

Warbler, Black-throated Gray 319

Warbler, Townsend's 320

Warbler, Macgillivray's 323

Warbler, Hermit 321

Warbler, Western Yellow-throat 324

Warbler, Pacific Yellow-throat 325

Warbler, Wilson's 327
Warbler, Pileolated 328
Warbler, Golden Pileolated 329
Water-thrush, Grinnell's 322
Waxwing, Bohemian 303
Waxwing, Cedar 304
Willett, Western 117
Woodpecker, Northern Hairy 190
Woodpecker, Harris's 191
Woodpecker, Cabanis's 192
Woodpecker, Queen Charlotte 193
Woodpecker, Gairdner's 194
Woodpecker, Batchelder's 195
Woodpecker, White-headed 196
Woodpecker, Black-backed Three-toed 197
Woodpecker, Alaskan Three-toed 198
Woodpecker, Alpine Three-toed 199
Woodpecker, Northern Pileated 203
Woodpecker, Lewis's 204
Wren, Rock 334
Wren, Vigor's 335
Wren, Parkman's 336
Wren, Western Winter 337
Wren, Tule 338

Y.

Yellow-legs, Greater 113
Yellow-legs, Lesser 114

*** END OF THE PROJECT GUTENBERG EBOOK CATALOGUE OF
BRITISH COLUMBIA BIRDS ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project

Gutenberg Literary Archive Foundation, the manager of the Project Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund.

If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you ‘AS-IS’, WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.