


Wild Birds in City Parks


Herbert Eugene Walter and Alice Hall

An abstract geometric pattern composed of various shapes including circles, triangles, squares, and lines, all rendered in shades of purple and magenta. The pattern is dense and covers the entire background of the lower half of the image.

Project Gutenberg

Wild Birds in City Parks

Herbert Eugene Walter and Alice Hall

An abstract geometric pattern in shades of pink and magenta. It features various shapes including circles, arcs, straight lines, and triangles, some of which are nested or overlapping. The pattern is dense and fills the lower two-thirds of the page.

Project Gutenberg

The Project Gutenberg eBook of Wild Birds in City Parks

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: Wild Birds in City Parks

Author: Herbert Eugene Walter
Alice Hall Walter

Release date: July 9, 2010 [eBook #33125]

Language: English

Other information and formats: www.gutenberg.org/ebooks/33125

Credits: Produced by Bryan Ness, Stephen Hutcheson, and the Online Distributed Proofreading Team at <http://www.pgdp.net>

*** START OF THE PROJECT GUTENBERG EBOOK WILD BIRDS IN
CITY PARKS ***

Wild Birds in City Parks

Being hints on identifying 145 birds, prepared primarily for the spring migration in Lincoln Park, Chicago

BY
HERBERT EUGENE WALTER
AND
ALICE HALL WALTER
REVISED AND ENLARGED EDITION
WITH CHART AND KEY

CHICAGO
A. W. MUMFORD & COMPANY, PUBLISHERS
378 Wabash Avenue
1906

[2

Copyright 1901, 1902 and 1904
by Alice Hall Walter

[3

Note to Third Edition.

With the continued success of this little booklet the authors have been led to revise and enlarge the present issue in the hope that it will prove more helpful and complete to those beginning the outdoor study of birds.

The new features of this edition are the addition of forty-five birds—the majority of which are shore and water birds—and a simple field key. The number of each bird as given in the check-list of the American Ornithologist's Union has also been inserted after its scientific name as an aid in referring to larger bird-books. The subject matter has been carefully rewritten and the order of the birds rearranged in accordance with data covering observations during the last seven years, as well as the charts which show the relative abundance of the birds and the height of the migration in Lincoln Park.

We wish to thank our friends for their kind support in furthering our efforts to enlarge the circle of admirers which the wild birds so richly deserve.

[4

“All nature is so full that that district produces the greatest variety which is most examined.”

—Gilbert White, 1768.

15

Preface.

The object of this little book is to furnish those who may be interested in making the acquaintance of wild birds with a simple letter of introduction to 145 birds, the majority of which are commonly seen during the spring migration.

Complete descriptions have been avoided, in the belief that the student should rely upon his own observation for the discovery of minor details. *The living bird* is the one important fact which will make the brief hints offered of value.

Anyone caring to make use of these hints may be assured that during the migrations of the birds city dwellers have one of the keenest delights of country life brought to their very doors, because many birds, migrating largely at night, are attracted by the lights of the city and stop off in their long journey to feed, so that a city park often contains a greater variety of feathered visitors than an equal area in the country.

We wish to remind those of our friends who have asked for pictures in a future edition that this book actually is *copiously illustrated* by hundreds of living birds every springtime in our parks and around our homes, illustrations that are all life size, absolutely accurate in detail and colored true to nature.

“As for the birds * * * they add immeasurably to the wholesome beauty of life.” 16

—Theodore Roosevelt, in *Bird Lore*, Vol. II, p. 98.

17

General Hints.

“A good observer is quick to take a hint and follow it up.”—*John Burroughs*.

The identification of birds depends quite as much upon accurate observation of their size, motions, flight, characteristic attitudes, manner of feeding, company, song, call-notes and haunts, as upon details of form and color. Especial care is necessary to insure correct estimates of size for the reason that living birds often appear smaller to the unpracticed eye than they actually are. The familiar *English Sparrow* is a convenient standard of size because it is usually at hand in our city parks for instant reference. Remembering that it is *6 inches long* a practical though rough division of wild birds may be made as follows:—

1. Birds smaller than the *English Sparrow*.
2. Birds about the size of the *English Sparrow*.
3. Birds decidedly larger than the *English Sparrow*.

If a few general characteristics of the principal bird-families be kept in mind, and these are quickly and almost unconsciously learned, the ¹⁸ identity of a strange bird may usually be narrowed down to a few possibilities. For example:

Woodpeckers climb up and down the trunks of trees bracing with their tails and tapping the bark vigorously;

Nuthatches are smaller than woodpeckers and have much the same habit of climbing up and down tree-trunks but with a freer wig-wagging motion, often descending head downward;

Flycatchers sit erect with drooping tails, watching alertly for insect prey upon which they pounce in mid-air, afterwards returning to their perch;

Swallows skim through the air in graceful and long sustained flights;

Sparrows have stout seed-cracking bills, feed upon the ground, seldom fly high or far at a time and are for the most part fine songsters;

Warblers are tiny, tireless, gaily-colored explorers of the twigs of trees and bushes;

Kinglets are smaller than warblers and quite as restless in their motions, but arrive earlier in the migration;

Wrens, with tails erect, slip mouse-like about brush heaps, crevices and bushes, though often perching in sight while singing;

Thrushes, who with the exception of the Robin and Bluebird are very plainly dressed, run about on the ground stopping suddenly in a listening attitude. When singing they fly up to some perch, although many of the unrivaled singers of this family are silent during their brief sojourn in city parks;

Vireos are most at home on the boughs of trees and sing freely as they glide in and out among the leaves to feed.

Female birds can usually be identified after the adult males have been ¹⁹ seen, although the females and young of many species are obscurely marked or quite different from the adult males, a fact especially true of the warblers. Immature birds are not considered in the present scheme of study since they are a source of confusion to the beginner and occur in any considerable number only during the fall migration.

The time of arrival, that is, when a bird may be expected during the migration, is a point worthy of particular attention. Many wild guesses may be avoided by simply noticing the dates when a bird has been known to arrive in any given locality from year to year.

As a rule birds are identified by the method of elimination. Suppose, for example, a small bird of lively motions is seen feeding among the twigs of a tree late in March. It is, of course, neither a warbler nor a vireo for these birds, although active and frequenting trees, arrive much later in the spring. Provided it does not brace its tail and climb up and around the trunk of the tree it is not likely to be a woodpecker, because the motions of woodpeckers are too characteristic to be mistaken. In similar manner, although sparrows and juncos may have arrived, it is probably neither of these since it does not feed upon the ground, while its size precludes the possibility of its being any bird larger than the *English Sparrow*. There are only a few birds therefore, which it might be and close observation together with the aid of a few

leading hints, will usually settle the question. In any case *watch the bird* for it is better to look at the bird than at a printed description of it.

The safest way, however, is to make careful notes about a stranger on the spot. Memory is never more treacherous than in the case of the description of a doubtful bird when one appeals to some book of reference, a museum collection or the judgment of others. It is not advisable to attempt extensive “field-notes” at first but it is essential to keep a daily, dated list of all identified birds. A blank chart for this purpose is inserted in the back part of the book.

Never *chase* birds. Have patience, stand still a great deal and use your common sense all the time.

Do not make yourself believe that you see a certain bird because it has been reported by others. Do your own looking and listening and do it well.

If you cannot go birding alone take along as few friends as possible because birds are suspicious of human beings in flocks, especially when they talk much.

A pair of field or opera-glasses is an invaluable aid, although practice is necessary in learning to adjust them rapidly and to fix them instantly upon the bird. Do not feel obliged to use the glasses when you are near enough to see well without them.

Keep the sun at your back, otherwise colors will deceive you.

Remember that birds do not stay “indoors” on account of rain, clouds or unfavorable weather. Warm spring rains literally fill the parks with feathered travelers who often tarry but a day.

[11]

Particular Hints.

NOTE:—The following one hundred and fourteen birds are arranged in the order of their *average first appearance in Lincoln Park*, based upon observations made during the last seven years.

For the convenience of those desiring access to reliable sources of information, the scientific name of each bird is placed in parenthesis below its common name, followed by its number in the official list of the American Ornithologist's Union.

Following the hints given about each bird are the names (in **black faced type**) of any bird or birds for which it might easily be mistaken.

1. BLUE JAY. 11-1/2 in.

(Cyanocitta cristata. 477.)

Black collar; *crested*; wings and tail deep blue, white-tipped and black-barred; bold; harsh, noisy call-note; stays the year around in Lincoln Park. [Loggerhead Shrike](#). [Kingfisher](#). [Bluebird](#).

2. LOGGERHEAD SHRIKE. 9 in.

(Lanius ludovicianus. 622.)

Slaty-gray; commonly white beneath, *not marked with dark transverse lines*; wings and tail black, marked with white; *hooked* bill; bar through the eye and *over forehead*, black; imitates notes of other birds. [Blue Jay](#).

3. ROBIN. 10 in.

(Merula migratoria. 761.)

Dark slate color; black head; yellow bill; throat white streaked with black; underparts bright chestnut red; sings “cheer-i-ly, cheer-i-ly, cheer-up;” nests in Lincoln Park. [Towhee](#).

[12

4. JUNCO. 6-1/4 in.

(Junco hiemalis. 567.)

Slate color; blackish bib (female brownish) over a white belly; *ivory bill*, *two white tail-feathers*; feeds in flocks on ground, often in company with other birds; “sucks its teeth” for a call-note; song, a melodious trill.

5. MEADOWLARK. 10-1/2 in.

(Sturnella magna. 501.)

Streaked; black crescent on a *bright yellow* breast; *outer tail-feathers white*; flies straight, *hovering* as it reaches the ground; noteworthy song. [Flicker.](#) [Dickcissel.](#)

6. CROW. 19 in.

(Corvus americanus. 488.)

Entirely black, including bill and feet; often seen in flocks; wings appear frayed and ragged in flight; note, a lusty “caw.” [Bronzed Grackle.](#)

7. BLUEBIRD. 6-1/2 in.

(Sialia sialis. 766.)

Sky-blue; brownish-red below; usually in pairs; sometimes nests in Lincoln Park; call, “pu-ri-ty, pu-ri-ty;” often heard before seen. [Blue Jay.](#)

8. SAVANNA SPARROW. 5-1/4 in.

(Passerculus sandwichensis savanna. 542a.)

Much streaked above and below; line over eye and edge of wing, yellowish; cheek sometimes suffused with yellow tinge; *tail short*, feathers pointed; movements stealthy; song, a weak trill. [Henslow Sparrow.](#)

9. BRONZED GRACKLE. 13 in.

(Quiscalus quiscula æneus. 511b.)

Iridescent black; *body* distinctly bronzy; often carries its tail rudderwise in flight; pompous walk; rusty, grating call; nests in Lincoln Park. [Crow.](#)

[13

10. SONG SPARROW. 6-1/2 in.

(Melospiza cinerea melodia. 581.)

Reddish-brown, showing black streaks; *ashy* line over eye; whitish below with dark-brown streaks which form a *spot in the middle of the breast*; noteworthy song. [Swamp Sparrow.](#) [Lincoln Sparrow.](#)

11. COWBIRD. 8 in.

(*Molothrus ater*. 495.)

Male black with glossy brown head and neck; sparrow-like bill; female brownish; fly in large flocks, uttering a greasy, squeaking note. [Bronzed Grackle.](#) [Rusty Blackbird.](#)

12. TREE SPARROW. 6-1/4 in.

(*Spizella monticola*. 559.)

Streaked; shows much grayish; two *showy white wing-bars*; chestnut-brown cap; breast whitish, *shading to a dark spot in the middle*; in flocks, often with *Juncos*; returns north early in the season; noteworthy song. [Chipping Sparrow.](#) [Field Sparrow.](#)

13. FOX SPARROW. 7-1/2 in.

(*Passarella iliaca*. 585.)

Fox-red back and tail; ashy about neck and head; white breast *thickly* streaked with dark spots; scratches like a hen; alert; noteworthy song. [Brown Thrasher.](#) [Hermit Thrush.](#)

14. PHOEBE. 7 in.

(*Sayornis phoebe*. 456.)

Dull olive-brown; darker on head and tail; whitish below; *bill* and feet black; tail drooping but jerking constantly; note, “phoe´be.” [Wood Pewee.](#) [Least Flycatcher.](#)

15. TOWHEE. 8-3/4 in.

(*Pipilo erythrophthalmus*. 587.)

Black above and black bib (female rich brown); reddish-brown sides; white underneath; *outer tail-feathers white*; scratches about under bushes; brilliant notes, “tow-hee´” and “che-wink´.” [Robin.](#)

16. DOWNY WOODPECKER. 6-1/4 in.

(*Dryobates pubescens medianus*. 394c.)

Bill small; closely resembles the *Hairy Woodpecker* but is *scarcely larger* than the *English Sparrow* and has the outer tail-feathers *barred with black*.

[Hairy Woodpecker. Yellow-bellied Sapsucker.](#)

17. GOLDEN-CROWNED KINGLET. 4 in.

(*Regulus satrapa*. 748.)

Olivaceous; two whitish wing-bars; *orange-yellow stripe edged with black on crown*; restless and fearless; often seen feeding in evergreens; note, rapid “tzee, tzee, tzee.” [Ruby-Crowned Kinglet.](#)

18. FLICKER. 12-1/2 in.

(*Colaptes auratus luteus*. 412a.)

Brown, barred with black; head grayish; back of neck marked with bright red; black crescent on cinnamon-brown spotted breast; in flight shows *white rump* and yellow lining of wings and tail; feeds much on ground, unlike other woodpeckers; call-note, “flee’-ker.” [Meadowlark.](#)

19. BROWN CREEPER. 5-1/2 in.

(*Certhia familiaris americana*. 726.)

Brown finely streaked with white; silky white below; long curved bill; *flies to the base of tree-trunks and works up in spirals* bracing with its long pointed tail-feathers; calls softly, “screeep, screeep.” [Nuthatches.](#)

20. CHICKADEE. 5-1/4 in.

(*Parus atricapillus*. 735.)

Ashy-gray; black throat and *cap*; white cheeks; frowsy; lively; often swings head downward from the tips of twigs; song, “chick-a-dee-dee”; call-note musical, “pewee,” sometimes confused with note of [Phoebe.](#)

21. WHITE-BREASTED NUTHATCH. 5-3/4 in.

(Sitta carolinensis. 727.)

Ashy-blue; cap and back of the neck, black; white below; constantly exploring the bark of trees but does not brace, woodpecker fashion, with its tail-feathers; note, nasal “yank, yank, yank.” [Red-breasted Nuthatch.](#) [Brown Creeper.](#)

22. BELTED KINGFISHER. 14 in.

(Ceryle alcyon. 390.)

Blue-gray; white collar; *big, crested head*; sits erect watching for fish and flies near the surface of the water, uttering a *rattling* call; summers in Lincoln Park.

23. YELLOW-BELLIED SAPSUCKER. 8-1/4 in.

(Sphyrapicus varius. 402.)

Mixed black and white; *yellowish-white underneath*; streaked on sides; white bar *lengthwise* the wing; scarlet patch, bordered with black on crown and throat (female with throat white); harsh, squealing note. [Downy Woodpecker.](#) [Hairy Woodpecker.](#)

24. RUSTY BLACKBIRD. 9-1/2 in.

(Scolopophagus carolinus. 509.)

Black, uniformly iridescent and more or less rusty; tail almost even; female smaller and rustier with pale line over eye. [Bronzed Grackle.](#) [Red-winged Blackbird.](#)

25. FIELD SPARROW. 5-3/4 in.

(Spizella pusilla. 563.)

Streaked; crown and *bill* reddish-brown; neck ashy; brownish-white *unspotted* breast; two indistinct white wing bars; song, “fe-u, fe-u, fu, fee, fee, fee,” beginning slowly and ending almost in a trill. [Chipping Sparrow.](#) [Tree Sparrow.](#)

26. HERMIT THRUSH. 7-1/4 in.

(Hylocichla guttata pallasii. 759b.)

Olive-brown; *tail reddish-brown*; below dull white tinged with buff and thickly spotted across the breast; shy, hiding in bushes; famous for its song but silent during migration. [Other Thrushes.](#)

[16

27. RED-WINGED BLACKBIRD. 9-1/4 in.

(Agelaius phoeniceus. 498.)

Black; scarlet shoulder-straps edged with yellow; female much streaked, marked with rusty; likes marshy places; note, “quonk-a-ree’.” [Cowbird.](#) [Rusty Blackbird.](#)

28. RUBY-CROWNED KINGLET. 4-1/4 in.

(Regulus calendula. 749.)

Dark greenish-olive; light underneath; ring about eye and wing-bars white; male shows at will a brilliant ruby crown-patch; remarkable song. [Golden-crowned Kinglet.](#) [Warblers.](#)

29. HAIRY WOODPECKER. 9 in.

(Dryobates villosus. 393.)

White band down the middle of black back; wings black spotted with white; beneath white; outer tail-feathers white *not barred*; red band on neck of male; large bill; note, sharp resonant “plick.” [Downy Woodpecker.](#) [Yellow-bellied Sapsucker.](#)

30. TREE SWALLOW. 6-1/4 in.

(Iridoprocne bicolor. 614.)

Metallic blue-green; *shining white below*; tail appears nearly square at end in flight; often seen skimming near the surface of still water. [Barn Swallow.](#) [Purple Martin.](#)

31. CARDINAL. 8-1/2 in.

(Cardinalis cardinalis. 593.)

Bright red, including large blunt bill; *crested*; black markings on face and throat; female olive-gray, tinged lightly with red and obscurely marked; fine songster. [Tanagers.](#)

32. WHITE-THROATED SPARROW. 7 in.

(Zonotrichia albicollis. 558.)

White throat; unmarked ashy breast; whitish on belly; crown striped with black and white; broad white stripe over the eye shading to yellow in front; in flocks about shrubbery; clear, sweet whistle, “pee-bod-dee-dee-dee.”

[White-crowned Sparrow.](#)

[17

33. MARSH HAWK. 22 in.

(Circus hudsonius. 331.)

Bluish gray, showing light streaks; gray below, belly white marked with brown; tail barred; *white rump patch*; female brown, barred and streaked more heavily; note shrill.

34. CHIPPING SPARROW. 5-3/4 in.

(Spizella socialis. 560.)

Streaked; *plain ashy breast*; two faint white wing-bars; *chestnut-brown cap*; whitish line over eye; song, a strong clear trill. [Field Sparrow.](#) [Tree Sparrow.](#)

35. MOURNING DOVE. 12 in.

(Zenaidura macroura. 316.)

Gray-blue, tinged with brownish; tail broadly tipped with white, *tapering to a narrow point*; sides of neck iridescent; marked with black on wings and sides of head; rapid whistling flight; note, “coo-oo.” **Pigeon.** [Cuckoos.](#)

36. WINTER WREN. 4-1/4 in.

(Olbiorchiles hiemalis. 722.)

Dark tawny-brown, barred on wings and tail; lighter below, barred posteriorly; pale line over eye; *stub tail, carried erect*; very fine song. [Other Wrens.](#)

37. VESPER SPARROW. 6-1/4 in.

(Poæcetes gramineus. 540.)

Streaked above and below; underparts whitish, tinged with buff across breast and on sides; shows *two conspicuous white tail feathers in flight*; noteworthy song. [Female Purple Finch.](#)

38. MYRTLE WARBLER. 5-1/2 in.

(Dendroica coronata. 655.)

Blue-gray, streaked; crown, *rump* and spot on each side of breast, *yellow*; white below, marked on breast and sides with black; usually in small flocks. [Magnolia Warbler.](#)

[18

39. SWAMP SPARROW. 5-3/4 in.

(Melospiza georgiana. 584.)

Very darkly colored and heavily streaked; chestnut cap, sometimes indistinctly streaked; *black forehead*; plain ashy breast; shy; frequents bushes along the water's edge. [Song Sparrow.](#) [Lincoln Sparrow.](#)

40. PURPLE FINCH. 6-1/4 in.

(Carpodacus purpureus. 517.)

Not "purple," but faded crimson-red, somewhat streaky; belly whitish; *large, heavy bill*; female mottled olive-brown, resembling sparrows; noteworthy song.

41. CEDAR WAXWING. 7 in.

(Ampelis cedrorum. 619.)

Smooth snuff-brown; *crested*; chin and line from bill to crest, black; end of tail banded with yellow; in flocks; “monotonous lisping note.”

42. WILSON THRUSH. 7-1/4 in.

(*Hylocichla fuscescens*. 756.)

Uniform tawny brown; no ring about the eye; below white, faintly tinged and lightly spotted; shy, about bushes; sometimes sings in Lincoln Park. [Other Thrushes.](#)

43. AMERICAN SPARROW HAWK. 10-1/2 in.

(*Falco sparverius*. 360.)

Reddish-brown, usually barred or spotted with black; head bluish, *marked on sides with black*; tail white-tipped with one broad black bar; below white, more or less spotted; narrow pointed wings; female, much more barred, darker and *streaked* below. *The smallest hawk.*

44. BARN SWALLOW. 7 in.

(*Hirundo erythrogastra*. 613.)

Blue-back; chestnut shading to white below; *deeply forked tail*. [Tree Swallow.](#)

[19

45. BROWN THRASHER. 11-1/4 in.

(*Toxostoma rufum*. 705.)

Bright rusty red; white below, much streaked; long bill and *long, sweeping tail*; song, loud, bold and varied; about bushes. [Fox Sparrow.](#)

46. HOUSE WREN. 5 in.

(*Troglodytes aedon*. 721.)

Brown, barred with darker; below grayish, tinged with brown, also barred; rusty-colored tail, *two inches long*; remarkable song. [Other Wrens.](#)

47. BLUE-GRAY GNATCATCHER. 5 in.

(Polioptila caerulea. 751.)

Blue-gray; white beneath; *long tail* edged with white; male has a black line over eye; fidgety, never long in a place; wheezy song. [Black-throated Blue Warbler.](#) [Cerulean Warbler.](#)

48. WHITE-CROWNED SPARROW. 7 in.

(Zonotrichia leucophrys. 554.)

Ashy-gray; rump, tail and wings, brown, streaked with black; *crown conspicuously striped* with black and white; no yellow; two white wing-bars; in flocks, often with *White-throated Sparrows*; scratches with both feet; notice the song. [White-throated Sparrow.](#)

49. PALM WARBLER. 5 in.

(Dendroica palmarum. 672.)

Streaky olive-gray; *chestnut cap*; yellow line over eye; beneath yellowish (brightest on throat and under tail) streaked with brown; feeds on ground in flocks; constantly *tilts its tail* up and down. [Water-Thrushes.](#)

[20

50. HENSLOW SPARROW. 5 in.

(Coturniculus henslowii. 547.)

Back conspicuously marked and streaked, showing much chestnut; head and neck a peculiar greenish tinge, *finely* lined with black; below tinged with buff and finely streaked; dark stripes and spots on sides of head; *sharp tail*; hides in the grass and can be carefully observed only with patience.

51. GRINNELL WATER-THRUSH. 5-3/4 in.

(Seiurus noveboracensis notabilis. 675a.)

Very dark olive-brown; below yellowish white, thickly streaked; *pale stripe* over the eye; skulks along the very edge of the water; *tilts its tail*; brilliant song. [Louisiana Water-Thrush.](#) [Ovenbird.](#) [Palm Warbler.](#)

52. CATBIRD. 8-3/4 in.

(Galeoscoptes carolinensis. 704.)

Slate-gray; bill, crown and tail, black; chestnut patch under tail; about shrubbery; willing to be looked at; calls like a cat; song, greatly varied.

53. PINE WARBLER. 5-3/4 in.

(Dendroica vigorsii. 671.)

Olive-green; yellow breast; sides of head suffused with yellowish—sometimes forming a *yellow line over eye*; two dull white wing-bars; song, a loud, penetrating trill. [Yellow-bellied Flycatcher. Female Western Yellow-throat. Female Wilson Black-cap.](#)

54. RED-BREASTED NUTHATCH. 4-3/4 in.

(Sitta canadensis. 728.)

Bluish-gray; crown and *stripe through eye, black*, with a white line over the eye; *below rusty-red*, throat white; climbs up and down and around trees; note, “cark, cark, cark.” [White-breasted Nuthatch.](#)

[21

55. BLACK AND WHITE CREEPER. 5-1/4 in.

(Mniotilta varia. 636.)

Black and white, *striped*; white belly; female shows more white below; creeps up and down trees like the [Nuthatches](#); song rapid and rasping, “he’-ho, he’-ho, he’-ho.” [Black-poll Warbler.](#)

56. BLACK-THROATED GREEN WARBLER. 5-1/4 in.

(Dendroica virens. 667.)

Olive-green; *bright yellow cheeks; throat and upper part of breast, black*; white wing-bars; song, repeated often and rapidly, “chee, chee, chee, chee, char, chee.” [Cape May Warbler. Hooded Warbler.](#)

57. RED-HEADED WOODPECKER. 9-1/2 in.

(Melanerpes erythrocephalus. 406)

Black and white, not streaked; *entire head, neck and upper part of breast, crimson*; pure white belly; note, a loud “kr-r-r-r r.” [Red-bellied Woodpecker.](#)

58. MOCKINGBIRD. 10 in.

(Mimus polyglottus. 703.)

Ashy-gray, wings and tail darker; whitish below; wing shows white patch; outer tail-feathers white; famous for its song. [Loggerhead Shrike.](#) [Cuckoos.](#) [Catbird.](#)

59. CHIMNEY SWIFT. 5-1/4 in.

(Chætura pelagica. 423.)

Dark sooty-gray; *bob-tail*; long, pointed wings, which often vibrate rapidly in flight; twittering note; never seen perching like other birds. [Swallows.](#)

60. OLIVE-BACKED THRUSH. 7-1/4 in.

(Hylocichla ustulata. 758a.)

Uniform dark olive; below white, darker on sides; cheeks, throat, breast and ring about eye, buff; spotted on breast; noteworthy song. [Gray-cheeked Thrush.](#) [Hermit Thrush.](#) [Wilson Thrush.](#)

[22

61. GREAT-CRESTED FLYCATCHER. 9 in.

(Myiarchus crinitus. 452.)

Olive, showing rusty brown on wings and tail; *crested head*; throat and breast, ashy-gray; wing-bars; belly sulphur-yellow; usually perches high; note, “loud whistle.” [Blue Jay.](#)

62. LOUISIANA WATER-THRUSH. 6-1/4 in.

(Seiurus motacilla. 676.)

Olive; white below, buff-tinged posteriorly and streaked, but not thickly, with dusky; *throat and middle of belly, unmarked*; *white* line over eye; loud,

ringing song. [Grinnell Water-Thrush](#). [Ovenbird](#). [Palm Warbler](#).

63. LEAST FLYCATCHER. 5-1/4 in.

(Empidonax minimus. 467.)

Olive-gray; wings, tail and head, darker; two narrow wing-bars; decidedly grayish below, whiter on throat; *jerks its tail* and utters a sharp, energetic call, “che-bec’.” [Wood Pewee](#). [Phoebe](#).

64. WOOD THRUSH. 7-3/4 in.

(Hylocichla mustelina. 755.)

Bright brown on head shading to olive-brown on rump and tail; white beneath, with conspicuous *round* spots except on throat and middle of belly; remarkable song. [Other Thrushes](#). [Fox Sparrow](#).

65. WESTERN YELLOW-THROAT. 5-1/4 in.

(Geothlypis trichas occidentalis. 681a.)

Olive-green, showing brownish tinge; *broad, black stripe across forehead and through eye*; bordered broadly with white; yellow beneath; *female* duller and without black stripe and easily confused with female [Mourning](#), [Connecticut](#) and [Pine Warblers](#); in low bushes; song, “witch-i-ty, witch-i-ty, witch-i-ty.”

[23

66. MAGNOLIA WARBLER. 5 in.

(Dendroica maculosa. 657.)

Back and tail, black; crown blue-gray; yellow rump; *yellow* below, streaked with black on breast and sides; white patch on wings; broad white band across tail, seen from below. [Myrtle Warbler](#). [Cape May Warbler](#).

67. AMERICAN GOLDFINCH. 5-1/4 in.

(Astragalinus tristis. 529.)

Canary-yellow, with black crown, tail and wings; white wing-bars; female olive-tinged, without pure black or yellow; in flocks; canary-like song; also sings in its undulating flight, “per-*chic*-o-ree, per-*chic*-o-ree.” [Yellow Warbler.](#)

68. YELLOW WARBLER. 5 in.

(Dendroica aestiva. 652.)

Yellow all over shading to olive on back; orange-brown streaks on breast; female not so yellow and less streaked; song, penetrating and unceasing; nests in Lincoln Park. [American Goldfinch.](#)

69. KINGBIRD. 8-1/2 in.

(Tyrannus tyrannus. 644.)

Dusky black; below white; *tail tipped with broad band of white*; fire-red patch on crown, usually concealed; spreads its tail like a fan when about to alight.

70. BALTIMORE ORIOLE. 7-1/2 in.

(Icterus galbula. 507.)

Bright orange, with black head, throat, back and wings; two white wing-bars; female smaller, brownish-yellow and showing little black; loud call, “co-weet’, co-weet’, co-weet’;” song noteworthy. [Orchard Oriole. Scarlet Tanager.](#)

71. SUMMER TANAGER. 7-1/2 in.

(Piranga rubra. 610.)

Vermillion all over; no black; female yellowish-olive and usually near the male; very rare. [Cardinal.](#)

72. YELLOW-BELLIED FLYCATCHER. 5-1/2 in.

(Empidonax flaviventris. 463.)

Bright olive-green, with dark-brown tail and wings; yellowish wing-bars; sulphur-yellow below, *brightest between legs* and tinged elsewhere with pale olive; yellow eye-ring; call, a soft, slow whistle, “pe´-a.” [Yellow-throated Vireo. Least Flycatcher. Pine Warbler.](#)

73. BLACKBURNIAN WARBLER. 5-1/4 in.

(Dendroica blackburniae. 662.)

Black streaked with white; middle of crown, throat and sides of head, *flaming orange*; black patch behind eye; white wing-bars; female duller; usually high up among the leaves. [Prothonotary Warbler.](#)

74. OVENBIRD. 6-1/4 in.

(Seiurus aurocapillus. 674.)

Olive-green showing yellow tinge; below pure white, thickly spotted; *orange-brown cap* bordered with black stripes; scratches like a hen; tilts tail up and down; under shrubbery; note, “teacher” repeated rapidly with increasing emphasis. [Water-Thrushes.](#)

75. RED-EYED VIREO. 6 in.

(Vireo olivaceus. 624.)

Olive-green; crown, slaty; white below, lightly tinged on sides; *dark line above a white one over eye*; clear, persistent song; nests in Lincoln Park. [Warbling Vireo.](#)

76. AMERICAN REDSTART. 5-1/4 in.

(Setophaga ruticilla. 687.)

Male black, with *six flame-colored spots*; female olive, with *six yellow spots*; spreads its tail like a fan; song, “chee, chee, chee, chew.”

77. CAPE MAY WARBLER. 5-1/4 in.

(Dendroica tigrina. 650.)

Yellowish-olive spotted with black on back; crown very dark; cheeks yellow with an *orange-brown ear patch*; yellow below, black streaks on breast; wing-bars; song all on one note. [Black-throated Green Warbler](#). [Magnolia Warbler](#).

78. SCARLET TANAGER. 7-1/4 in.

(*Piranga erythromelas*. 608.)

Scarlet, with black wings and tail; female olive-green, usually near by; rather sluggish; note, “chip, chree;” song noteworthy. [Cardinal](#). [Summer Tanager](#).

79. YELLOW-THROATED VIREO. 5-1/2 in.

(*Vireo flavifrons*. 628.)

Bright olive-green, shading to ashy-blue on rump; *yellow throat* and breast, sharply defined against white belly; line around eye and to the bill, yellow; two white wing-bars; fine song. [Yellow-bellied Flycatcher](#). [Blue-headed Vireo](#).

80. ROSE-BREASTED GROSBEAK. 8-1/4 in.

(*Zamelodia ludoviciana*. 595.)

Black, including throat, with white rump and wing-patches; breast white with a brilliant rose-red blotch; lining of wings also rose-red; *large beak*; female yellowish-brown, darkly mottled, no red; sluggish; remarkable song.

81. PURPLE MARTIN. 7-1/2 in.

(*Progne subis*. 611.)

Blue-black all over; female lighter and streaked with grayish-white below; often nests in bird houses; twittering song. [Tree Swallow](#). [Chimney Swift](#).

82. YELLOW-BREASTED CHAT. 7-1/4 in.

(*Icteria virens*. 683.)

Olive-green; bright yellow below, belly *white*; region about eye black, marked with a white line above eye; white spot and shorter line below eye; the largest warbler; remarkable song. [Dickcissel. Western Yellow-Throat. Yellow-Throated Vireo.](#)

83. BLACK-THROATED BLUE WARBLER. 5-1/4 in.

(Dendroica caerulescens. 654.)

Slaty-blue; below white, with black throat and sides; *conspicuous white patch on wing*; female grayish, no black, but may be recognized by *wing patch*; hoarse note, “kree, kree.” [Blue-gray Gnatcatcher.](#)

84. NASHVILLE WARBLER. 4-3/4 in.

(Helminthophila rubricapilla. 645.)

Olive-green; head and neck bluish gray; indistinct brown patch on crown; *bright yellow below*; no wing bars; *white eye ring*; song, not loud, “wee´-see, wee´-see, wit´-a-wit´-a-wit´.” [Tennessee Warbler.](#)

85. WARBLING VIREO. 5-1/4 in.

(Vireo gilvus. 627.)

Mouse gray, with olive rump; white below, tinged with greenish yellow; creamy-white around the eye; song, rippling warble. [Red-eyed Vireo. Least Flycatcher.](#)

86. BLUE-HEADED VIREO. 5-1/2 in.

(Vireo solitarius. 629.)

Olive-green; *ashy-blue head*; pure white below, sides yellowish; line around the eye and to the bill, white; two white wing-bars; song, noteworthy. [Yellow-throated Vireo.](#)

87. ORCHARD ORIOLE. 7 in.

(Icterus spurius. 506.)

Male, rich *chestnut-brown* with black head, wings and tail, narrow white wing-bar; female olivaceous above and greenish-yellow below with two white wing-bars; young male like female but with black throat; high trees; loud, clear song. [Baltimore Oriole.](#)

88. LONG-BILLED MARSH WREN. 5-1/4 in.

(Telmatodytes palustris. 725.)

Brown, almost black on head, tail and between shoulders, the latter streaked with white; barred only on wings and tail; *line over eye* and middle of breast white; *long bill*; spluttering song. [Other Wrens.](#)

89. BAY-BREASTED WARBLER. 5-1/2 in.

(Dendroica castanea. 660.)

Ashy-brown, black-streaked; black cheeks and forehead; throat, upper part of breast, sides and crown, *rich chestnut brown*; white wing-bars. [Chestnut-sided Warbler.](#)

90. PARULA WARBLER. 4-1/2 in.

(Compothlypis americana. 648.)

Grayish-blue, *yellowish across the back*; throat and breast yellow, latter crossed by brownish band; wing-bars; peculiar song. [Blue Golden-winged Warbler.](#)

91. WOOD PEWEE. 6-1/4 in.

(Contopus virens. 461.)

Dark olive-brown; brownish black on wings and tail; frowsy head; below white, showing olive tinge on sides; two narrow white wing-bars; plaintive note repeated slowly and often, “pee´to-way´, pee´-to-way´, hee´-rue.” [Phoebe. Least Flycatcher.](#)

92. BOBOLINK. 7-1/2 in.

(Dolichonyx orizivorus. 494.)

Male black with light back, creamy buff on neck; female, sparrow colored, streaked above; long merry musical song; open fields; in Lincoln Park usually seen in flocks flying overhead, singing on the wing.

[28

93. BLUE GOLDEN-WINGED WARBLER. 5 in.

(*Helminthophila chrysoptera*. 642.)

Blue-gray; crown and *patch on wing*, golden yellow; *throat*, and stripe through eye, *black*; white below; song, drawling “zee, zee, zee, zee.” [Parula Warbler.](#)

94. INDIGO BUNTING. 5-3/4. in.

(*Cyanospiza cyanea*. 598.)

Indigo blue; wings and tail blackish; female brown; in flocks; often feeds in the grass; noteworthy song. [Bluebird.](#)

95. PROTHONOTARY WARBLER. 5-1/4 in.

(*Prothonotaria citrea*. 637.)

Entire head, neck and breast, *bright orange yellow*; blue-gray wings, rump and tail; female duller; prefers shrubbery near water. [Blackburnian Warbler.](#) [Yellow Warbler.](#)

96. TENNESSEE WARBLER. 4-1/2 in.

(*Helminthophila peregrina*. 647.)

Olive-green, brightest on rump; head and neck, blue gray; *dull white below*; faint white line over eye; no wing-bars; song, shrill and feeble. [Nashville Warbler.](#)

97. MOURNING WARBLER. 5-1/2 in.

(*Geothlypis philadelphia*. 679.)

Bright olive-green; head, neck, throat and upper part of breast, *bluish ash*; yellow belly; black band on breast; about bushes and in the grass;

noteworthy song. [Connecticut Warbler.](#)

98. CHESTNUT-SIDED WARBLER. 5 in.

(*Dendroica pennsylvanica*. 659.)

Back shows black streaks mixed with olive-green; yellow cap; white below with dainty *chestnut markings on sides of breast*; song, “chee, chee, chee, chee’ar.” [Bay-breasted Warbler.](#)

[29

99. WHIP-POOR-WILL. 9-3/4 in.

(*Antrostomus vociferus*. 417.)

Mottled, strongly resembling the bark of certain trees; narrow white band across upper part of breast; big head; in daytime lies *motionless, lengthwise a branch*. [Nighthawk.](#)

100. CERULEAN WARBLER. 4-1/2 in.

(*Dendroica caerulea*. 658.)

Azure-blue, marked with black on back and sides of head; white below, streaked with blue; two white wing-bars; frequents tree-tops; song resembles that of [Parula](#). [Blue-gray Gnatcatcher.](#)

101. HOODED WARBLER. 5-1/2 in.

(*Wilsonia mitrata*. 684.)

Olive-green; yellow below; head, neck and upper breast, black; *yellow “mask” across forehead and cheeks*; white spots on tail; female less black. [Western Yellow-throat.](#) [Wilson Black-cap Warbler.](#) [Canadian Warbler.](#)

102. BLACK-POLL WARBLER. 5-1/2 in.

(*Dendroica striata*. 661.)

Streaked black and white; *black crown*; white cheeks; below white with line of black spots down sides; tree-tops; song, a staccato musical “chink” repeated rapidly. [Black and White Creeper.](#)

103. YELLOW-BILLED CUCKOO. 12-1/4 in.

(Coccyzus americanus. 387.)

Bronzy olive-gray; white below; *lower half of curved bill yellow* tipped with black; outer tail-feathers black tipped *broadly* with white; wings show a *reddish-brown tinge in flight*; glides stealthily from tree to tree and keeps concealed among the leaves; slowly droops and raises its tail when perching; song, loud “kuk-kuk, kuk-kuk.” [Black-billed Cuckoo. Mourning Dove.](#)

[30

104. CANADIAN WARBLER. 5-1/2 in.

(Wilsonia canadensis. 686.)

Bluish-lead color; yellow below with a necklace of black spots; no wing-bars; noteworthy song. [Magnolia Warbler.](#)

105. WILSON BLACK-CAP WARBLER. 4-3/4 in.

(Wilsonia pusilla. 685.)

Bright olive-green; yellow below; *black cap*; in low bushes. [Pine Warbler. Female Western Yellow-throat.](#)

106. LINCOLN SPARROW. 5-3/4 in.

(Melospiza lincolni. 583.)

Finely streaked with black and brown; white beneath, with **broad buff band** across the breast and down the sides; narrow streaks on throat, upper breast and sides; dark stripe on each side of throat from corner of mouth; very shy. [Song and Swamp Sparrows.](#)

107. RUBY-THROATED HUMMINGBIRD. 3-1/4 in.

(Trochilus colubris. 428.)

Iridescent green and purple; male has ruby-red throat; frequents flowering shrubs and plants; darts here and there and poises in mid air with wings vibrating so rapidly that they make a humming noise; “squeaky note.”

108. NIGHTHAWK. 9-1/2 in.

(Chordeiles virginianus. 420.)

On the wing overhead after insects just before or at dusk; long, pointed wings *showing a white patch*; flight resembles that of [Chimney Swift](#); sharp note, “skeep;” often nests on the flat roofs of city buildings. [Whip-poor-will.](#)

109. BLACK-BILLED CUCKOO. 12 in.

(Coccyzus erythrophthalmus. 388.)

Bronzy olive-gray; white below; bill *entirely* black; *only a little white* on tail; movements like those of *Yellow-billed Cuckoo*; song, soft “coo-coo, coo-coo.” [Yellow-billed Cuckoo. Mourning Dove.](#)

[31]

110. GRAY-CHEEKED THRUSH. 7-1/2 in.

(Hylocichla aliciae. 757.)

Uniform olive; below white, *very palely* suffused with buff; cheeks *gray tinged*; breast and sides of throat, spotted; usually silent in Lincoln Park. [Olive-backed](#) and [other Thrushes.](#)

111. YELLOW-HEADED BLACKBIRD. 10-1/4 in.

(Xanthocephalus xanthocephalus. 497.)

Black; head, throat and breast, *yellow*; white wing patch; female dark brown, yellow markings duller, no wing patch; female smaller than male.

112. DICKCISSEL. 6 in.

(Spiza americana. 604.)

Streaked; ashy on head and neck; white chin; *black throat patch*; *yellow breast*; white belly; *line over eye, yellow*; wings show chestnut brown; female has less yellow and no throat patch. [Yellow-breasted Chat. Meadowlark.](#)

113. CONNECTICUT WARBLER. 5-1/2 in.

(Geothlypis agilis. 678.)

Olive-green; head ashy with throat darker; *white ring* around eye; belly yellow; wings long and pointed; low bushes and swampy places; penetrating call-note. [Mourning Warbler](#).

114. RED-BELLIED WOODPECKER. 9-1/2 in.

(Centurus carolinus. 409.)

Black and white, back, wings and tail *barred transversely* and very evenly; below grayish-white, *tinged with red*; top of head and back of neck crimson; female, crown gray; rare. [Red-headed Woodpecker](#). [Hairy Woodpecker](#).

[32]

Table of Occurrence.

In this table the birds are arranged according to the order of their comparative frequency or rarity, based upon personal observations in *Lincoln Park*. The abundance of the birds—that is, the number of individual birds of different species—is in no wise shown, but simply the relative representation of each species. For instance, out of a total of 454 mornings, covering observations during six years, the [Robin](#) was seen on 383 different mornings.

The record for 1902, during the absence of the authors, is that of Mr. H. V. Bozell.

[33]

**NUMBER OF MORNINGS SEEN IN
LINCOLN PARK.**

	1898	1899	1900	1901	1902	1903	Total
1 Robin	62	52	57	69	61	82	383
2 Bronzed Grackle	57	58	57	60	57	72	361
3 Blue Jay	49	58	61	65	40	51	324
4 Cowbird	41	34	44	44	39	54	256
5 Junco	33	35	32	32	51	42	225

6 Towhee	35	18	24	44	45	48	214
7 Song Sparrow	27	33	19	38	37	36	190
8 Flicker	29	18	21	33	37	46	184
9 Ruby-crowned Kinglet	31	28	23	30	22	39	173
10 White-throated Sparrow	30	23	23	31	26	35	168
11 Bluebird	26	1	7	31	35	50	150
12 Wilson Thrush	34	35	23	21	11	20	144
13 Yellow-bellied Sapsucker	23	12	15	30	27	34	141
14 Yellow Warbler	27	30	24	18	13	27	139
15 Belted Kingfisher	25	21	24	27	21	19	137
16 Red-headed Woodpecker	23	23	16	26	16	26	130
17 Myrtle Warbler	19	25	11	21	22	31	129
18 Brown Thrasher	27	8	20	21	16	29	121
19 Brown Creeper	18	15	19	14	28	27	121
20 Catbird	22	27	16	17	10	17	109
21 Golden-crowned Kinglet	18	14	13	26	15	23	109
22 Chipping Sparrow	25	11	11	12	17	32	108
23 American Goldfinch	21	19	12	20	5	18	95
24 Tree Swallow	13	11	16	19	13	22	94
25 American Redstart	18	16	19	8	10	20	91
26 Barn Swallow	15	14	10	19	14	19	91
27 Fox Sparrow	11	9	14	9	27	21	91
28 Chimney Swift	16	11	14	15	11	22	89
29 Blue-gray Gnatcatcher	18	16	11	10	16	17	88
30 White-crowned Sparrow	9	5	18	22	16	17	87
31 Palm Warbler	8	21	7	11	19	20	86

32 Hermit Thrush	13	9	10	18	9	27	86
33 Black and White Creeper	15	15	11	13	12	18	84
34 Least Flycatcher	18	18	8	14	7	18	83
35 Magnolia Warbler	13	13	13	5	16	20	80
36 Black-throated Blue Warbler	12	14	9	13	12	16	76
37 Olive-backed Thrush	8	4	14	16	11	23	76
38 Phoebe	15	4	5	16	15	19	74
39 Baltimore Oriole	18	6	10	15	11	13	73
40 Grinnell Water-Thrush	17	6	9	14	8	17	71
41 Field Sparrow	..	2	12	17	23	17	71
42 Swamp Sparrow	4	4	14	12	17	17	68
43 Crow	4	11	7	11	16	18	67
44 Chestnut-sided Warbler	11	6	11	13	9	16	66
45 Ovenbird	9	11	11	12	11	11	65
46 White-breasted Nuthatch	12	6	17	21	6	2	64
47 Western Yellow-throat	18	17	10	8	3	15	61
48 Wood Pewee	12	14	9	6	7	13	61
49 Downy Woodpecker	14	7	4	8	17	10	60
50 House Wren	1	..	10	11	13	24	59
51 Red-eyed Vireo	13	10	5	6	6	16	56
52 Loggerhead Shrike	5	1	16	13	7	13	55
53 Indigo Bunting	12	17	2	6	5	11	53
54 Black-throated Green Warbler	4	9	4	9	6	12	44
55 Blackburnian Warbler	11	4	9	7	5	6	42

56 Black-poll Warbler	6	7	8	3	3	15	42
57 Kingbird	8	1	5	3	6	17	40
58 Scarlet Tanager	7	3	7	4	3	10	34
59 Ruby-throated Hummingbird	7	9	4	3	1	10	34
60 Mourning Dove	5	3	6	10	2	8	34
61 Rose-breasted Grosbeak	5	2	6	8	6	6	33
62 Tree Sparrow	..	..	10	3	12	8	33
63 Winter Wren	3	5	11	..	3	10	32
64 Meadowlark	3	..	7	13	5	4	32
65 Purple Martin	1	..	..	6	4	20	31
66 Cape May Warbler	3	2	9	2	6	7	29
67 Warbling Vireo	7	1	4	3	4	10	29
68 Mourning Warbler	5	8	2	4	4	5	28
69 Wilson Black-cap Warbler	9	1	3	2	5	8	28
70 Yellow-billed Cuckoo	8	7	1	3	1	7	27
71 Great-crested Flycatcher	5	2	..	1	10	9	27
72 Bobolink	2	..	9	10	1	4	26
73 Canadian Warbler	8	3	6	4	1	4	26
74 Red-winged Blackbird	4	..	2	8	3	7	24
75 Wood Thrush	8	3	..	5	3	3	22
76 Orchard Oriole	4	3	5	1	1	6	20
77 Red-breasted Nuthatch	..	..	10	..	3	7	20
78 Marsh Hawk	..	2	1	8	5	4	20
79 Hairy Woodpecker	3	1	4	..	4	7	19
80 Bay-breasted Warbler	1	1	2	1	4	10	19

81	Parula Warbler	6	..	..	2	9	2	19
82	Chickadee	3	..	7	8	..	..	18
83	Pine Warbler	..	..	3	1	5	9	18
84	Henslow Sparrow	3	2	1	2	3	6	17
85	Nashville Warbler	1	1	4	1	..	9	16
86	Yellow-throated Vireo	2	..	2	2	3	7	16
87	Cedar Waxwing	6	5	..	1	2	1	15
88	Purple Finch	4	..	4	..	1	5	14
89	Vesper Sparrow	..	..	6	..	4	3	13
90	Blue-headed Vireo	..	..	2	1	6	4	13
91	Gray-cheeked Thrush	..	..	..	..	3	10	13
92	American Sparrow Hawk	1	1	..	4	2	4	12
93	Yellow-bellied Flycatcher	..	..	5	..	2	4	11
94	Nighthawk	..	2	1	1	..	7	11
95	Lincoln Sparrow	..	..	1	..	1	8	10
96	Black-billed Cuckoo	..	1	2	2	2	2	9
97	Rusty Blackbird	..	2	..	3	..	2	7
98	Yellow-breasted Chat	..	..	..	..	1	6	7
99	Long-billed Marsh Wren	1	2	..	..	..	3	6
100	Louisiana Water-Thrush	..	..	1	..	3	2	6
101	Prothonotary Warbler	..	1	..	..	3	1	5
102	Savanna Sparrow	..	..	..	..	..	5	5
103	Blue Golden-winged Warbler	1	..	1	1	1	..	4
104	Whip-poor-will	1	1	..	1	..	1	4

105 Connecticut Warbler	..	..	..	..	..	3	3
106 Dickcissel	..	..	..	1	2	..	3
107 Tennessee Warbler	2	..	..	..	..	..	2
108 Mockingbird	..	..	..	2	..	..	2
109 Cerulean Warbler	..	..	1	..	1	..	2
110 Summer Tanager	..	1	..	..	..	..	1
111 Cardinal	..	..	1	..	..	..	1
112 Red-bellied Woodpecker	1	..	..	..	..	..	1
113 Hooded Warbler	..	..	..	..	..	1	1
114 Yellow-headed Blackbird	..	..	..	1	..	..	1
Total number of days	66	77	74	78	72	87	454
Av'ge number of kinds seen daily	18.83	12.75	14.59	16.23	16.45	19.53	16.34

[35]

Table of Arrival.

NOTE.—The following table gives the dates of the first arrival of each bird mentioned in the preceding list of one hundred and fourteen for the seven years from 1897 to 1903 inclusive, *as observed in Lincoln Park*. The birds are arranged in their respective families, and space is left for recording their first appearance in any succeeding year.

The record for 1902, in the absence of the authors, was kept by Mr. Harold V. Bozell.

Attention is called to the fact that many birds which are common summer residents in the neighboring country are rare migrants in the Park. Among these may be mentioned the [Red-winged Blackbird](#), [Bobolink](#) and [Warbling Vireo](#). Some fine singers also, for example, the [Vesper Sparrow](#), [Yellow-throated Vireo](#) and [Hermit Thrush](#), are usually silent in the Park.

[36]

	1897	1898	1899	1900	1901	1902	1903	_____
Family COLUMBIDAE								
35 Mourning Dove	..	5/9	3/25	4/6	4/10	4/26	4/1	_____
Family FALCONIDAE								
33 Marsh Hawk	..	..	4/25	4/19	3/25	4/21	4/1	_____
43 Sparrow Hawk	..	5/18	4/25	4/18	3/24	3/21	4/30	_____
Family CUCULIDAE								
103 Yellow-billed Cuckoo	..	5/19	5/1	5/16	5/17	5/17	5/12	_____
109 Black-billed Cuckoo	..	..	5/27	5/12	5/15	5/16	5/12	_____
Family ALCEDINIDAE								
22 Belted Kingfisher	3/28	3/25	3/22	4/10	3/26	4/11	4/10	_____
Family PICIDAE								
29 Hairy Woodpecker	..	4/11	5/24	4/19	..	3/25	3/6	_____
16 Downy Woodpecker	4/7	4/8	3/27	4/25	4/5	3/10	3/6	_____
23 Yellow-bellied Sapsucker	4/25	3/30	4/8	4/6	3/19	3/27	4/1	_____
114 Red-bellied Woodpecker	..	5/22	..	..	..	..	..	_____
57 Red-headed Woodpecker	4/26	4/29	4/28	5/6	4/27	4/27	4/28	_____
18 Flicker	4/15	3/25	4/11	4/7	3/23	3/24	3/18	_____
Family CAPRIMULGIDAE								
99 Whip-poor-will	..	5/21	4/30	..	5/9	..	5/17	_____
108 Nighthawk	..	..	5/10	5/13	5/18	5/18	5/18	_____
Family MICROPODIDAE								
59 Chimney Swift	5/1	4/30	5/11	4/30	4/17	4/22	5/5	_____
Family TROCHILIDAE								
107 Ruby-throated Hummingbird	..	5/19	5/9	5/13	5/16	5/17	5/13	_____
Family TYRANNIDAE								
69 Kingbird	4/29	4/30	5/10	5/6	5/10	4/21	5/5	_____
61 Great-crested Flycatcher	..	5/3	4/30	..	5/11	4/21	4/29	_____
14 Phoebe	4/7	3/20	4/17	4/1	3/18	3/15	3/17	_____

91	Wood Pewee	4/27	5/19	5/10	5/11	5/16	4/28	5/12	_____
72	Yellow-bellied Flycatcher	..	..	..	4/29	..	5/1	5/12	_____
63	Least Flycatcher	5/12	4/30	4/20	4/30	5/6	4/21	5/9	_____
Family CORVIDAE									
1	Blue Jay	..	..	..	..	..	..	..	_____
6	Crow	..	3/9	3/26	3/12	3/3	3/6	3/8	_____
Family ICTERIDAE									
92	Bobolink	..	5/14	..	5/8	5/4	5/12	5/8	_____
11	Cowbird	4/4	3/16	4/10	4/1	3/19	3/1	3/14	_____
111	Yellow-headed Blackbird	..	..	..	..	5/17	..	..	_____
27	Red-winged Blackbird	..	5/7	..	4/18	3/22	3/26	3/17	_____
5	Meadowlark	..	3/14	..	3/27	3/19	3/10	3/13	_____
87	Orchard Oriole	..	5/15	5/6	5/11	5/10	5/2	5/5	_____
70	Baltimore Oriole	..	4/30	5/2	5/6	5/4	5/2	5/5	_____
24	Rusty Blackbird	..	..	..	4/19	4/5	..	3/17	_____
9	Bronzed Grackle	3/29	3/14	3/20	3/24	3/23	3/10	3/14	_____
Family FRINGILLIDAE									
40	Purple Finch	..	4/24	..	4/15	..	4/11	4/15	_____
67	American Goldfinch	4/25	5/6	4/28	5/7	4/30	5/2	5/7	_____
37	Vesper Sparrow	..	..	..	4/22	..	4/18	4/1	_____
8	Savanna Sparrow	..	..	..	..	..	..	3/19	_____
50	Henslow Sparrow	..	4/29	4/26	4/21	4/30	4/24	4/15	_____
48	White-crowned Sparrow	5/5	5/9	4/29	5/2	4/20	3/11	4/29	_____
32	White-throated Sparrow	4/26	4/13	4/19	4/21	4/17	3/24	3/21	_____
12	Tree Sparrow	..	..	..	4/6	4/5	3/7	3/13	_____
34	Chipping Sparrow	4/15	4/13	4/18	4/8	4/8	4/13	4/12	_____
25	Field Sparrow	..	..	4/23	4/17	4/5	3/11	3/19	_____
4	Junco	3/28	3/9	3/14	3/23	3/18	3/8	3/13	_____
10	Song Sparrow	4/8	3/15	3/19	3/27	3/18	3/8	3/12	_____
106	Lincoln Sparrow	..	..	..	5/10	5/23	5/12	5/10	_____

39 Swamp Sparrow	..	5/7	4/20	4/21	4/18	3/11	4/12	_____
13 Fox Sparrow	4/8	3/30	3/14	4/5	3/18	3/10	3/17	_____
15 Towhee	4/7	3/17	4/11	4/7	3/25	3/10	3/18	_____
31 Cardinal	3/28	..	..	4/25	..	..	..	_____
80 Rose-breasted Grosbeak	..	..	5/10	5/11	5/4	4/28	5/9	_____
94 Indigo Bunting	..	5/14	5/10	5/21	5/10	5/2	5/6	_____
112 Dickcissel	..	..	..	..	5/16	5/18	..	_____
Family TANAGRIDAE								
78 Scarlet Tanager	..	5/14	5/4	5/2	5/5	5/3	5/5	_____
71 Summer Tanager	..	..	5/4	..	..	..	..	_____
Family HIRUNDINIDAE								
81 Purple Martin	..	5/25	5/30	..	4/27	4/23	4/17	_____
44 Barn Swallow	4/18	5/1	4/23	4/26	4/10	4/18	4/24	_____
30 Tree Swallow	..	5/11	4/12	4/18	4/12	3/11	4/1	_____
Family AMPELIDAE								
41 Cedar Waxwing	..	4/11	3/27	..	5/16	4/24	4/5	_____
Family LANIIDAE								
2 Loggerhead Shrike	..	3/14	4/11	3/9	3/3	3/9	3/1	_____
Family VIREONIDAE								
75 Red-eyed Vireo	..	5/10	4/29	5/12	5/4	5/2	5/4	_____
85 Warbling Vireo	4/27	5/15	5/12	5/6	5/11	5/7	5/5	_____
79 Yellow-throated Vireo	..	5/1	..	5/12	5/5	4/28	5/12	_____
86 Blue-headed Vireo	..	..	..	5/17	5/5	5/2	5/8	_____
Family MNIOTILTIDAE								
55 Black and White Creeper	4/28	4/30	4/21	4/26	5/4	4/28	4/27	_____
95 Prothonotary Warbler	..	..	5/16	..	..	4/27	5/19	_____
93 Blue Golden-winged Warbler	..	5/14	..	5/12	5/12	5/2	..	_____
84 Nashville Warbler	..	5/10	5/4	5/9	5/19	5/2	5/1	_____
96 Tennessee Warbler	..	5/11	..	..	..	..	..	_____
90 Parula Warbler	..	5/11	..	..	5/12	5/3	5/9	_____
77 Cape May Warbler	..	5/13	5/1	5/2	5/5	5/2	5/9	_____

68	Yellow Warbler	5/11	4/30	4/29	4/30	5/6	5/2	4/29	_____
83	Black-throated Blue Warbler	5/11	5/11	5/4	5/8	5/5	5/2	5/5	_____
38	Myrtle Warbler	4/14	4/16	4/14	4/19	4/14	4/17	4/8	_____
66	Magnolia Warbler	4/26	5/11	4/27	5/8	5/6	4/21	5/5	_____
100	Cerulean Warbler	..	..	..	5/12	..	..	..	_____
98	Chestnut-sided Warbler	5/21	5/10	5/30	5/3	5/7	5/2	5/9	_____
89	Bay-breasted Warbler	..	5/20	5/4	5/11	5/15	4/22	5/9	_____
102	Black-poll Warbler	..	5/19	5/10	5/13	5/16	5/12	5/6	_____
73	Blackburnian Warbler	5/12	5/10	5/3	4/28	5/10	4/28	4/30	_____
56	Black-throated Green Warbler	4/18	5/17	4/23	4/27	4/26	4/28	5/2	_____
53	Pine Warbler	..	..	..	4/26	4/30	4/21	4/22	_____
49	Palm Warbler	..	4/29	4/20	4/27	4/26	4/21	4/22	_____
74	Ovenbird	5/11	5/1	5/3	5/5	5/5	5/1	5/6	_____
51	Grinnell Water-Thrush	..	4/30	4/28	4/23	4/17	4/21	4/29	_____
62	Louisiana Water-Thrush	..	..	..	..	..	4/27	5/5	_____
113	Connecticut Warbler	..	..	..	..	..	..	5/20	_____
97	Mourning Warbler	..	5/20	5/13	5/6	5/16	5/2	5/11	_____
65	Western Yellow-throat	4/26	5/1	4/28	5/5	5/5	5/2	5/1	_____
82	Yellow-breasted Chat	..	..	..	..	..	5/2	5/11	_____
101	Hooded Warbler	..	..	..	..	..	..	5/12	_____
105	Wilson Black-cap Warbler	5/21	5/6	5/22	5/10	5/16	5/3	5/17	_____
104	Canadian Warbler	..	5/19	5/16	5/12	5/16	5/6	5/12	_____
76	American Redstart	5/11	5/1	4/28	5/11	5/6	5/2	5/5	_____
Family									
TROGLODYTIDAE									
58	Mockingbird	..	..	..	..	4/29	..	..	_____
52	Catbird	4/26	5/1	4/28	4/30	4/28	3/26	5/5	_____
45	Brown Thrasher	4/26	4/24	4/27	4/16	4/24	4/22	4/12	_____
46	House Wren	4/14	5/4	4/30	..	4/21	4/23	4/12	_____

36 Winter Wren	4/8	4/25	4/30	..	3/22	4/19	4/8	_____
88 Long-billed Marsh Wren	..	4/14	..	4/12	..	..	3/29	_____
Family CERTHIIDAE								
19 Brown Creeper	4/7	4/7	4/6	4/2	4/4	3/21	3/18	_____
Family PARIDAE								
21 White-breasted Nuthatch	4/7	3/29	4/11	4/2	3/18	3/31	4/2	_____
54 Red-breasted Nuthatch	..	..	..	5/3	..	4/22	4/28	_____
20 Chickadee	..	4/1	..	3/19	4/13	..	..	_____
Family SYLVIIDAE								
17 Golden-crowned Kinglet	4/7	3/14	4/11	4/6	4/5	3/28	3/19	_____
28 Ruby-crowned Kinglet	4/14	4/1	4/13	4/7	4/17	4/13	3/19	_____
47 Blue-gray Gnatcatcher	..	4/25	4/22	4/18	4/28	4/22	4/25	_____
Family TURDIDAE								
64 Wood Thrush	4/18	5/3	5/1	..	5/4	4/29	5/12	_____
42 Wilson Thrush	5/11	4/10	4/12	4/7	4/11	4/26	4/8	_____
110 Gray-cheeked Thrush	..	..	..	..	..	5/21	5/11	_____
60 Olive-backed Thrush	..	5/1	5/4	5/8	4/30	4/22	4/29	_____
26 Hermit Thrush	..	3/31	4/11	4/7	4/5	4/13	3/19	_____
3 Robin	3/28	3/9	3/4	3/24	3/14	3/10	3/7	_____
7 Bluebird	3/28	3/8	4/4	3/31	3/17	3/7	3/7	_____

WATER AND SHORE BIRDS.

General Hints.

The [Loon](#) and [Grebes](#) are common representatives of the order of Diving Birds. Their legs are set far back on the body, making it difficult for them to walk at all.

[Grebes](#) look like small, tailless ducks. They have long, slender necks, short wings, smooth glossy plumage and flat, lobed feet. They rarely leave the water and can dive or sink out of sight instantly when disturbed, swimming to a distance with only the tip of the bill out of water.

[Loons](#) are very large, with flat, heavy bodies, short tails and long, tapering bills. They are never crested, but are conspicuously marked, and are equally expert with the grebes in diving and sinking.

[Gulls](#) and [Terns](#) belong to the order of the *Long-winged Swimmers*, but are better described by the name of *Skimmers*.

[Terns](#) are much slenderer and usually smaller than gulls, have very pointed bills and wings, and forked tails. They rarely if ever swim, but skim swallow-like over the water, bill downward, plunging into the water for their prey.

[Gulls](#) are plumper than terns, with heavier bills and tails usually even. They do not dive or plunge suddenly into the water, but fly and soar or float about on the surface, sitting well up out of the water.

Ducks, Geese and Swans have webbed feet and short legs, and in walking ¹⁴² carry themselves almost horizontal. As a rule they have very strong wings, enabling them to fly long distances at great speed. Our ducks are most easily remembered in three groups:—

1. **The fish-eating Mergansers**, whose plumage is largely black and white and which have saw-edged bills and, usually, conspicuous crests.
2. **The Sea Ducks**, which are conspicuously marked but plainly colored, frequent open water or the sea coast, diving, often to great depths, for their food. Descriptions are given of the [Lesser Scaup](#), [Redhead](#), *Canvas-back*, [American Golden-eye](#), [Oldsquaw](#) and [Buffle-head](#).
3. **The River Ducks**, which have broad, rounded bills, are much variegated in color and markings, and have a peculiar habit of feeding head downwards, as though standing on their heads with the body tipped up. The [Teals](#), [Mallard](#), [Wood Duck](#) and [Pintail](#) belong to this group. The females of these ducks, though differing noticeably in size, may easily be confused.

[Geese](#) are larger than ducks, feed mostly on land and are usually seen during migration flying at a height in regular ranks after a leader.

[Hérons](#) and [Bitterns](#) belong among the *Wading Birds*. They have long, pointed bills, long legs, loose, baggy plumage, broad wings and long necks, curved s shaped in flight.

[Hérons](#) are tall, crested birds, with very long legs, bills and necks and small tails. Their notes are harsh and squawking. Large numbers sometimes nest together.

[Bitterns](#) have shorter legs, necks and bills than herons and are very rarely crested. They utter hoarse, resounding calls, and are ordinarily solitary in habit.

Rails and Coots are known as *Marsh Birds*, although the latter are fine swimmers. [43]

Rails as a rule are smaller than bitterns and frequent grassy marshes where they steal stealthily about, effectively concealed by their dull streaky plumage. Short turned-up tails, short wings and long legs are noticeable characteristics of these rapid runners.

Coots are ducklike in appearance and smoothly plumaged. Lobed feet enable them to swim easily. Their bills spread out at the base in the form of a plate or shield which extends up on to the forehead.

Plover, **Sandpipers** and the **Woodcock** are strikingly different representatives of the great order of *Shore Birds*. The plover family have stout, short bills, while the sandpipers and woodcock belong to a family which have soft probing bills of greatly varying length, for obtaining their food in soft mud.

Sandpipers have pointed wings, are dull-colored, and are usually found slipping gracefully along the water's edge in search of food. They skim rapidly over the water on outspread wings, and their clear, reed-like notes are distinctly musical.

Plovers are small and plump, with long pointed wings, short necks and rather flat heads. They run and fly very rapidly, generally in flocks piping loudly but sweetly. They frequent the inland as well as the shore.

Shore and Water Birds are often extremely shy and they can detect the presence of an observer at a distance. The caution is therefore emphasized to approach them slowly and quietly.

[44]

Particular Hints.

Note:—Owing to a lack of sufficient data the following *Shore and Water Birds* are arranged according to the American Ornithologist's Union order

instead of their average first appearance. They have all been seen, however, in Lincoln Park by the authors.

115. HORNED GREBE. 14 in.

(Colymbus auritus. 3.)

Brownish, sooty black, extending in narrow line up *back of neck*; rest of neck, sides and upper breast rich chestnut-brown; *silky* white below; head and chin deep black, set off by buffy “horns” or crests, which slant abruptly backwards from the eyes; black bill, tipped with yellow; eye, red. WINTER PLUMAGE: Black parts sooty; brown replaced by white; grayer below; crests and ruff less conspicuous. [Pied-billed Grebe.](#)

116. PIED-BILLED GREBE. 13-1/2 in.

(Podilymbus podiceps. 6.)

Brownish-black, showing gray on head and neck; belly, dusky white, otherwise brownish below; black throat-patch and a conspicuous black band across bill. WINTER PLUMAGE: Throat whitish; browner below; no band on bill. [Horned Grebe.](#)

117. LOON. 33 in.

(Gavia imber. 7.)

Black, showing greenish on head and neck, spotted with square or oval patches of white on back and wings and striped with white in front of wings; white below; two conspicuous white-striped bars across the black neck give the effect of a *broad black collar*; tail, very short; eye, red; large black bill, long and pointed; dives and sinks like the [Grebes](#); note, an eerie, prolonged cry.

[45]

118. AMERICAN HERRING GULL. 24 in.

(Larus argentatus smithsonianus. 51a.)

White; wings and back, pearly blue-gray; bill yellow, showing a small vermillion spot on either side; the longest wing feathers are partly black,

marked and *tipped with white*; the closed wing therefore shows black towards the end, marked with a regular line of white spots and tipped with white at the extreme point; eyelids, bright yellow; looks large and heavy in flight. WINTER PLUMAGE: Streaked on head and neck with gray. IMMATURE PLUMAGE: Dark and much streaked with brownish; bill darker. [Ring-billed Gull.](#)

119. RING-BILLED GULL. 19 in.

(*Larus delawarensis*. 54.)

Similar to the *American Herring Gull*, but *smaller*; bill, greenish-yellow, bright at tip, *banded with black around the middle*; wing tipped at extreme point with black; feet greenish-yellow; eyelids vermillion-red. [American Herring Gull.](#)

120. BONAPARTE GULL. 13 in.

(*Larus philadelphia*. 60.)

Back and wings, light pearl-gray; head and throat, dark slate color; back of head, neck, underparts and square tail, white; wings tipped, and *bordered narrowly on the outside edge with black*; feet and legs, red; bill black. WINTER PLUMAGE: Hood, grayish white. Large flocks seen in Lincoln Park. [Common Tern.](#)

121. COMMON TERN. 15 in.

(*Sterna hirundo*. 70.)

Pearly gray back and wings, whiter on rump; *top of head, shining black*; pure white on throat; dusky-white below; tail forked, *outer edge darkest*; bill long and red, blackening towards tip; feet reddish. WINTER PLUMAGE: Front of head and under-parts, white; bill nearly black. [Bonaparte Gull.](#)

[46

122. AMERICAN MERGANSER. 25 in.

(*Merganser americanus*. 129.)

Black; rump and tail, ashy gray; head, throat and upper part of neck, greenish black; wing, *largely white*, edged with black and crossed by a black bar; white below, extending around the neck in a narrow collar; long red bill tipped with black; eye, red; shows salmon tinge below in flight. FEMALE: Ashy-gray, with sharply defined brown head and neck and a pale salmon or brownish tinge across upper breast; *throat, white*; wings largely dark with a white patch; small crest on back of neck; feet orange; eye yellow; rare; “pursues and catches food under water.” [American Golden-eye. Red-breasted Merganser.](#)

123. RED-BREASTED MERGANSER. 22 in.

(Merganser serrator. 130.)

Long ragged crest; head and neck black; broad white collar; broad cinnamon band streaked with black across the breast, otherwise white below, showing salmon tinge in flight; wing largely white, edged and barred twice with black; eye and bill, red. FEMALE: Back and wings dark gray turning to brown on head and neck; much paler on sides of neck and throat and gray across breast; crest, less prominent; white wing-patch; common. [American Merganser.](#)

124. HOODED MERGANSER. 18 in.

(Lophodytes cucullatus. 131.)

Black, including throat and neck; large, circular crest, white, bordered with black; white below running up in front of the wings in two points; sides brownish, finely lined with black; white wing-patch, crossed by black bar; also lengthwise white streaks on end of wings; *short* black bill; eye yellow. FEMALE: Grayish-brown; throat white; crest small; sides unmarked. [Buffle-head.](#)

[47

125. MALLARD. 23 in.

(Anas boschas. 132.)

Head, throat and neck glossy green; *narrow* white collar; breast, rich brown; back, dark brownish; underparts, silver-gray; tail white, set off by black feathers which curl up from either side of the black rump; wing-patch

purple, bordered on either side with a black and then a white bar. FEMALE: Buffy-brown and black; much streaked and speckled; lighter on throat; shows wing-patch as in male; common.

126. GREEN-WINGED TEAL. 14-1/2 in.

(*Nettion carolinensis*. 139.)

Gray, finely lined on sides and shoulders with black; *white bar in front of wing*; head, including throat, brown with a broad green stripe from eye to back of head, ending in a small tuft; wings, gray-brown with brilliant green and black wing-patch, bordered by buffy bars; breast very pale reddish-brown, speckled with round black spots; buffy patches on sides of tail; white belly. FEMALE: Mottled brown; top of head and back of neck dark brown; sides of head and neck buff-colored and finely streaked; throat buff, unmarked; no green on head nor white bar in front of wings; wing-patch as in male, but smaller. [Blue-winged Teal.](#)

127. BLUE-WINGED TEAL. 15 in.

(*Querquedula discors*. 140.)

Back and underparts thickly mottled brownish and black, lightest below; head slaty, showing purplish gloss; a conspicuous crescent-shaped white stripe in front of eye; *shows blue* on bend of wing, followed by a white bar and a bright green wing-patch; white patch on sides of tail; bill black. FEMALE: Dusky-brown; black on top of head; streaked and whitish on neck and sides of head; throat and about base of bill, *white*; back and underparts, mottled and spotted; wing shows blue but no green; white bars on head are wanting. [Green-winged Teal.](#)

[48

128. PINTAIL. 27 in.

(*Dafila acuta*. 143.)

Head, including throat, brown, darkest on top; long, swan-like neck, black above, finely waved white and dusky on back and sides; *long middle tail feathers*, black; white below with a conspicuous curved white stripe running up sides of neck to head and ending in a point; bronzy patch on wings.

FEMALE: Tail much shorter but pointed; dusky, everywhere streaked; no white stripe on neck; whitish wing-bars; smaller than male. [Old-squaw](#).

129. WOOD DUCK. 18 in.

(Aix sponsa. 144.)

Highly variegated; long, smooth, glossy crest, showing green and purple iridescence and marked by two very narrow white parallel lines, curving from bill and behind eye almost to end of crest; throat white, extending irregularly in two stripes, one up behind eye, the other nearly around neck; breast, glossy brown, spotted with white and set off on either side by a conspicuous white stripe bordered with black; wings highly iridescent, marked by green patch bordered with white. FEMALE: Much duller and less conspicuously marked; head brownish-gray, slightly crested showing greenish tints; throat and line extending from it around base of bill, with space about eye, white.

130. REDHEAD. 20 in.

(Aythya americana. 146.)

Puffy head and *upper* part of neck and throat, bright reddish brown; breast and upper part of back with rump, *black*; belly white; middle back and *sides* evenly and finely waved black and white, *which shows gray at a distance*. FEMALE: Grayish-brown, almost white on throat; brown on head, light brown on neck. **Canvas-back.**

[49]

131. LESSER SCAUP DUCK. 16 in.

(Aythya affinis. 149.)

Head, neck and breast black, showing *purple* reflections on head; back black and white, very narrowly barred; upper parts of wings dark, finely mottled with white; lower part of wing brownish-black, showing a small white patch; lower breast and belly white; bill, bluish, tipped with black. FEMALE: Black replaced by brown, lighter on head; region around bill white. [Female Redhead](#), easily confused with female Lesser Scaup.

132. AMERICAN GOLDEN-EYE. 19 in.

(Clangula americana. 151.)

Black, glossed with green; white below, extending around the neck in a collar; head bunchy, slightly crested and marked *below eye* and just back of eye by a nearly oval *white spot*; wings largely white; short bill. FEMALE: Brown, snuff-colored on head; white collar; white wing patch; band of bluish gray across breast and down sides. [American Merganser.](#)

133. BUFFLE-HEAD. 15 in.

(Charitonetta albeola. 153.)

Head, throat and upper neck, iridescent black; conspicuous black crest broadly banded with white over head from eye to eye; back black, ashy on tail; wings black with a very large white patch; *collar and entire under parts, white*. FEMALE: Smaller; brownish, lighter below with no crest and only a white patch on cheeks and a small white wing patch. [Hooded Merganser.](#)

134. OLD-SQUAW. 21 in.

(Harelda hiemalis. 154.)

WINTER PLUMAGE: Back, long tail-feathers and breast with conspicuous *patch on sides of neck*, black; throat and upper breast, top of head, neck and belly, white; cheeks grayish-brown; wings black and gray; bill crossed by yellowish band. FEMALE: Dark brown, lighter on head, grayish on tail; throat, breast and region about the eye, gray-white; below white, brown on lower part of throat; no long tail-feathers; smaller than male. SUMMER PLUMAGE: Head and neck, dusky black; grayish-white on sides of head, and female shows more brown. [Pintail.](#)

[50]

135. CANADA GOOSE. 40 in.

(Branta canadensis. 172.)

Head and neck, black with white throat-patch extending up to and just behind eye; brownish-gray marked with lighter, darkest on back; rump and

tail, black above and white below; fly after a leader in harrow-shaped ranks, often high up, crying, “honk, honk, honk.”

136. AMERICAN BITTERN. 30 in.

(*Botaurus lentiginosus*. 190.)

Streaky tawny- and dark-brown; tawny-yellowish below, mixed with white and streaked with dark brown; conspicuous black streak on sides of neck; short brown tail; breast feathers loose and baggy; yellow eye; call, loud resounding squawk or booming notes which have given it the common name of “stake-driver.”

137. LEAST BITTERN. 13 in.

(*Ardetta exilis*. 191.)

Crown, with back, tail and upper half of wings, shining greenish-black; rest of wing shows buff and bright chestnut; throat and baggy breast, whitish, shading into yellowish-brown on sides of neck, and chestnut on back of neck; buffy white below; tufts of dark feathers on either side of breast; yellow eye. [Green Heron.](#)

138. GREEN HERON. 17 in.

(*Butorides virescens*. 201.)

Wings and back dark green, latter tinged with blue-gray; crown, greenish-black; neck and breast, rich reddish brown; throat and line down middle of neck to breast whitish, ending in light streaks on breast; the smallest heron; carries neck curved in flight; squawking note, “scow.” [Least Bittern.](#)

[51

139. SORA. 8-1/2 in.

(*Porzana carolina*. 214.)

Olive-brown with lengthwise markings of black and some white; short tail turned up, showing buffy white underside; sides *barred* posteriorly with white; front of head and *throat, black*; slaty line over eye; sides of neck and breast, grayish slate; bill *short and yellowish*; legs long. YOUNG: No black;

throat whitish and brown on breast; very stealthy; skulks and crouches in grass.

140. AMERICAN COOT. 15 in.

(Fulica americana. 221.)

Smooth blackish slate color, lighter below and black on head and neck; white bill shows *dark marks near the tip*; turned up tail; white underneath; wing edged with white; eye red; young show white below and no marks on bill; toes with scalloped edges. May be mistaken for a small duck.

141. AMERICAN WOODCOCK. 10-1/2 in.

(Philohela minor. 228.)

Black, mixed black and rusty slate; below reddish-brown, no markings; *large* head with short neck and *very long straight bill*; eye set *high up and far back*; crown barred crosswise with black and rusty; short tail; burrows in soft mud for earthworms; nocturnal.

142. SEMI-PALMATED SANDPIPER. 6 in.

(Ereunetes pusillus. 246.)

Grayish-brown, marked with black and buffy; rump very dark; tail tapering; underparts *pure white*, slightly tinged and streaked across breast; white line over eye and dusky line beneath; note, “peep-peep.” [Other Sandpipers.](#)

[52

143. SOLITARY SANDPIPER. 8-1/2 in.

(Helodromas solitarius. 256.)

Back, dusky olive-brown, *finely spotted with white*; crown and back of neck showing dark streaks; white below, throat unmarked; sides of head and neck with breast slightly buffy and *distinctly streaked*; sides lightly barred; wings dark brown with one narrow white wing-bar; middle of tail very dark; outer feathers *white, barred with black*; note, a soft whistle. [Spotted Sandpiper.](#)

144. SPOTTED SANDPIPER. 7-1/2 in.

(Actitis macularia. 263.)

Greenish-ash marked lightly with black; *long white line over eye*; pure white below everywhere with dark round spots; wings brownish, marked broadly with white bar; flies close to the water with wings full-spread, showing white wing-bar; note, penetrating “pee-weet, weet;” walks with tilting motion. [Solitary Sandpiper.](#)

145. KILLDEER. 10 in.

(Oxyechus vociferus. 273.)

Gray-brown; rump rusty-brown; tail rather long; white below; white collar followed by a black collar, and a *black band across breast*; forehead, line over eye and wing-bar, white; bill black; runs very swiftly; note, loud and persistent “kill-dee” often heard high overhead.

[53]

Explanation of Chart.

The “height of the migration” usually comes during the second or third week in May. This chart makes a comparison of the observations of six years graphic.


For instance, the largest number of different species seen on any one morning in 1903 was 68 (May 12), while in 1901 it was 48 (May 16).

The sudden rise or fall in the migration may also be seen at a glance. May 9, 1899, for example, only 16 different species were observed and May 10 the record was 40. Similarly, in 1901, the number of species seen May 17 dropped from 45 to 15 in three days.

[54]

**A chart showing the number of different kinds of birds seen in *LINCOLN PARK* during the
*HEIGHT OF THE MIGRATION***

*A chart showing the number of different kinds of
birds seen in LINCOLN PARK during the*



[55]

Note to Key.

In response to many demands, the following simple field-key has been devised, to the end that the beginner may be aided in learning to observe correctly those points about a bird that appear most conspicuous in the field or which are particularly distinctive.

Technical analyses and measurements have not been employed, but instead, the birds are roughly placed in three groups according to size, the Robin and English Sparrow marking respectively the 10-inch and 6-inch lines of division.

The first part of the key deals with *general* differences in color and markings, while the second seeks to emphasize the most noteworthy *special* points which distinguish the different species. Incidentally a few striking peculiarities of bill, wings and tail have been included.

It will readily be seen that a key embracing so little cannot do more than reduce wild guessing to a few reasonable chances in identifying a strange bird. Some practice is of course necessary to enable the student to use the key readily. Its helpfulness will largely depend upon accuracy of observation and a careful application of the points which it suggests.

Shore and water birds have been omitted—first, in order to keep the key as simple as possible; second, because quite full descriptions of these birds are given in the “Particular Hints;” and, third, for the reason that beginners, as a rule, meet with far better success by becoming familiar with the common land birds before attempting the study of water and shore birds.

EXPLANATIONS.

1. The heading, “Dull Colors,” includes all shades of gray or olive, very dull dark-brown and black not showing conspicuous iridescence. Birds in this group are very rarely streaked or spotted, never barred and seldom show any trace of bright colors. 156
2. To make the key compact and to avoid referring to the index the birds are indicated by their respective numbers as given under “Particular Hints.”
3. Females differing greatly *in coloration* from the males are denoted by heavy figures. Differences in markings are taken into account only when the female is particularly obscure or liable to be mistaken for another species.
4. Every bird appears once under the first six general headings, and a few twice, some of which come under apparently contradictory headings in order to cover incomplete as well as complete observations. For example, the [Robin](#) has a streaked throat and the [Fox Sparrow](#) an indistinctly streaked back, both inconspicuous points, often overlooked in the field, yet useful if complete observations have been made. Again, the [Yellow-bellied Flycatcher](#), though brighter colored than most of its kind, would scarcely be

described as having either brilliant or very dull plumage, and is therefore difficult to tabulate exclusively in one place.

HOW TO USE THE KEY.

Two illustrations will most briefly indicate the easiest method of using the key—

Observation: A dull-colored bird of medium size, nearer six inches than ten, much streaked all over, showing nothing distinctive unless a kind of spot on the breast. Turning to the key, pass by “*Bright Colors*,” “*Iridescent*” and “*Dull Colors*,” choosing “*Streaked, Barred or Spotted*.” Not being sure of the size, it may be necessary to try all the birds between “6 and 10 inches,” as well as those “less than 6 inches.” Before going to so much trouble, however, pass on to “*Under Parts*” and see whether an easier approach can be made there. The sub-heading “Spot or patch on throat, sides, breast or sides of neck” looks comprehensive enough to apply, while in this case it contains fewer examples. Comparing the numbers there given with those under the first heading selected, it is found that only [10](#) and [23](#) appear in both places. On looking up [10](#) and [23](#) it is found that they refer to the [Song Sparrow](#) and [Sapsucker](#), two species so widely different in coloration, ¹⁵⁷ habits and movements that it ought not to be necessary to study the key further in order to be sure of the bird in question—namely, the [Song Sparrow](#).

Observation: A bird in flight, pure white below, without doubt over 10 inches on account of its long tail. Not having seen the general color with any certainty, but being fairly sure of the white breast and long tail, try “*Under Parts*” first. Only two birds over 10 inches are given under the sub-heading “Pure white or ashy,” and these are [103](#) and [109](#). Turning to “*Tail*,” both numbers appear under “very long, sometimes keeled or forked,” but only 103, the [Yellow-billed Cuckoo](#), comes under “Outer tail-feathers conspicuously white or spotted.” *If conspicuous tail spots had been seen* the key would have helped to identify the bird, but the key alone cannot make up for lack of observations.

The second illustration may give force to the suggestion that certain birds can be satisfactorily identified only after very careful observation and some further study.

BRIGHT COLORS

[58

STREAKED BARRED SPOTTED

Showing BLUE, blue-gray or bluish-ash [A1](#)

Showing RED, chestnut, light or reddish-brown [A2](#)

Showing YELLOW, orange or buff [A3](#)

Showing conspicuous BLACK markings [A4](#)

NEVER STREAKED BARRED SPOTTED

Showing BLUE, blue-gray or bluish-ash [A5](#)

Showing RED, chestnut, light or reddish-brown [A6](#)

Showing YELLOW, orange or buff [A7](#)

Showing much BLACK [A8](#)

IRIDESCENT

Showing iridescent black, blue, green or brownish [B1](#)

DULL COLORS

FEW MARKINGS

Sharply defined [C1](#)

Obscure [C2](#)

NO MARKINGS

[D1](#)

STREAKED BARRED OR SPOTTED ON BREAST BACK OR BOTH

ABOVE AND BELOW, rarely bright colors [E1](#)

PLAIN ABOVE

Heavily streaked or spotted below [E2](#)

Lightly streaked or spotted below [E3](#)

PLAIN BELOW, rarely with one conspicuous spot on breast [E4](#)

HIGHLY VARIEGATED [E5](#)

HOOD

[F1](#)

UNDER PARTS

NO MARKINGS

Pure white or ashy [G1](#)

Dull colors, indistinctly shaded [G2](#)

FEW, IF ANY, MARKINGS

Distinctive red, yellow or brown BREAST [G3](#)

BELLY white or yellow, sharply contrasting with breast [G4](#)

DISTINCTIVE MARKINGS

A line of streaks down sides or across breast	<u>G5</u>
Spotted, finely streaked, or throat and middle of belly plain	<u>G6</u>
Bib, collar, or band across breast	<u>G7</u>
Spot or patch on throat, sides, breast, or sides of neck	<u>G8</u>
Noticeably tinged with red, yellow, buff or brownish	<u>G9</u>
BILL	
Hooked, or noticeably long and sometimes curved	<u>H1</u>
Large and stout, or noticeably short and thick	<u>H2</u>
Used for hammering	<u>H3</u>
HEAD	
TOP	
CRESTED, or red crown-patch displayed at will	<u>I1</u>
Crown conspicuously STRIPED	<u>I2</u>
Distinct CAP, no line over eye	<u>I3</u>
Distinct CAP or crown-patch, with line over, through or back of eye	<u>I4</u>
FOREHEAD showing black, sometimes with bar through eye	<u>I5</u>
SIDES	
Conspicuous line over or through eye	<u>I6</u>
Inconspicuous but distinctive line over eye	<u>I7</u>
Conspicuous eye-ring, or black or yellow mask	<u>I8</u>
Distinctively marked or colored on cheeks or sides of head	<u>I9</u>
WINGS	
Long and pointed	<u>J1</u>
One or two conspicuous WING-BARS, white or yellowish	<u>J2</u>
Spots, patches or bright markings	<u>J3</u>
BACK	
STREAKED	
Entirely streaked or mottled, no distinct cap	<u>K1</u>
Rump plain, cap or striped crown	<u>K2</u>
Gray or ashy about head or neck	<u>K3</u>
BARRED AND SPOTTED	
Barred, including wings	<u>K4</u>
Black and white, wings barred or spotted	<u>K5</u>
No MARKINGS, back and wings alike	<u>K6</u>
DISTINCTIVE MARKINGS OR COLORS	
Rump patch, white, yellow or brown	<u>K7</u>

	On back of neck, between shoulders, or middle of back	K8
TAIL		
COLOR		
	Outer feathers conspicuously white or spotted	L1
	Reddish-brown, or tipped with white or yellow	L2
MARKINGS:	Barred or banded with black or white	L3
SIZE		
	Very long, sometimes keeled or forked	L4
	Very short	L5
	Narrow and pointed	L6
MOVEMENTS:	Used for bracing, or, jerking or tilting motions.	L7

	10 Inches or more	Between 6 and 10 Inches	Less than 6 Inches
BRIGHT COLORS			
A1	1 , 22 .	...	38 , 66 , 100 , 104 .
A2	3 , 43 , 45 .	13 , 40 , 80 .	76 , 89 , 98 .
A3	5 .	112 .	49 , 56 , 66 , 68 , 73 , 77 , 104 .
A4	1 , 5 , 43 .	112 .	38 , 56 , 66 , 73 , 76 , 77 , 89 , 98 , 104 .
A5	...	7 , 44 .	21 , 47 , 54 , 83 , 84 , 86 , 90 , 93 , 94 , 95 , 97 , 113 .
A6	3 .	7 , 15 , 27 , 31 , 41 , 57 , 71 , 78 , ... 87 .	
A7	111 .	27 , 41 , 61 , 70 , 82 , 92 .	53 , 65 , 67 , 79 , 84 , 90 , 93 , 95 , 97 , 101 , 105 , 113 .
A8	111 .	15 , 27 , 57 , 70 , 78 , 87 , 92 .	67 , 83 , 101 .
IRIDESCENT			
B1	9 , 35 .	11 , 24 , 30 , 44 , 81 .	107 .
DULL COLORS			
C1	35 , 58 .	2 , 4 , 52 , 69 , 87 .	17 , 20 , 28 , 72 , 83 .
C2	103 , 109 , 111 .	11 , 14 , 24 , 31 , 75 , 78 , 81 , 91 .	28 , 59 , 63 , 67 , 85 , 94 , 96
NO MARKINGS			

D1	6 .	11 , 14 , 24 , 71 , 81 .	59 , 65 , 85 , 94 .
STREAKED BARRED OR SPOTTED ON BREAST BACK OR BOTH			
E1	18 , 33 , 43 .	10 , 13 , 23 , 27 , 37 , 40 , 80 , 92 , 99 , 108 .	8 , 36 , 46 , 50 , 55 , 77 , 102 , 106 .
E2	43 , 45 .	13 , 26 , 60 , 62 , 64 , 74 , 110 .	51 , 104 .
E3	33 .	42 , 81 .	49 , 68 , 104 .
E4	5 .	12 , 16 , 29 , 32 , 48 , 112 , 114 .	19 , 25 , 34 , 39 , 88 .
E5	...	80 .	38 , 56 , 66 , 73 , 76 , 77 , 89 , 98 , 100 .
HOOD			
F1	111 .	11 , 57 , 70 , 87 .	95 , 101 .
UNDER PARTS			
G1	103 , 109 .	16 , 29 , 30 , 48 , 69 , 75 .	19 , 21 , 34 , 85 .
G2	1 , 58 .	2 , 14 , 24 , 70 , 87 , 91 .	17 , 20 , 25 , 28 , 39 , 47 , 63 , 88 , 96 .
G3	3 , 5 .	7 , 44 , 82 , 112 .	53 , 65 , 79 , 84 , 95 , 101 , 105 .
G4	...	4 , 15 , 61 , 82 .	79 , 83 , 97 , 113 .
G5	...	23 .	38 , 66 , 73 , 98 , 100 , 102 , 104 .
G6	43 .	62 , 64 .	50 .
G7	22 .	4 , 15 , 99 , 108 .	56 , 83 , 89 , 90 , 100 , 101 , 106 .
G8	5 , 18 .	10 , 12 , 23 , 32 , 80 , 112 .	76 , 93 , 97 , 107 .
G9	...	23 , 40 , 60 , 114 .	25 , 36 , 49 , 53 , 54 , 72 , 86 , 106 .
BILL			
H1	33 , 43 , 45 , 103 , 109 .	2 .	19 , 88 , 107
H2	...	11 , 31 , 40 , 80 .	67 .
H3	18 .	16 , 23 , 29 , 57 , 114 .	...
HEAD			
I1	1 , 22 .	31 , 41 , 61 , 69 .	28 .
I2	...	32 , 48 , 74 .	17 , 55 .
I3	...	52 , 114 .	20 , 21 , 67 , 89 , 102 , 105 .
I4	...	12 .	25 , 34 , 38 , 49 , 54 , 93 , 98 .

I5 ...	31 , 41 .	39 , 47 , 65 , 89 .
I6 5 .	2 , 16 , 29 , 32 , 62 , 82 , 112 .	47 , 51 , 53 , 56 , 66 , 77 , 79 , 88 .
I7 111 .	24 , 75 .	8 , 96 , 100 .
I8 ...	60 .	20 , 65 , 72 , 79 , 84 , 86 , 101 , 113 .
I9 35 , 43 .	110 , 114 .	20 , 49 , 50 , 73 , 77 , 89 , 93 , 100 , 102 .
WINGS		
J1 ...	30 , 44 , 81 , 99 , 108 .	59 .
J2 45 .	12 , 48 , 70 , 87 , 91 .	19 , 28 , 53 , 56 , 63 , 67 , 72 , 79 , 86 , 89 , 90 , 98 , 100 , 102 .
J3 58 , 111 .	2 , 15 , 27 , 57 , 80 , 108 .	66 , 76 , 83 , 93 .
BACK		
K1 5 .	10 , 37 , 40 , 80 , 92 , 99 , 108 .	8 , 50 , 55 , 106 .
K2 ...	32 .	19 , 54 , 58 , 73 , 77 , 102 .
K3 ...	12 , 48 , 112 .	25 , 39 .
K4 18 , 43 .	114 .	36 , 46 .
K5 ...	16 , 23 , 29 .	...
K6 3 , 6 , 9 , 35 , 109 .	4 , 7 , 11 , 14 , 26 , 42 , 60 , 62 , 64 , 74 , 75 , 81 , 82 , 110 .	20 , 51 , 54 , 59 , 65 , 84 , 85 , 96 , 97 , 101 , 104 , 105 , 113 .
K7 18 , 33 .	80 .	38 , 66 , 77 , 88 .
K8 18 .	16 , 23 , 29 , 114 .	50 , 88 , 90 .
TAIL		
L1 3 , 5 , 22 , 58 , 103 .	2 , 4 , 15 , 29 , 37 .	38 , 47 , 49 , 53 , 56 , 76 , 89 , 100 .
L2 1 , 35 .	13 , 26 , 41 , 61 , 69 .	...
L3 1 , 33 , 43 .	16 , 23 , 114 .	36 , 46 , 66 , 88 .
L4 9 , 35 , 45 , 103 , 109 .	44 .	19 , 47 .
L5 ...	...	21 , 36 , 54 , 59 , 107 .
L6 35 .	...	8 , 50 .
L7 18 .	16 , 23 , 29 , 57 , 62 , 74 , 114 .	19 , 49 , 51 .

Glossary.

Barred—Marked with transverse lines.

Bib—Whole throat, with upper breast of same color, sharply defined.

Cap—Entire top of head defined by an oval patch of distinct and contrasting color.

Crest—A tuft of feathers carried erect on top or back of head.

Crown-patch—Smaller and less sharply defined than cap.

Hood—Whole head and neck all around of same color.

Mask—Forehead, together with broad stripe through eye, of same color.

Mottled—Colors and markings blended in no distinct patterns.

Patch—An oddly-shaped and conspicuous mark, never round.

Rump—The extreme lower portion of the back next the tail (easily seen in flight).

Spotted—Marked with nearly round spots or one irregular spot.

Streaked—Marked with *longitudinal* lines or streaks.

Variegated—Showing a variety of colors and markings.

Wing-Bar—A line, usually white, running obliquely across wing.

[63]

Supplementary List.

NOTE.—Owing to the impossibility of determining what additional species are most likely to stray into a city park, the following list has been made to include certain species known to occur in Cook County, Illinois, some of which are of common occurrence in the vicinity of Chicago but have not yet been seen in Lincoln Park by the authors:

Red-throated Loon (*Gavia lummei*. 11.)

Glaucous Gull. (*Larus glaucus*. 42.)

Franklin Gull. (*Larus franklinii*. 59.)

Forster Tern. (*Sterna forsteri*. 69.)

Black Tern. (*Hydrochelidon nigra surinamensis*. 77.)

Shoveller. (*Spatula clypeata*. 142.)

American Scaup Duck. (*Aythya marila*. 148.)

Ring-necked Duck. (*Aythya collaris*. 150.)

Great Blue Heron. (*Ardea herodias*. 194.)

Black-crowned Night Heron. (*Nycticorax nycticorax naevius*. 202.)

King Rail (*Rallus elegans*. 208.)

Virginia Rail. (*Rallus virginianus*. 212.)

Wilson Snipe. (*Gallinago delicata*. 230.)

Pectoral Sandpiper. (*Actrodramas maculata*. 239.)
 Least Sandpiper. (*Actrodramas minutilla*. 242.)
 Greater Yellow-legs. (*Totanus melanoleucus*. 254.)
 Yellow-legs. (*Totanus flavipes*. 255.)
 Bartramian Sandpiper. (*Bartramia longicauda*. 261.)
 Black-bellied Plover. (*Squatarola squatarola*. 270.)
 American Golden Plover. (*Charadrius dominicus*. 272.)
 Sharp-shinned Hawk. (*Accipiter velox*. 332.)
 Cooper Hawk. (*Accipiter cooperi*. 333.)
 Red-tailed Hawk. (*Buteo borealis*. 337.)
 Red-shouldered Hawk. (*Buteo lineatus*. 339.)
 Broad-winged Hawk. (*Buteo platypterus*. 343.)
 American Rough-legged Hawk. (*Archibuteo lagopus sancti-johannis*. 347a.)
 Pigeon Hawk. (*Falco columbarius*. 357.)
 Short-eared Owl. (*Asio accipitrinus*. 367.)
 Barred Owl. (*Syrnium varium*. 368.)
 Saw-whet Owl. (*Nyctala acadia*. 372.)
 Screech Owl. (*Megascops asio*. 373.)
 Acadian Flycatcher. (*Empidonax virescens*. 465.)
 Traill Flycatcher. (*Empidonax traillii*. 466.)
 Prairie Horned Lark. (*Otocoris alpestris praticola*. 474b.)
 Evening Grosbeak. (*Hesperiphona vespertina*. 514.)
 Redpoll. (*Acanthis linaria*. 528.)
 Snowflake. (*Passerina nivalis*. 534.)
 Lapland Longspur. (*Calcarius lapponicus*. 536.)
 Grasshopper Sparrow. (*Coturniculus savannarum passerinus*. 546.)
 Eave Swallow. (*Petrochelidon lunifrons*. 612.)
 Bank Swallow. (*Riparia riparia*. 616.)
 Northern Shrike. (*Lanius borealis*. 621.)
 White-rumped Shrike. (*Lanius ludovicianus excubitorides*. 622a.)
 Philadelphia Vireo. (*Vireo philadelphicus*. 626.)
 White-eyed Vireo. (*Vireo noveboracensis*. 631.)
 Blue-winged Warbler. (*Helminthophila pinus*. 641.)
 Orange-crowned Warbler. (*Helminthophila celata*. 646.)
 Water-Thrush. (*Seiurus noveboracensis*. 675.)
 American Pipit. (*Anthus pensilvanicus*. 697.)
 Short-billed Marsh Wren. (*Cistothorus stellaris*. 724.)
 Tufted Titmouse. (*Bæolophus bicolor*. 731.)

[64]

[65]

Index of Common Names.

A · [B](#) · [C](#) · [D](#) · E · [F](#) · [G](#) · [H](#) · I · [J](#) · [K](#) · [L](#) · [M](#) · [N](#) · [O](#) · [P](#) · Q · [R](#) · [S](#) · [T](#) · U ·
[V](#) · [W](#) · X · [Y](#) · [Z](#)

Page

[Bittern, American,](#)

50

<u>Least,</u>	50
<u>Blackbird, Red-winged,</u>	16
<u>Rusty,</u>	15
<u>Yellow-headed,</u>	31
<u>Bluebird,</u>	12
<u>Bobolink,</u>	27
<u>Buffle-head,</u>	49
<u>Bunting, Indigo,</u>	28
<u>Cardinal,</u>	16
<u>Catbird,</u>	20
<u>Chat, Yellow-breasted,</u>	26
<u>Chickadee,</u>	14
<u>Coot, American,</u>	51
<u>Cowbird,</u>	13
<u>Creeper, Black and White,</u>	21
<u>Brown,</u>	14
<u>Crow,</u>	12
<u>Cuckoo, Black-billed,</u>	30
<u>Yellow-billed,</u>	29
<u>Dickcissel,</u>	31
<u>Dove, Mourning,</u>	17
<u>Duck, Lesser Scaup,</u>	49
<u>Wood,</u>	48
<u>Finch, Purple,</u>	18
<u>Flicker,</u>	14
<u>Flycatcher, Great-crested,</u>	22
<u>Least,</u>	22
<u>Yellow-bellied,</u>	24
<u>Gnatcatcher, Blue-gray,</u>	19
<u>Golden-eye, American,</u>	49
<u>Goldfinch, American,</u>	23
<u>Goose, Canada,</u>	50
<u>Grackle, Bronzed,</u>	12
<u>Grebe, Horned,</u>	44

<u>Pied-billed,</u>	44
<u>Grosbeak, Rose-breasted,</u>	25
<u>Gull, American Herring,</u>	45
<u>Bonaparte,</u>	45
<u>Ring-billed,</u>	45
<u>Hawk, American Sparrow,</u>	18
<u>Marsh,</u>	17
<u>Heron, Green,</u>	50
<u>Hummingbird, Ruby-throated,</u>	30
<u>Jay, Blue,</u>	11
<u>Junco,</u>	12
<u>Killdeer,</u>	52
<u>Kingbird,</u>	23
<u>Kingfisher, Belted,</u>	15
<u>Kinglet, Golden-crowned,</u>	14
<u>Ruby-crowned,</u>	16
<u>Loon,</u>	44
<u>Mallard,</u>	47
<u>Martin, Purple,</u>	25
<u>Meadowlark,</u>	12
<u>Merganser, American,</u>	46
<u>Hooded,</u>	46
<u>Red-breasted,</u>	46
<u>Mockingbird,</u>	21
<u>Nighthawk,</u>	30
<u>Nuthatch, Red-breasted,</u>	20
<u>White-breasted,</u>	15
<u>Old-squaw,</u>	49
<u>Oriole, Baltimore,</u>	23
<u>Orchard,</u>	27
<u>Ovenbird,</u>	24

<u>Pewee, Wood,</u>	27
<u>Phoebe,</u>	13
<u>Pintail,</u>	48
<u>Redhead,</u>	48
<u>Redstart, American,</u>	24
<u>Robin,</u>	11
<u>Sandpiper, Semi-palmated,</u>	51
<u>Solitary,</u>	52
<u>Spotted,</u>	52
<u>Sapsucker, Yellow-bellied,</u>	15
<u>Shrike, Loggerhead,</u>	11
<u>Sora,</u>	51
<u>Sparrow, Chipping,</u>	17
<u>Field,</u>	15
<u>Fox,</u>	13
<u>Henslow,</u>	20
<u>Lincoln,</u>	30
<u>Savanna,</u>	12
<u>Song,</u>	13
<u>Swamp,</u>	18
<u>Tree,</u>	13
<u>Vesper,</u>	17
<u>White-crowned,</u>	19
<u>White-throated,</u>	16
<u>Swallow, Barn,</u>	18
<u>Tree,</u>	16
<u>Swift, Chimney,</u>	21
<u>Tanager, Scarlet,</u>	25
<u>Summer,</u>	23
<u>Teal, Blue-winged,</u>	47
<u>Green-winged,</u>	47
<u>Tern, Common,</u>	45
<u>Thrasher, Brown,</u>	19
<u>Thrush, Gray-cheeked,</u>	31
<u>Hermit,</u>	15

<u>Olive-backed,</u>	21
<u>Wilson,</u>	18
<u>Wood,</u>	22
<u>Towhee,</u>	13
<u>Vireo, Blue-headed,</u>	26
<u>Red-eyed,</u>	24
<u>Warbling,</u>	26
	[66]
<u>Yellow-throated,</u>	25
<u>Warbler, Bay-breasted,</u>	27
<u>Blackburnian,</u>	24
<u>Black-poll,</u>	29
<u>Black-throated Blue,</u>	26
<u>Black-throated Green,</u>	21
<u>Blue Golden-winged,</u>	28
<u>Canadian,</u>	30
<u>Cape May,</u>	25
<u>Cerulean,</u>	29
<u>Chestnut-sided,</u>	28
<u>Connecticut,</u>	31
<u>Hooded,</u>	29
<u>Magnolia,</u>	23
<u>Mourning,</u>	28
<u>Myrtle,</u>	17
<u>Nashville,</u>	26
<u>Palm,</u>	19
<u>Parula,</u>	27
<u>Pine,</u>	20
<u>Prothonotary,</u>	28
<u>Tennessee,</u>	28
<u>Wilson Black-cap,</u>	30
<u>Yellow,</u>	23
<u>Water-Thrush, Grinnell,</u>	20
<u>Louisiana,</u>	22
<u>Waxwing, Cedar,</u>	18
<u>Whip-poor-will,</u>	29

<u>Woodcock, American,</u>	51
<u>Woodpecker, Downy,</u>	14
<u>Hairy,</u>	16
<u>Red-bellied,</u>	31
<u>Red-headed,</u>	21
<u>Wren, House,</u>	19
<u>Long-billed Marsh,</u>	27
<u>Winter,</u>	17
<u>Yellow-throat, Western,</u>	22

Colored plates, 7-1/2 x 9-1/2 inches in size, of all the birds mentioned in 168
this little book, except numbers 50, 53, 61, 63, 105, 106, 113, 115, 122,
142 and 143, will be sent by the publishers at 2 cents each, or a portfolio of
134 pictures for \$2.00. Both book and pictures, if ordered together, \$2.25.

A. W. MUMFORD & COMPANY, Publishers
378 Wabash Avenue
Chicago, Ill.

Transcriber's notes to the Electronic Edition

- Spelling, capitalization, and punctuation of bird names was made consistent (*e.g.*, “Redhead” vs. “Red-head” vs. “Redheaded”). All spelling changes were justified by usage elsewhere in the text. Unambiguous abbreviated names (“American Sparrow Hawk” vs. “Sparrow Hawk”) were not changed.
- Note that the “Yellow Warbler” is also once called “Summer Warbler”.
- The “Canvas-back Duck” was, apparently unintentionally, omitted from the bird list, but is referenced several times in the text.
- Two tables, the “Identification Key” and “Tables of Arrival”, were reformatted for vertical scrolling.
- In the ASCII text file, an * (asterisk) indicates that the following bird name or number was printed in **bold font**.

*** END OF THE PROJECT GUTENBERG EBOOK WILD BIRDS IN
CITY PARKS ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, “Information about donations to the Project Gutenberg Literary Archive Foundation.”
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.