

THE
BOTANICAL MAGAZINE;
OR,
Flower-Garden Displayed :

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the
Open Ground, the Green-House, and the Stove, are
accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according
to the celebrated LINNÆUS; their Places of Growth,
and Times of Flowering :

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A W O R K

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as
wish to become scientifically acquainted with the Plants they cultivate.

By *WILLIAM CURTIS*,
Author of the *FLORA LONDINENSIS*.

V O L. X.

“ What nature, alas ! has denied
“ To the delicate growth of our illc,
“ Art has in a measure supplied ;
“ E'en winter is deck'd with a smile.

COWPER.

L O N D O N :

PRINTED BY STEPHEN COUCHMAN,

For W. CURTIS, N^o 3. *St. George's-Crescent*, Black-Friars-Road;
And Sold by the principal Booksellers in Great-Britain and Ireland.

M D C C C X V I.

THE
BOTANICAL MAGAZINE;
OR,
Flower-Garden Displayed:
IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the
Open Ground, the Green-House, and the Stove, are
accurately represented in their natural Colours.

TO WHICH ARE ADDED,
Their Names, Class, Order, Generic and Specific Characters, according
to the celebrated LINNÆUS; their Places of Growth,
and Times of Flowering :

TOGETHER WITH
THE MOST APPROVED METHODS OF CULTURE.

A W O R K

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as
wish to become scientifically acquainted with the Plants they cultivate.

By WILLIAM CURTIS,
Author of the FLORA LONDINENSIS.

VOL. X.

" What nature, alas ! has denied
" To the delicate growth of our idle,
" Art has in a measure supplied ;
" E'en winter is deck'd with a smile.

COWPER.

L O N D O N :
PRINTED BY STEPHEN COUCHMAN,
For W. CURTIS, N^o 3. *St. George's-Crescent*, Black-Friars-Road ;
And Sold by the principal Bookfellers in Great-Britain and Ireland.
MDCXCVI.

The Project Gutenberg eBook of The Botanical Magazine, Vol. 10

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The Botanical Magazine, Vol. 10

Author: William Curtis

Release date: May 18, 2020 [eBook #62172]

Most recently updated: October 18, 2024

Language: English

Other information and formats: www.gutenberg.org/ebooks/62172

Credits: Produced by Karin Spence, Jason Isbell and the Online Distributed Proofreading Team at <https://www.pgdp.net>
(This book was produced from images made available by the HathiTrust Digital Library.)

*** START OF THE PROJECT GUTENBERG EBOOK THE
BOTANICAL MAGAZINE, VOL. 10 ***

THE **BOTANICAL MAGAZINE;** OR, Flower-Garden Displayed:

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the Open Ground, Green-House and the Stove, are accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according to the celebrated LINNÆUS; their place of Growth, and Times of Flowering:

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A W O R K

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as wish to become scientifically acquainted with the Plants they cultivate.

By *WILLIAM CURTIS*,

Author of the FLORA LONDINENSIS.

VOL. X.

"What nature, alas! has denied
"To the delicate growth of our isle,
"Art has in a measure supplied;
"E'en Winter is deck'd with a smile."
COWPER.

LONDON

PRINTED BY STEPHEN COUCHMAN,

For W. CURTIS, N^o 3, *St. George's-Crescent*, Black-Friars-Road; And Sold by the principal Booksellers in Great-Britain and Ireland.

MDCCXCVI.

TABLE OF CONTENTS

<u>325—JUSTICIA NASUTA.</u>
<u>326—MESEMBRYANTHEMUM VIRIDIFLORUM.</u>
<u>327—CHRYSANTHEMUM INDICUM.</u>
<u>328—TRIFOLIUM INCARNATUM.</u>
<u>329—ONONIS NATRIX.</u>
<u>330—SIDA CRISTATA.</u>
<u>331—KALMIA ANGUSTIFOLIA.</u>
<u>332—OENOTHERA FRUTICOSA.</u>
<u>333—CERINTHE MAJOR.</u>
<u>334—HYPERICUM MONOGYNUM.</u>
<u>335—ONONIS ROTUNDIFOLIA.</u>
<u>336—LOTUS HIRSUTUS.</u>
<u>337—PRUNELLA GRANDIFLORA.</u>
<u>338—ALLAMANDA CATHARTICA.</u>
<u>339—ARUM TRILOBATUM.</u>
<u>340—POLYGALA HEISTERIA.</u>
<u>341—SCILLA AMÆNA.</u>
<u>342—ERICA PERSOLUTA.</u>
<u>343—ANTHOLYZA CUNONIA.</u>
<u>344—ASPALATHUS PEDUNCULATA.</u>
<u>345—POLYGALA BRACTEOLATA.</u>
<u>346—PROTEA MELLIFERA.</u>
<u>347—OENOTHERA ROSEA.</u>
<u>348—CALCEOLARIA FOTHERGILLII.</u>
<u>349—SOLANUM LACINIATUM.</u>
<u>350—ERICA VENTRICOSA.</u>

351—SAXIFRAGA MUTATA.

352—OENOTHERA PURPUREA.

353—MAHERNIA INCISA.

354—MIMULUS AURANTIACUS.

355—OENOTHERA PUMILA.

356—ERICA MASSONI.

357—BRIZA MAXIMA.

358—ERICA BACCANS.

359—CONVOLVULUS ALTHÆOIDES.

360—HIBISCUS SPECIOSUS.

INDEX—Latin Names of the Plants.

INDEX—English Names of the Plants.

INDEX—Latin Names of the Plants contained in the first Ten Volumes.

INDEX—English Names of the Plants contained in the first Ten Volumes.

INDEX—Hardy Trees contained in the first Ten Volumes.

INDEX—Hardy Shrubs contained in the first Ten Volumes.

INDEX—Hardy Herbaceous Perennial Plants contained in the first Ten Volumes.

INDEX—Annual and Biennial Plants contained in the first Ten Volumes.

INDEX—Greenhouse Plants contained in the first Ten Volumes

INDEX—Stove Plants contained in the first Ten Volumes

INDEX—Plants contained in the first Ten Volumes arranged according to the System of LINNÆUS

[325]

JUSTICIA NASUTA. DICHOTOMOUS JUSTICIA.

Class and Order.

DIANDRIA MONOGYNIA.

Generic Character.

Cor. ringens. *Caps.* 2-locularis: seminum retinaculis 2-valvis: valvis longitudinalibus unculatis. *Stamina* anthera solitaria.

Specific Character and Synonyms.

JUSTICIA *nasuta* foliis lanceolato-ovatis integerrimis pedunculis dichotomis. *Linn. Sp. Pl. Ed.* 3. p. 23. *Syst. Vegetab. ed. 14. Murr. p. 63.*

PULCOLLI *Rheed. Mal. 9. p. 135. t. 69.*



N^o. 325

This species of *Justicia*, long since figured in the *Hortus Malabaricus*, and described in the *Species Plantarum* of LINNÆUS, has been introduced to the Royal Garden at Kew, since the publication of the *Hortus Kewensis*, by

Mr. AITON: it is a stove plant, producing, almost the year through, abundance of flowers, distinguished not less for their singularity than their snowy whiteness; the following description, taken from the living plant, is fuller than that of LINNÆUS in the *Sp. Pl.*

DESCR. Stalk shrubby, somewhat angular, three feet or more in height, green, very much branched, slightly pubescent; Leaves opposite, standing on short footstalks, ovato-lanceolate, running out to a short blunt point, entire, veiny, a little downy; Peduncles from the alæ of the leaves, alternate, dichotomous, two small bracteæ are placed at each bifurcation; Flowers pure white, inodorous; Calyx composed of five lanceolate, entire leaves, continuing; these, as well as the whole of the plant, are beset with minute transparent globules, visible with a magnifier. Corolla deciduous, tube linear, grooved, pale green, slightly villous, bending a little upward; upper lip very narrow, bent back, bifid at the extremity, the edges toward the base rolled back, so as to make it appear in that part almost tubular; lower lip depending, trifid, segments equal, obtuse at the base, round the mouth of the tube marked with fine purple dots; Filaments two, short, projecting from the mouth of the tube, finally bending back; Antheræ at first yellow, afterwards livid; Germen oblong, smooth; Style capillary, slightly hairy; Stigma bifid.

The plant is increased by cuttings; in the *Hort. Malab.* the bruised leaves are described as in use for the cure of cutaneous eruptions.

[326]

MESEMBRYANTHEMUM VIRIDIFLORUM. GREEN-FLOWERED FIG-MARIGOLD.

Class and Order.

ICOSANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-fidus. *Petala* numerosa linearia. *Caps.* carnosae inferae polyspermae.

Specific Character and Synonyms.

MESEMBRYANTHEMUM *viridiflorum* foliis semicylindraceis papuloso-pilosis, calycibus quinquefidis hirsutis. *Ait. Kew.* v. 2. p. 196. *Haworth Mesemb.* p. 199. *Syst. Nat. ed. Gmel.* p. 848.



N°. 326

Of the seventy species of *Mesembryanthemum* described in the *Hortus Kewensis*, this is the only one with green flowers.

Mr. MASSON introduced it from the Cape in 1774.

The flowers are not only remarkable for their colour, but the extreme fineness of the florets; they begin to come forth in July, and continue to be produced till the end of September; the plant is easily increased by cuttings, is of ready growth, and blows freely.

In Mr. HAWORTH'S *Observations on the Genus Mesembryanthemum*, lately published, there is a very ample description of it, to which we refer such of our readers as wish for more minute information respecting it.

[327]

CHRYSANTHEMUM INDICUM. INDIAN CHRYSANTHEMUM.

Class and Order.

SYNGENESIA POLYGAMIA SUPERFLUA.

Generic Character.

Recept. nudum. Pappus marginatus. Cal. hemisphæricus, imbricatus: squamis marginalibus membranaceis.

Specific Character and Synonyms.

CHRYSANTHEMUM *indicum* foliis simplicibus ovatis sinuatis angulatis serratis acutis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 773. Spec. Pl. 1253. Thunb. Jap. p. 320.*

TSJETTI-PU. *Rheed. Mal. t. 44.*

MATRICARIA *sinensis*. *Rumph. Amb. p. 259. t. 91.*

MATRICARIA *japonica* maxima flore roseo sive suave rubente pleno elegantissimo. *Breyn. Prod. p. 66?*

KIK, KIKF, vel KIKKU. *Kæmpf. Amæn. Ex. p. 875.*



N^o. 327

We rejoice in the opportunity afforded us, of presenting our readers with the coloured engraving of a plant recently introduced to this country, which, as an ornamental one, promises to become an acquisition highly valuable.

This magnificent species of *Chrysanthemum*, which we have hitherto seen only in the collection of Mr. COLVILL, Nurseryman, King's-Road, Chelsea, began to flower with him early in November last, 1795; and as there were many buds on the plant, at that time, yet unopened, it appeared as if it would continue to flower during the early part of the winter at least.

It is a plant of strong growth; the stem rising to the height of two or three feet, somewhat woody, much branched, beset with numerous leaves, having some resemblance to those of Mugwort, of a greyish hue; the flowers, on being smelt to, discover an agreeable fragrance, they are produced on the summits of the branches in a loose sort of cluster (those which terminate the main stem, grow to the size of a large carnation) of a dark purple colour; they are, it is to be observed, double, or rather between semidouble and double; the florets of the radius at first perfectly tubular or quilled, as they advance split gradually downward on the inside, their outside is of a greyish tint, which being visible in most of the florets, especially the younger ones, gives them a particoloured appearance; these florets do not so completely occupy the receptacle but that room is left for a few others in the centre, of a different form, and yellow colour, which, on examination, appear to have their parts perfect, as indeed do those of the radius; the receptacle is beset with membranous paleæ or chaffy scales, a circumstance which would lead us to consider this plant rather as an *Anthemis* than a *Chrysanthemum*, of which it has the calyx, with the foliage of *Mugwort*.

New as this plant is to us, it appears to have been cultivated in China for ages: LINNÆUS, who describes it in his *Species Plantar.* refers us to a figure in the *Hortus Malabaricus*; this figure, and the description accompanying it, agree generally with our plant, but the flowers are more double, much smaller, less clustered, and do not correspond in point of colour, yet there can be no doubt but our plant is a variety of the same; it is there described as growing in sandy situations, and having green petals.

RUMPHIUS, in his highly interesting work, the *Herbarium Amboinense* is much more minute in his information; he observes, that these plants were originally brought from China, where they flower in May and June; that there are two sorts principally cultivated in India, the white and yellow-flowered, and a third sort, differing only in the colour of its flowers which are red (the variety, as we suppose, here figured) began to be known among them at Amboyna; the flowers there do not expand well, owing to their

being produced at the rainy season, and they decay without producing any seed.

He tells us further, that it is cultivated chiefly for pleasure; that the natives and the Dutch plant it only in the borders of their gardens, in which it does not succeed so well as in pots; and that, if it remains more than two years in the same spot, it degenerates, becomes less woody, and often wholly perishes; that the Chinese, by whom it is held in high estimation, pay great attention to its culture; they set it in pots and jars, and place it before the windows of their apartments, and that it is not unusual for them when they invite their friends to an entertainment to decorate their tables with it; on those occasions, he that produces the largest flower, is considered as conferring the greatest honour on his guests; besides these three varieties already mentioned, they have a fourth, which is more rare, whose flowers are of a greenish ash colour (is not this the var. figured in the *Hort. Malab.*?) all these varieties growing in separate pots, they place in certain quarters which they particularly wish to decorate, and the effect they produce is highly pleasing: in the cultivation of this plant they spare no pains, the shorter it is and the larger its flowers, the more it is esteemed; to make it dwarfish, and at the same time productive of flowers, they check its growth; for, if suffered to grow rude, it assumes a wild nature, and produces little but leaves when it is coming into flower, of the three blossoms which usually terminate each branch, they pluck off two, and thus the remaining flower grows larger; by this, and other management, they cause the flowers to grow to the breadth of one's hand: he enumerates still a fifth sort with white flowers, which is extremely rare, and smaller than the others, called *Tsehuy say si*, that is the *drunken woman*; its flowers morning and evening flag, and hang down as if debilitated by intoxication, in the middle of the day they become erect, and follow the course of the sun; but this (most probably a distinct species) is not exported from China. Finally, he remarks, that the Chinese and Malays are so attached to these flowers, that they even decorate their hair with them.

THUNBERG, in his *Flora Japonica*, enumerates it among the natives of that country, and describes it as growing spontaneously in Papenberg, near Nagasaki, and elsewhere, observing, that it is cultivated for the extreme beauty of its flowers in gardens and houses throughout the whole empire of

Japan, and that the flowers vary infinitely in point of colour, size, and plenitude. KÆMPFER's account of it in his *Amæn. Exot.* is very similar.

This Chrysanthemum appears to be a hardy greenhouse plant, and it is highly probable that, like the *Camellia* and *Aucuba*, it will bear the cold of our mild winters without injury.

As it flowers so late, there is but little prospect of its producing seeds with us, but it may be increased by cuttings, and parting of the roots.

[328]

TRIFOLIUM INCARNATUM. CRIMSON TREFOIL.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Flores subcapitati. Legumen vix calyce longius non dehiscens deciduum.

Specific Character and Synonyms.

TRIFOLIUM *incarnatum* spicis villosis oblongis obtusis aphyllis, foliolis subrotundis crenatis.
Linn. Syst. Vegetab. ed. 14. Murr. p. 689. Ait. Kew. v. 3. p. 87.

TRIFOLIUM spica rotunda rubra. *Bauh. Pin. p. 328.*

LAGOPUS maximus flore rubro. The greatest Haresfoot. *Park. Th. p. 1106. f. 1.*

TRIFOLIUM album incarnatum spicatum f. Lagopus maximus. *Bauh. Hist. 2. p. 376.*



N^o. 328

Of the annual species of *Trifolium* cultivated by the curious here, this is one of the largest, as well as one of the most shewy; the blossoms in the plants which we have had an opportunity of observing, have been of a

bright crimson colour, and have therefore corresponded badly with the name of *incarnatum*, originally applied to the plant by some of the old botanists, and adopted by LINNÆUS; like other Trefoils with red flowers, the blossoms are doubtless found with different shades of colour, and sometimes wholly white.

It is a native of Italy, a hardy annual, cultivated here by PARKINSON in 1640, flowers in July, and readily ripens its seeds, by which it is easily raised. It appears to have been lost out of this country since its first introduction, as it is not mentioned by MILLER; Mr. AITON enumerates it among the productions of Kew-Garden, and we last summer saw several plants of it raised from foreign seeds flowering in the garden of JOHN SYMMONS, Esq. Paddington-House, Paddington, who has to boast a collection of hardy herbaceous plants superior to most in this country; his readiness to oblige me with specimens for drawing on this occasion, and his various acts of kindness exerted in the promotion of my botanical views, I have to acknowledge with much gratitude.

[329]

ONONIS NATRIX. YELLOW-FLOWERED REST-HARROW.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum sessile. *Filamenta* connata absque fissura.

Specific Character and Synonyms.

ONONIS *Natrix* pedunculis unifloris aristatis, foliis ternatis viscosis stipulis integerrimis caule fruticoso. *Linn. Syst. Vegetab. ed. 14. Murr. p. 653. Ait. Kew. v. 3. p. 24.*

ANONIS viscosa spinis carens lutea major. *Bauh. Pin. 389.*



Nº. 329

The *Ononis Natrix*, a plant usually to be met with in all general collections of greenhouse plants, is a native of Spain, and the South of France, where it is said to grow wild in the corn-fields.

The general practice sanctioned by that of Mr. AITON, is to consider this species as tender; Mr. MILLER says it is very hardy, and recommends it to be planted in the open border, a treatment likely to suit it in mild winters; there is, however, one part of his account evidently erroneous, he describes the root as perennial, and the stem as herbaceous, this is not only contrary to LINNÆUS's specific description, but to fact, the stalk being undoubtedly shrubby.

As this plant in the course of a year or two is apt to grow out of form, it is advisable either to renew it frequently by seed, which it produces in abundance, or to keep it closely cut in.

It flowers from the middle of summer till towards the close, and is propagated readily either by seeds or cuttings.

Is no novelty in this country, having been cultivated by Mr. JAMES SUTHERLAND in 1683^[1].

[330]

SIDA CRISTATA. CRESTED SIDA.

Class and Order.

MONADELPHIA POLYANDRIA.

Generic Character.

Cal. simplex, angulatus. *Stylus* multipartitus. *Caps.* plures 1-spermæ.

Specific Character and Synonyms.

SIDA *cristata* foliis angulatis, inferioribus cordatis, superioribus panduriformibus, capsulis multilocularibus. *Sp. Pl. ed. 3. p. 964. Syst. Veg. ed. 14. Murr. p. 623. Ait. Kew. v. 2. p. 444. Cavanill. Diss. 1. t. 11. f. 2.*

ABUTILON *Lavateræ* flore, fructu cristato. *Dill. Elth. t. 2.*

ANODA hastata. *Linn. Syst. Nat. ed. Gmel. p. 1040.*



N^o. 330

DILLENIIUS has figured and described this plant in his *Hortus Elthamensis* as an *Abutilon*: LINNÆUS in his *Sp. Pl.* has ranked it with the *Sida*'s, in which he has been followed by Prof. MURRAY, Messrs. AITON and CAVANILLE; but

Prof. GMELIN, in the last edition of LINNÆUS' *Syst. Nat.* has made another new genus of it, by the name of *Anoda*; as his reasons for so doing are by no means cogent, we join the majority in continuing it a *Sida*.

It flowered in the garden of Mr. SHERARD, at Eltham, in 1725, and was introduced from Mexico, where it is a native: Mr. AITON considers it a stove plant, as he does the *Tropæolum majus*, and other natives of South-America; strictly speaking they may be such, but if raised early, and treated like other tender annuals, this plant will flower and ripen its seeds in the open ground, as we have experienced at Brompton.

It grows to the height of three feet, or more, producing during the months of July and August a number of blossoms in succession, which are large and shewy; the stigmata in this flower are curious objects, resembling the heads of Fungi in miniature.

[331]

KALMIA ANGUSTIFOLIA. NARROW-LEAV'D KALMIA.

Class and Order.

DECANDRIA MONOGYNIA.

Generic Character.

Cal. 5-partitus. *Cor.* hypocrateriformis, limbo subtus quinque corni. *Caps.* 5-locularis.

Specific Character and Synonyms.

KALMIA *angustifolia* foliis lanceolatis, corymbis lateralibus. *Linn. Syst. Veget. ed. 14. Murr. p. 404. Ait. Kew. v. 2. p. 64. Gronov. Fl. Virg. p. 65.*

CHAMÆDAPHNE sempervirens, foliis oblongis angustis, foliorum fasciculis oppositis e foliorum alis. *Catesb. Carol. app. t. 17. f. 1.*

LEDUM floribus bullatis fasciculatim ex alis foliorum oppositis nascentibus, foliis lanceolatis integerrimis glabris. *Trew. Ehr. t. 38.*



Nº. 331

In this work we have already given three different species of *Kalmia*, two commonly, and one more rarely cultivated with us, we mean the *hirsuta*, and which indeed we are sorry to find is scarcely to be kept alive in

this country by the most skilfull management; to these we now add another species, a native also of North-America, introduced by PETER COLLINSON, Esq. in 1736, two years after he had introduced the *latifolia*; CATESBY mentions its having flowered at Peckham in 1743; it is a low shrub, rarely rising above the height of two feet, growing spontaneously in swampy ground, and flowering with us from May to July; there are two principal varieties of it, one with pale and another with deep red flowers; these two plants differ also in their habits, the red one, the most humble of the two, not only produces the most brilliant flowers, but those in greater abundance than the other; Mr. WHITLEY, who has these plants in great perfection, assures me that it usually blows in the autumn as well as summer.

This shrub is extremely hardy, thriving best in bog earth, and is propagated most commonly by layers.

Like the *latifolia*, it is regarded in America as poisonous to sheep.

[332]

OENOTHERA FRUTICOSA. SHRUBBY OENOTHERA.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Calix 4-fidus. Petala 4. Capsula cylindrica infera. Semina nuda.

Specific Character and Synonyms.

OENOTHERA *fruticosa* foliis lanceolatis subdentatis, capsulis pedicellatis acutangulis, racemo pedunculato. *Linn. Syst. Veget. ed. 14. Murr. p. 358. Ait. Kew. v. 2. p. 4. L'Herit. Stirp. nov. t. 2. t. 5.*

OENOTHERA florum calyce monophyllo, hinc tantum, aperto. *Gron. virg. 42.*

LYSIMACHIA lutea caule rubente, foliis salicis alternis nigro maculatis, flore specioso amplo, vasculo seminali eleganter striato insidente, *Clayt. n. 36.*



N°. 332

Most of the *Oenothera* tribe are annual, have large yellow flowers, which open once only, and that in the evening, displaying their beauty, and exhaling their fragrance at a time which will not admit of their being much

enjoyed; the present species in some respects deviates from many of the others, the root is perennial, the flowers which are large and shewy, though they open in the evening, remain expanded during most of the ensuing day; the flower-buds, the germen, and the stalk are enlivened by a richness of colour which contributes to render this species one of the most ornamental and desirable of the tribe.

It is a hardy perennial, growing to the height of three or four feet, with us altogether herbaceous, and therefore improperly called *fruticosa*; a native of Virginia, flowering from June to August: was cultivated in 1739 by Mr. MILLER.

May be propagated by seeds, by parting of the roots, and also by cuttings.

CERINTHE MAJOR. GREAT HONEY-WORT.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corollæ limbus tubulato ventricosus: fauce pervia. *Semina* 2, bilocularia.

Specific Character and Synonyms.

CERINTHE *major* foliis amplexicaulibus, corollis obtusiusculis patulis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 187. Ait. Kew. v. 1. p. 183.*

CERINTHE *glaber* foliis oblongo-ovatis glabris amplexicaulibus, corollis obtusiusculis patulis. *Mill. Dict. ed. 6. 4to.*

CERINTHE flore ex rubro purpurascente. *Bauh. pin. p. 258.*

CERINTHE major. Great Honiewoort. *Ger. Herb.*



Nº. 333

Ancient writers on plants, supposing that the flowers of this genus produced abundantly the material of which bees form their wax, gave it the name of *Cerinthe*, which rendered into English would be wax-flower or

waxwort, not honeywort, by which the genus has long been, and is now, generally called.

Of this genus there are only two species known, the *major* and the *minor*, both happily distinguished by the different form of their flowers, a part from which it is not common to draw specific differences, though in some instances they afford the best.

The *major* varies much, the leaves being sometimes spotted, very rough, and the flowers of a more yellow hue; this is the sort figured by GERARD in his Herbal, who mentions its growing in his garden (1597). MILLER considers this as a species but LINNÆUS, HALLER, AITON, and others, regard it as a variety; our figure represents the *Cerinth glaber* of MILLER.

This is an annual, remarkable for the singular colour of its foliage; its flowers, though not very brilliant, possess a considerable share of beauty; both combined render it worthy a place in our gardens, more especially as it is a plant of easy culture, coming up spontaneously from self-sown seeds, and being a native of Switzerland, as well as the more southern parts of Europe, seedling plants produced in the Autumn rarely suffer by our winters. It flowers in July, August, and September.

[334]

HYPERICUM MONOGYNUM. CHINESE ST. JOHN'S-WORT.

Class and Order.

POLYADELPHIA POLYANDRIA.

Generic Character.

Cal. 5-phyllus. *Petala* 5. *Nect.* 0. *Capsula.*

Specific Character and Synonyms.

HYPERICUM *monogynum* floribus monogynis, staminibus corolla longioribus, calycibus coloratis, caule fruticoso. *Linn. Syst. Vegetab. ed. 14. Murr. p. 702. Hort. Kew. v. 3. p. 108.*



N°. 334

Of this genus 28 species are enumerated in the *Hortus Kewensis* of Mr. AITON, 42 in Prof. MURRAY's *ed.* of the *Systema Vegetab.* and 64 in Prof. GMELIN's 13th ed. of *Linn. Syst. Nat.* of the latter number 14 are described

with 5 styles, 46 with 3, 2 with 2 styles, and 2 with 1; when the term *monogynum* was first applied to this species, it was a proper one, there being then only one in that predicament, another having since been discovered it ceases to be so now; some have indeed doubted the propriety of using the word *monogynum* at all, alleging that in reality there are five styles, which manifestly shew themselves above, though they coalesce below; such is the opinion of my friend, Dr. GWYN; this is a point on which Botanists will think differently.

This elegant native of China, now common in our greenhouses, appears from Mr. MILLER to have been first introduced to this country in 1753, by HUGH, Duke of NORTHUMBERLAND; he tells us, that the plants were raised in his Grace's curious garden at Stanwick, from whence the Apothecaries garden at Chelsea was furnished with it.

Mr. MILLER has given us a minute description of this plant, which he observes is the more valuable, as it continues in flower great part of the year; he observes further, that if planted in a very warm situation, it will live in the open air, but that those plants which stand abroad will not flower in winter, as those do which are removed into shelter in autumn.

It may be propagated by slips from the root, or by layers.

ONONIS ROTUNDIFOLIA. ROUND-LEAVED REST-HARROW.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum sessile. *Filamenta* connata absque fissura.

Specific Character and Synonyms.

ONONIS *rotundifolia* fruticosa, foliis ternatis ovatis dentatis, calycibus triphyllo-bracteatis, pedunculis subtrifloris. *Linn. Syst. Vegetab. ed. 14. Murr. p. 653. Ait. Kew. v. 3. p. 25.*

ONONIS *rotundifolia* fruticosa, pedunculis trifloris, calycibus triphyllo-bracteatis, foliis ternatis subrotundis. *Linn. Sp. Pl. ed. 3. p. 1010.*

CICER sylvestre latifolium triphyllum. *Bauh. Pin. 347.*

CICER sylvestre tertium. *Dod. Pempt. 525.*



N^o. 335

Prof. JACQUIN, and most modern writers on Botany, consider the *Ononis* here figured, as the *rotundifolia* of LINNÆUS; it accords certainly with the figure of *Dodon.* to which that author refers, but is irreconcilable with his

description; the leaves for example are neither *parva*, *integerrima*, nor *glabra*, the words by which LINNÆUS describes them; they are indeed evidently serrated in the figure of *Dodon.* which he quotes: by the name of *rotundifolia*, however, this plant is now very generally known in our nurseries, to which its beauty has gained it admission. LOBEL tells us in his *Adversaria*, printed in 1576, that the plant was then growing in the garden of a Mr. MORGAN; as it is not enumerated in Mr. MILLER's Dictionary, ed. 6, 4to, ^[2] we suspect that it has been lost out of the country and re-introduced.

Baron HALLER informs us, that it is found wild in abundance at the bottom of the Alps in Switzerland; it is found also in other parts of Europe.

It flowers in our open borders from May to July, in which it ripens its seeds, by which it is in general propagated, as also by slips; it grows to about the height of a foot and a half, is very hardy, and easy of culture.

[336]

LOTUS HIRSUTUS. HAIRY BIRD'S-FOOT-TREFOIL.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Legumen cylindricum striatum. *Alæ* sursum longitudinaliter conniventes. *Cal.* tubulosus.

Specific Character and Synonyms.

LOTUS *hirsutus* capitulis subrotundis, caule erecto hirto, leguminibus ovatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 691. Ait. Kew. v. 3. p. 93.*

LOTUS *polyceratos* frutescens incana alba, siliquis erectis crassioribus et brevioribus rectis. *Moris. Hist. 2. p. 177. s. 2. t. 18. f. 14.*

LOTUS *pentaphyllos* siliquosus villosus. *Bauh. Pin. 332.*



Nº. 336

The *Lotus hirsutus*, according to LINNÆUS, is a native of the South of France, Italy, and the East.

In mild winters it will bear the open border with us, but it is more generally kept in the greenhouse, of which indeed it has long had possession, being cultivated, according to Mr. AITON, in 1683, by Mr. JAMES SUTHERLAND.

It flowers from June to August.

Its blossoms are of a white colour, which being contrasted with the redness of the calyx, gives them a pleasing appearance; CASP. BAUH. informs us, that they are purple also.

This shrub will acquire the height of several feet; its flowers are usually succeeded by seed-vessels, which in favourable situations produce perfect seeds, by which the plant is easily propagated, as also by cuttings.

[337]

PRUNELLA GRANDIFLORA. GREAT-FLOWERED SELF-HEAL.

Class and Order.

DIDYNAMIA GYMNOSPERMIA.

Generic Character.

Filamenta bifurca; altero apice antherifera. Stigma bifidum.

Specific Character and Synonyms.

PRUNELLA *grandiflora* foliis omnibus ovato oblongis subserratis, calicis labio superiore profunde trilobo. *Linn. Syst. Nat. ed. 13. Gmel. p. 920. Jacq. Fl. Austr. v. 4. t. 377.*

PRUNELLA *prima*. *Clus. Hist. xlii.*

PRUNELLA *cærulea* magno flore. *Bauh. Pin. 261.*



N^o. 337

The plant here figured, which we have several years cultivated in our garden at Brompton, without discovering in it the least disposition to vary, is undoubtedly the same as is figured by Prof. JACQUIN, in his *Fl. Austr.*

under the name of *grandiflora*; he regards it as a distinct species, and as such it is introduced in Prof. GMELIN's ed. of the *Syst. Nat.* of LINNÆUS: Mr. AITON, in his *Hort. Kew.* following LINNÆUS, makes it a variety of the *vulgaris*, a common English plant, which we have never seen to vary much in the size of its flowers.

Prof. JACQUIN informs us, that it grows wild, mixed with the *laciniata* (a kindred species with yellowish flowers) on the Alps; it is found also in similar situations in various other parts of Europe.

In July and August, it puts forth its large shewy blossoms, of a fine purple colour.

Such as are partial to hardy herbaceous plants, of ready growth, which are ornamental, take up but little room, and are not apt to entrench on their neighbours, will be induced to add this to their collection.

It is propagated by parting its roots in autumn.

[338]

ALLAMANDA CATHARTICA. WILLOW-LEAV'D ALLAMANDA.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Contorta. Caps. lenticularis erecta echinata 1-locularis polysperma. *Semina* bractæata.

Specific Character and Synonyms.

ALLAMANDA *cathartica*. Linn. Mant. p. 214. Suppl. p. 165. Syst. Vegetab. ed. 14. Murr. p. 252. Ait. Kew. V. 3. p. 489.

ORELIA *grandiflora*. Aubl. Guian. p. 271. t. 106.



N^o. 338

This beautiful stove plant (a native of Cayenne and Guiana, where it is said to grow by the sides of rivers) was introduced to this country in 1785,

by Baron HAKE^[3]; it has since flowered in many of our collections, usually in June and July.

Stem or trunk shrubby, upright, climbing to a considerable height; bark of the old wood pale brown, of the young wood green and smooth; leaves generally growing four together, sometimes only two, and those opposite, sessile, smooth and glossy above, paler green beneath, the midrib on the underside evidently villous, veiny, veins as in *Plumeria*, terminating before they reach the margin, chewed discovering little taste; flowers very large, produced even on young plants, of a fine yellow colour, and somewhat spicy smell, not terminal, but growing from the sides of the branches, sometimes singly, more frequently three or four together, standing on short footstalks; calyx composed of five leaves, lanceolate, smooth, unequal, short compared with the flower; corolla mono-petalous, funnel-shaped, tube nearly cylindrical, limb dilated below and bellying out, on the inside striped with orange-coloured veins, above divided into five segments, which roll over each other before they open, when Open broad, somewhat truncated, one side rounded, the other terminating in a short point; at the base Of the divisions the yellow colour on the inside of the flower becomes nearly white, forming so many spots; the mouth of the tube is perfectly closed with villi converging to a point, and which serve as a covering or species of thatch to five long, pointed, rigid, arrow-shaped antheræ beneath them, which sit on the top of so many ridges, projecting from the inside of the tube, and which just beneath the antheræ are villous; germen nearly round, with a slight ridge on each side of it, surrounded at its base by a greenish glandular substance; style filiform, the length of the tube of the corolla, enlarging as it comes near to the stigma; stigma divisible into three distinct parts, the lowermost of a yellowish hue, in the form of a rim turning downwards, the middle of a dark green colour, secreting honey in considerable quantity; the summit, which perhaps is the true stigma, a short conical point, rising from the centre of the middle part, bifid at top with a furrow running down each side of it; seed-vessels according to the figure in AUBLET, are very large and prickly.

Of this genus, which has a considerable affinity with that of *Vinca* and *Plumeria*, only one species is described in LINNÆUS's works, and this is usually increased by cuttings.

[339]

ARUM TRILOBATUM. THREE-LOBED ARUM.

Class and Order.

GYNANDRIA POLYANDRIA.

Generic Character.

Spatha monophylla cucullata. *Spadix* supra nudus, inferne femineus, medio stamineus.

Specific Character and Synonyms.

ARUM *trilobatum* acaule, foliis sagittato trilobis, flore sessili. *Linn. Syst. Veget. ed. 14. Murr. p. 828. Ait. Kew. V. 3. p. 316. Mill. Icon. t. 52. f. 2.*

ARISARUM *amboynicum*. *Rumph. Amb. 5. p. 320. t. 110. f. 2.*



Nº. 339

Mr. MILLER in his figure of this plant, to which LINNÆUS refers, has been more happy in his representation than in that of many others; RUMPHIUS' figure and description accord also with our plant, some of his leaves indeed

are more perfectly three-lobed than any we have seen here on the living plant and to this variation, he informs us, they are subject.

We learn from Mr. MILLER, that roots of this Arum were brought from Ceylon in the year 1752. It flowers in May and June, and is regarded both by Mr. MILLER and Mr. AITON as a stove plant; we have seen it succeed very well with the treatment of a tender greenhouse plant.

It is one of the least of the tribe; its root is like that of the common Arum, and extremely acrid: but the plant is more particularly distinguished by the rich, brown, velvety appearance of its flowers; the length of its tapering spadix, which on its lower part is full of little cavities, and resembles a piece of metal corroded by long exposure; and by the intolerable stench which the whole of the flower, but more especially the spadix, sends forth.

It is a native of Amboyna, as well as of Ceylon. RUMPHIUS informs us that the roots, sometimes eaten raw by mistake, cause violent inflammations of the mouth and throat, and that they do not lose their acrimony even when boiled.

The plant increases freely by offsets from its roots.

[340]

POLYGALA HEISTERIA. HEATH-LEAVED MILKWORT.

Class and Order.

DIADELPHIA OCTANDRIA.

Generic Character.

Cal. 5-phyllus: foliolis 2 majoribus alæformibus, ante maturitatem seminis coloratis. *Capsula* obcordata 2-ocularis, *Semina* solitaria.

Specific Character and Synonyms.

POLYGALA *Heisteria* floribus imberbibus lateralibus, caule arborescente, foliis triquetris mucronato-spinosis. *Linn. Syst. Veg. ed. 14. Murr. p. 639. Ait. Kew. v. 3. p. 6.*

SPARTIUM africanum frutescens ericæ folio. *Comm. Hort. 2. p. 193. t. 97.*

THYMBRA capensis nepetæ Theophrasti foliis aculeatis flore parvo purpureo. *Pluk. Alm. 366. t. 229. f. 5.*

HEISTERIA. *Hort. Cliff. 352.*



N^o. 340

After receiving various names, this plant has been finally placed by LINNÆUS among the Milkworts; it retains the trivial name of *Heisteria*,

instead of the generic one previously bestowed on it, in honour of Prof. HEISTER, the celebrated German Surgeon.

In Holland it appears to have been long since known; but was a stranger here, till introduced from the Cape, by Mr. MASSON, in 1787. *Ait. Kew.*

In the course of a few years it becomes a shrub of considerable size, equalling a small furze bush, to which, in its habit, it bears a distant resemblance; and furze-leaved, in our humble opinion, would have been a more expressive name than heath-leaved, which Mr. AITON has given it in accordance with COMMELIN's idea.

The purple of the flowers is brilliant in the extreme, and as those are plentifully produced almost the year through, it has very generally obtained a place in collections of greenhouse plants about London.

It is commonly increased by cuttings, but not easily.

SCILLA AMÆNA. BYZANTINE SQUILL.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. 6-petala patens, decidua. *Filamenta* filiformia.

Specific Character and Synonyms.

SCILLA *amæna* floribus lateralibus alternis subnutantibus, scapo angulato. *Linn. Syst. Veg. ed. 14. Murr. p. 328. Ait. Kew. v. 1. p. 444. Jacq. Fl. Austr. v. 3. t. 218.*

HYACINTHUS *stellaris cæruleus amænus. Bauh. Pin. 46.*

HYACINTHUS *stellaris Bizantinus. Hort. Eyst. Vern. 2d. ord. fol. 13.*

HYACINTHUS *stellatus Byzantinus alter sive flore Boraginis. The other Starry Jacinth of Constantinople. Park. Parad. p. 128. t. 131. f. 4.*

HYACINTHUS *stellaris, caulibus pluribus ex eodum bulbo ortis singulis pluribus floribus oneratis. Moris. Hist. 2. p. 374. s. 4. t. 12. f. 17.*



Nº. 341

The *Scilla amœna*, a hardy bulbous plant of small growth, is a native of the Levant, and has long been cultivated in this country for ornament, being introduced about the year 1600, by ED. LORD ZOUCH, (vid. *Park. Parad.* &

Lobel. Avers.); when it grows luxuriantly, many flowering stems will spring from the same root, which, when the plant is in flower, are altogether upright; as the seed-vessels advance in size and weight, they bend down, the blossoms are violet-colored, and the germen in the centre of each is distinguished for the paleness of its colour, a character which did not escape the observation of PARKINSON.

It flowers early in May, and, in favourable situations, ripens its seeds in the beginning of June.

Is usually increased by offsets, which are produced in tolerable abundance; will grow in almost any soil or situation; but will succeed best if planted in a sandy loam, and a warm sheltered part of the garden.

Its seeds are of a pale amber colour, and drop soon.

Though Professor JACQUIN figures it in his *Fl. Austriaca*, he suspects that it had originally been introduced from Constantinople.

[342]

ERICA PERSOLUTA. BLUSH-FLOWERED HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* bifidæ. *Caps.* 4-locularis.

Specific Character and Synonyms.

ERICA *persoluta* antheris aristatis inclusis, corollis campanulatis, calycibus ciliatis, foliis quaternis. *Linn. Mant. p. 230. Ait. Kew. v. 2. p. 17.*

ERICA *subdivaricata*. *Berg. Cap. 114.*

ERICA *persoluta* aristata foliis ternis quaternisque trigonis glabris, floribus umbellatis, calice ciliato. *Thunb. Prod. Pl. Cap. p. 73.*



N^o. 342

The different species and varieties of the African Heaths are now become so numerous, that there is scarcely any period of the year in which some of them may not be found to delight the eye with their blossoms: the *persoluta*

here figured is one of those which flower early in the spring; through the months of March, April and May, its branches are loaded with a profusion of bright purple flowers, which, joined to the lively verdure of its foliage, places it among the most desirable of the tribe.

If suffered to grow, it will form a shrub of considerable height: there is a variety of it, called *alba*, whose flowers are nearly white, yet not wholly divested of a red tinge; to this variety the term blush-coloured, which Mr. AITON has given to the species, seems most applicable.

This heath was introduced, from the Cape, by Mr. MASSON, in 1774. *Ait. Kew.*

It is increased without much difficulty from cuttings, and is more easily preserved than many of the others.

ANTHOLYZA CUNONIA. SCARLET-FLOWERED ANTHOLYZA.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Cor. tubulosa irregularis recurvata. *Caps.* infera.

Specific Character and Synonyms.

ANTHOLYZA *Cunonia* corollis rectis, labii quinquepartiti lobis duobus externis latioribus adscendentibus. *Linn. Sp. Pl. ed. 3 p. 54. Ait. Kew. v. 1 p. 66.*

ANTHOLYZA *Cunonia* corollis subpapilionaceis: labii lobis duobus externis latioribus adscendentibus. *Linn. Syst. Veg. ed. 14. Murr. p. 87.*

ANTHOLYZA *Cunonia* foliis ensiformibus glabris striatis, spica oblonga disticha. *Thunb. Prod. Pl. Cap. p. 7.*

CUNONIA floribus sessilibus spathis maximis. *Butt. cun. 211. t. 1.*

CUNONIA *Antholyza*. *Mill. Dict.*



N°. 343

Amidst that variety of tender bulbous plants which the Cape affords, there are none whose flowers can boast a richer scarlet, than the *Antholyza* here figured; in their form also, there is a great singularity, and in their

mode of growth much elegance; to us indeed it is matter of surprise that this plant should not be more generally cultivated, especially as its culture is attended with so little trouble: Mr. MILLER, who grew it in 1756, has minutely described it in his Dictionary; he informs us, that it is easily propagated by offsets, or raised from seeds; the latter, as the plant flowers in April and May, ripen about the middle of June, and should be sown about the middle of August; the plant with the pot in which it has grown, after it has ripened its seeds, should be set by the shed or some dry place, and in the autumn the largest roots should be selected, and three of them planted in a pot of a moderate size, keeping them in an open frame till the approach of winter, when they are to be removed into the house with other greenhouse plants.

This species is found wild in Persia as well as at the Cape.

[344]

ASPALATHUS PENDUNCULATA. SMALL-LEAVED ASPALATHUS.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 5-fidus: lacinia superiore majore. *Legumen* ovatum muticum subdispermum.

Specific Character and Synonyms.

ASPALATHUS *pendunculata* foliis fasciculatis subulatis glabris, pedunculis filiformibus folio duplo longioribus. *L'Herit. Sert. Angl. t. 26. Ait. Kew. v. 3. p. 16.*



N^o. 344

Of the five species of *Aspalathus* enumerated in the *Hortus Kewensis*, four are natives of the Cape; of those, three have been introduced by Mr. MASSON: the *albens* and *candicans* in 1774, the present species in 1775.

This is one of the plants figured by Mons. L'HERITIER, in a work of his, entitled *Sertum Anglicum*, containing new species observed by him during his excursion to this country a few years since, most of which flowered in the royal garden at Kew.

It will grow to the height of several feet, produces abundance of blossoms, which open during most of the summer months, and is propagated by cuttings.

[345]

POLYGALA BRACTEOLATA. SPEAR-LEAVED MILKWORT.

Class and Order.

DLADELPHIA OCTANDRIA.

Generic Character.

Cal. 5-phyllus; foliolis alæformibus coloratis. *Legumen* obcordatum biloculare.

Specific Character and Synonyms.

POLYGALA *bracteolata* floribus cristatis racemosis, bractæis triphyllis, foliis lineari-lanceolatis, caule fruticoso. *Linn. Syst. Veget. ed. 14. Murr. p. 638. Berg. Cap. 182. Buxb. Cent. 3. p. 4. t. 71. Ait. Kew. v. 3. p. 4.*



N°. 345

To the vast number of ornamental plants introduced from the Cape of Good Hope by Mr. MASSON, we here add another, imported in 1787; when Mr. AITON published his *Hortus Kewensis*, it most probably had not

flowered with him, as he does not notice its particular time of blowing: we now see it in most of the Nurserymen's collections of greenhouse plants, flowering from May to July. It is a plant of great singularity and beauty, the purple of its flowers is brilliant in the extreme.

It is increased by cuttings.

[346]

PROTEA MELLIFERA. HONEY-BEARING PROTEA.

Class and Order.

TETRANDRIA MONOGYNIA.

Generic Character.

Cor. 4-petala: interdum basi vel apice coherentibus. *Antheræ* lineares vel oblongæ petalis infra apicem insertæ. *Cal.* propius 0. *Nux* supera, 1-locularis, evalvis.

Specific Character and Synonyms.

PROTEA *mellifera* foliis lanceolato-ellipticis capituloque terminali oblongo glabris. *Thunb. Prot. n. 37. Linn. Syst. Veget. ed. 14. Murr. p. 139. Ait. Kew. vol. 1. p. 127.*

LEUCADENDRON repens foliis lanceolatis, floribus oblongis, calycum squamis glabris. *Linn. Sp. Pl. ed. 3. p. 135.*

LEPIDOCARPODENDRON foliis angustis brevioribus salignis, calycis squamis elegantissimi ex roseo aureo albo atro rubro variegatis, florum plumis albis. *Boerh. Lugd. 2. p. 187.*



N°. 346

To a magnificent appearance, the blossom of the Protea joins a structure extremely curious and interesting; but, at the same time, difficult to be understood by students, who are apt to consider the whole as one great

simple flower, while in reality it is composed of a number of florets enclosed within a common calyx formed of numerous leaves or scales placed one over the other, and sitting on one common receptacle, being in fact what Botanists term an aggregate flower, approaching indeed near to a compound one, there being in the structure and union of the antheræ a considerable similarity to those of the Syngenesia class, much more so than in the flowers of Plantain, Scabious, Teasel, and others.

The florets of the present species correspond extremely well with the character of the genus *Leucadendron* in the sixth edition of the *Genera Pl.* of LINNÆUS, but not with that of *Protea*, as given in the *Hort. Kew.* and *Gmel. ed. Linn. Syst. Nat.* to which it is now united; the corolla being most evidently composed of two (not four) petals, the largest of these is trifid at top, each segment of it, as well as the summit of the smaller petal, terminates in a twisted kind of plume, not peculiar to this species; of the antheræ, which are long, linear, and form a kind of cylinder, three are attached to the largest petal, the fourth (which appears to be less perfect than the others) to the smaller petal; the germen is enveloped with numerous orange-coloured hairs, having the gloss of the richest sattin; the antheræ terminate in small appendages of a brown colour.

This magnificent shrub, a principal ornament of the Cape-House at Kew, is a native of the Cape of Good Hope, from whence it was introduced by Mr. MASSON, in 1774 (*Ait. Kew.*); it flowers chiefly in the Spring, and often during the Summer; is propagated principally by layers.

Our drawing was made from a plant raised from Cape seeds, which flowered this season, among a number of others equally curious, in the collection of Mr. BARR, Nurseryman and Florist, near Ball's-Turnpike, Kingsland.

[347]

OENOTHERA ROSEA. ROSE-COLOURED OENOTHERA.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-fidus. *Petala* 4. *Caps.* cylindrica infera. *Sem.* nuda.

Specific Character and Synonyms.

OENOTHERA *rosea* foliis ovatis dentatis, inferioribus lyratis: capsulis clavatis. *Ait. Kew.* v. 2. p. 3.

OENOTHERA *rosea*. *L'Herit. Stirp. Nov. tom.* 2. t. 6.



N. 347

Of the several different species of this genus growing in our garden at this present writing (eleven in number) two only are of a red or purple

colour, the one here figured, and another with a larger flower, which we purpose soon to publish under the name of *purpurea*.

The present species, Mr. AITON informs us, was introduced in 1783 by Mons. THOUIN, from Peru.

It has been considered as a greenhouse plant, and a perennial; we find it to be more hardy than greenhouse plants in general, and scarcely entitled to the distinction of a perennial.

It may be increased by cuttings and seeds, the latter of which are plentifully produced. It rarely exceeds a foot in height; its rose-coloured flowers expand during the whole of the day, and are produced during most of the Summer months.

To guard against accidental severity of weather, sow its seeds in the Spring with tender annuals; when the plants have acquired a proper age and the season is favourable, plant them out singly in the open border.

[348]

CALCEOLARIA FOTHERGILLII. FOTHERGILL'S SLIPPER-WORT.

Class and Order.

DIANDRIA MONOGYNIA.

Generic Character.

Corolla ringens inflata. *Capsula* 2-locularis, 2 valvis. *Calyx* 4-partitus æqualis.

Specific Character and Synonyms.

CALCEOLARIA *Fothergillii* foliis spathulatis integerrimis, pedunculis scapiformibus unifloris.
Ait. Kew. v. 1. p. 30. tab. 1. Linn. Syst. Nat. ed. 13. Gmel. p. 39.



N°. 348

To this species of *Calceolaria* the trivial name of *Fothergillii* has been given in honour of Dr. JOHN FOTHERGILL, who introduced it in 1777 with several other rare plants from Falkland-Islands, where it is a native.

In the *Hortus Kewensis* of Mr. AITON it was first figured, and minutely described.

This plant of great singularity and beauty particularly merits a place in the collections of the curious; its principal flowering season is in May, but it will often blossom much later in the Summer.

It is not to be had in perfection, nor even preserved, without great attention, joined to a peculiar treatment; several times it has been nearly lost from this country. Mr. AITON makes it a biennial greenhouse plant; it, no doubt, is one of those plants which are liable to be killed in very severe weather, and therefore it is necessary to keep a pot of it at least, during the Winter, either in the greenhouse or a well-secured frame; but the plant will be found to succeed best in the open ground, provided it be planted in bog-earth, in a situation that is moist and moderately shady; with us it has proved perfectly perennial.

With care it can be propagated by layers and cuttings, but with most certainty in the former mode; possibly it may be raised from seeds, they ripen here, but are as fine as dust.

[349]

SOLANUM LACINIATUM. CUT-LEAV'D NIGHTSHADE.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corolla rotata. *Antheræ* subcoalitæ, apice poro gemino dehiscentes. *Bacca* 2-locularis.

Specific Character and Synonyms.

SOLANUM *laciniatum* caule fruticoso inermi glaberrimo, foliis pinnatifidis: laciniis lanceolatis acutis, paniculis axillaribus binis ternisve. *Ait. Hort. Kew. v. 1. p. 247.*



N^o. 349

Mr. AITON in his *Hort. Kew.* mentions this plant as being a native of New-Zealand, on the authority of Sir JOSEPH BANKS, and that it was introduced in 1772; he regards it as a stove plant; it has been since found to

be more hardy than he imagined, and not to require more heat than the greenhouse affords; in the Summer it will stand abroad, and even ripen its fruit in the open air.

It is a plant of some beauty, but is more remarkable for having its antheræ separating widely from each other, and thereby losing the character of a *Solanum*, so far as it depends on that circumstance. The berries when ripe are of the size of a small plum, and of a yellowish green hue; their pulp is sweet, in some small degree resembling that of a fig, whether it be so innocent we do not take on us to assert.

It flowers during most of the Summer, and is easily increased by cuttings or seeds.

[350]

ERICA VENTRICOSA. PORCELAIN HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Calyx 4-phyllus. *Corolla* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* bifidæ. *Caps.* 4-ocularis.

Specific Character and Synonyms.

ERICA *ventricosa* mutica, foliis quaternis trigonis ciliatis, floribus ventricosis glabris. *Thunb.*
Prod. Cap. p. 71.



Nº. 350

Of the many new Heaths which have been introduced within these few years, none have excited greater admiration than the present one; its blossoms, though they cannot boast the grandeur or richness of colour so

strikingly displayed in the *cerinthoides*, and some others, please more on a near inspection, they have indeed a delicacy and beauty which are indescribable; we have given to it the English name of Porcelain Heath, as the flowers have somewhat the appearance of porcelain, or enamel.

This species has been introduced from the Cape, since the publication of the *Hortus Kewensis*; we saw it in blossom many years since, in the Royal Garden at Kew; several varieties of it have been raised from Cape seeds by Mr. LODDIGES, Nurseryman, at Hackney, differing in the hairiness of their leaves, size and colour of their blossoms; but the best variety we have seen is the one here represented.

In point of form the blossoms resemble those of the *ampullacea*, but they are not so large, and have no viscidty; the stigma is enclosed within the mouth of the flower, and the peduncles are usually naked; in some very luxuriant specimens we have observed a scale or two on them.

It is with Heaths, in some respects, as it is with fruit trees: one season they will produce blossoms most abundantly, they seem indeed to overblow themselves, the next few appear; in different years the blossoms of the same heath will vary also considerably in size. The *ventricosa* is a free blower, and will in general produce flowers in abundance, for two or three months, from June to September.

Those who possess the knack of striking heaths, raise it by cuttings without much difficulty.

[351]

SAXIFRAGA MUTATA. SAFFRON-COLOUR'D SAXIFRAGE.

Class and Order.

DECANDRIA DIGYNIA.

Generic Character.

Calyx 5-partitus. *Corolla* 5-petala. *Caps.* 2-rostris 1-ocularis polysperma.

Specific Character and Synonyms.

SAXIFRAGA *mutata* foliis radicalibus aggregatis lingulatis cartilagineo-serratis, caule racemoso folioso, calyce villosa et viscido. *Linn. Syst. Vegetab. ed. 14. Murr. p. 411. Jacq. Ic. Pl. rar. Fasc. 9.*



Nº. 351

LINNÆUS has arranged several distinct species of Saxifrage under the *Saxifraga Cotyledon*, but very properly separates this; though in the general form of its foliage it is as nearly related to it as most of the others; from four

of them it is indeed perfectly distinct, instead of the sawed cartilaginous edge, it has only fine soft hairs, more especially towards the base; towards the extremity there is some appearance of little teeth, but widely differing from those of the others. The whole plant is covered with viscid hairs; the stalk about a foot high, is much branched; but the great peculiarity of this plant consists in its flowers, the petals being long, narrow, and pointed, and of a saffron colour, of a deeper tint when they first open, and gradually changing to a pale yellow; the beauty of the flowers is heightened by a glandular substance in the centre of each, which when the flower expands is of a bright purple colour.

This rare species of Saxifrage is a native of the Alps of Switzerland and Italy; we raised it with difficulty from seeds sent us by Mr. DAVAL, of Orbe, in Switzerland; it was three years before the plants flowered; those now in bloom in my garden, June 15, bid fair to produce seeds in abundance; the plant may also be produced from offsets, in the same way as the *Saxifraga Cotyledon*, vulgarly called *Pyramidal Sedum*, and requires a similar treatment; though an Alpine plant, it must be sheltered from much wet and severe frosts.

Baron HALLER describes it with spots on the petals; those certainly did not exist on our plants.

OENOTHERA PURPUREA. PURPLE OENOTHERA.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-fidus. *Petala* 4. *Caps.* cylindrica infera. *Sem.* nuda.

Specific Character and Synonyms.

OENOTHERA *purpurea* foliis ovato-lanceolatis glaucescentibus integerrimis, capsulis sessilibus, stigmatе atro-purpureo.



N^o. 352

The species of *Oenothera* here figured is the one we mentioned in treating of the *rosea*, a very ornamental species, introduced to the Royal Garden at Kew, from the western coast of North-America, in 1791.^[4] We

have given to it the name of *purpurea*, it being the only one in our gardens with purple flowers.

It is distinguished by three principal characters, the glaucous appearance of its foliage, the purple hue of its blossoms, and the dark colour of its stigma.

It flowers during the months of July and August, and ripens its seeds in September; is an annual of ready growth, and worthy of being more generally introduced to the flower-garden; its seeds should be sown in the spring, with other tender annuals, and the seedlings when of a proper age planted out singly in the open border, where they will acquire the height of two feet, and produce abundance of blossoms which, like those of the *rosea*, open during the day as well as in the evening.

MAHERNIA INCISA. CUT-LEAV'D MAHERNIA.

Class and Order.

PENTANDRIA PENTAGYNIA.

Generic Character.

Calix 5-dentatus. *Petala* 5. *Nect.* 5 obcordata filamentis supposita. *Capsula* 5-locularis.

Specific Character.

MAHERNIA *incisa* caule hispidulo, foliis lanceolatis incisis, stipulis integerrimis.



N°. 353

In point of size and mode of growth, this beautiful species comes near to the *pinnata* already figured in this work; but differs essentially in the

singular hispidity of its stalks, the form of its leaves, and the colour of its flowers.

The stalks to the naked eye discover a manifest roughness, a magnifying glass shews this roughness to be of a singular kind, that they are beset on every side with little protuberances, from whence issue tufts of pellucid hairs, and here and there a single hair is discoverable with a small red viscid globule at its extremity; a portion of the stalk, when highly magnified, somewhat resembles that of the creeping *Cereus*. The leaves, which are not so manifestly hairy as the stalk and calyces, are deeply jagged on the edges, and bear some little affinity in their appearance to those of the *Pelargonium tricolor*. The flowers, when in bud, are of the richest crimson imaginable; as they open they incline to a deep orange, and finally become of a yellowish hue.

As this plant produces abundance of blossoms, they may be seen in all their states during most of the Summer and Autumnal months.

The plant from whence our drawing was taken, flowered this Summer with Mr. COLVILL, Nurseryman, King's-Road, it requires the same treatment as the *pinnata*, is probably a native of the same country, and may be propagated by cuttings in the same manner.

[354]

MIMULUS AURANTIACUS. ORANGE MONKEY-FLOWER.

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Calyx 4-dentatus prismaticus. *Cor.* ringens: labio superiore lateribus replicato. *Caps.* 2-ocularis polysperma.

Specific Character.

MIMULUS *aurantiacus* caule erecto fruticoso tereti, foliis ovato-lanceolatis obtusiusculis.



Nº. 354

The present species of *Mimulus*, equal in point of shew to most of the inhabitants of our greenhouses, to which situation it is adapted, flowered this Summer with Mr. COLVILL, Nurseryman, King's-Road.

Stalk about three feet high, much branched, shrubby, round, the young wood green, with a tinge of purple toward the lower part of each joint, slightly viscid, as it becomes older changing to a light brown colour, and discovering manifest fissures; branches alternately opposite, flower-bearing quite to the base; leaves opposite, sessile, slightly connate, ovato-lanceolate, somewhat blunt at the extremity, this bluntness is particularly apparent when contrasted with a leaf of the *ringens*, toothed or slightly sawed on the edge, smooth, veiny; flowers inodorous, large, nearly twice the size of those of the *ringens*, uniformly pale orange, growing in pairs from the alæ of the leaves, standing on footstalks about half the length of the calyx; calyx five-angled and five-toothed, tube of the flower within the calyx, narrow, cylindrical, pale yellow, bent a little downward, gradually expanding, and dividing into two lips, the upper lip divided into two, the lower lip into three segments, all of them irregular, the two uppermost very much so; at the base of the middle segment of the lower lip are two prominent ridges, of a somewhat deeper colour; stamina four, two long, two short; antheræ deep orange, cruciform, within the flower; stigma white, two-lip'd, lips closed or expanded according to its age; style filiform; germen oblong; at the base of the germen is a gland of considerable size which secretes much honey.

This plant flowers during most of the Summer, and is increased by cuttings. We know not with certainty of what country it is a native.

[355]

OENOTHERA PUMILA. DWARF OENOTHERA.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-fidus. *Petala* 4. *Caps.* cylindrica infera. *Sem.* nuda.

Specific Character and Synonyms.

OENOTHERA *pumila* foliis lanceolatis obtusis glabris subpetiolatis, caulibus prostratis, capsulis acutangulis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 358. Ait. Kew. v. 2. p. 4.*

OENOTHERA foliis radicalibus ovatis caulinis lanceolatis obtusis, capsulis ovatis obtusis. *Mill. Ic. t. 188.*

LYSIMACHIA siliquosa glabra minor mariana angustioribus foliis. *Pluk. Mant. 123.*



N°. 355

From all the species of *Oenothera* with yellow flowers hitherto introduced to this country, the present one is distinguished by the inferiority of its size; in its most luxuriant state it rarely exceeds a foot in height, and

as it commonly grows it is far more humble: from a perennial root it sends out many flowering stems, which are productive of blossoms in regular succession from April to July: these are open in the morning as well as in the evening, which renders the plant more ornamental and desirable: for the border of a small garden, or for rock-work, it is extremely suitable, if not so shewy as some others; it is very hardy, of ready growth, not apt to encroach, flowers during most of the summer months, and is readily propagated by seeds, cuttings, or dividing of the roots in autumn.

Is a native of North-America, and was cultivated here by Mr. MILLER in 1757. *Ait. Kew.*

The stalks cannot be considered as prostrate, though LINNÆUS describes them as such; *adscendentibus* would be a more appropriate term.

[356]

ERICA MASSONI. MASSON'S HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* bifidæ. *Caps.* 4-locularis.

Specific Character and Synonyms.

ERICA *Massoni* antheris muticis inclusis, corollis cylindricis grossis, floribus capitatis, foliis octofariis imbricatis pubescentibus. *Linn. Suppl. p. 221. Ait. Kew. v. 2. p. 23.*

ERICA *Massoni* foliis quaternis oblongis serratis villosis, floribus cylindricis viscosis. *Thunb. Prod. Cap. p. 71.*



Nº. 356

This very magnificent African Heath is first described in the *Suppl. Pl.* of the younger LINNÆUS, and named in honour of Mr. MASSON, whose exertions at the Cape, where he resided many years as collector of the natural

productions of that fertile spot, have so eminently contributed to render the royal collection at Kew, in this tribe of plants especially, rich in the extreme.

This Heath grows to a considerable height, and becomes much branched, the branches are long and upright, covered closely with numerous hoary leaves, nearly cylindrical, most of these terminate in a large umbel of flowers, which continue a long while in bloom, and are so extremely viscous that scarcely a winged insect can settle on them and escape with its life; the formidable wasp sometimes becomes its victim, as we once had an opportunity of seeing.

When the *Hort. Kew.* was first published, this species had not flowered with Mr. AITON, it has since done so in many collections near town, and no where in greater perfection than at Messrs. LEE and KENNEDY'S, Hammersmith, on some one or other of whose plants blossoms may be seen during most of the summer.

It is raised from cuttings with extreme difficulty, is not so easily kept as some others, and is more susceptible of injury from moisture.

[357]

BRIZA MAXIMA. GREAT QUAKING GRASS.

Class and Order.

TRIANDRIA DIGYNIA.

Generic Character.

Cal. 2-valvis, multiflorus. *Spicula* disticha: valvulis cordatis, obtusis: interiore minuta.

Specific Character and Synonyms.

BRIZA *maxima* spiculis cordatis, flosculis septendecim. *Linn. Syst. Veget. ed. 14. Murr. p. 115.*
Ait. Kew. v. 1. p. 103.

GRAMEN tremulum maximum. *Bauh. Pin. 2.*

PHALARIS pratensis altera. *Pearle Grasse. Ger. emac. p. 87. f. 3.*



N^o. 357

The Grasses are in general more regarded for utility than ornament, yet in the latter point of view many of them have engaged the attention of the curious, and long held a place in the flower-garden, to which they have a

twofold claim, as they not only decorate the garden when fresh, but the mantle-shelf when dry; to these purposes the present species of Briza has long been applied: JOHNSON tells us, in his time, 1633, "it was sown yearly in many of the London gardens."

This species, a native of Spain and Italy, blossoms in June and July, and ripens its seeds in August.

Where it has once seeded it comes up spontaneously, without the trouble of sowing it; autumnal seedlings make the strongest plants, they are liable however to be cut off in very severe seasons; should that happen, sow more seed in the spring with your other annuals.

ERICA BACCANS. ARBUTUS-FLOWERED HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* apice bifidæ pertusæ. *Caps.* 4-locularis. 4-valvis polysperma.

Specific Character and Synonyms.

ERICA *baccans* antheris cristatis inclusis, corollis globoso-campanulatis calyce colorato inclusis, foliis imbricatis. *Linn. Mant. p.* 233.

ERICA *baccans* antheris cristatis, corollis globoso-campanulatis tectis, stylo incluso, foliis ternis imbricatis. *Linn. Syst. Veget. ed. 14. Murr. p.* 366. *Ait. Kew. v. 2. p.* 18.

ERICA *baccans* cristata foliis quaternis linearibus serrulatis, floribus umbellatis calyce ovato æquante. *Thunb. Prod. p.* 74.

ERICA Africana glabra fruticosa arbuti flore. *Seb. Mus. 1. p.* 32. *t.* 21. *f.* 3.



N^o. 358

SEBA, a Dutch writer, appears first to have noticed this Heath; he figures it in his Museum, and distinguishes it by the name of Arbutus-flowered, which Mr. AITON has retained.

This very elegant and ornamental species grows to a considerable height, and in favourable situations produces abundance of flowers early in the summer, which are remarkable, though not peculiarly so, for being enveloped with a calyx of same colour.

It is a native of the Cape, and was introduced by Mr. MASSON in 1774^[5].

Is raised from seeds, which it ripens with us more freely than most of the African Heaths, a fortunate circumstance, as it is scarcely possible to strike its cuttings.

Seedling plants rarely flower till they are three years old.

[359]

CONVOLVULUS ALTHÆOIDES. SILKY-LEAVED CONVOLVULUS.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. campanulata plicata. *Stigmata* 2. *Caps.* 2-locularis: loculis dispermis.

Specific Character and Synonyms.

CONVOLVULUS *althæoides* foliis cordatis sinuatis sericeis: lobis repandis; pedunculis bifloris.
Linn. Syst. Vegetab. ed. 14. Murr. p. 202. Ait. Kew. v. 1. p. 211. var. [Greek: b].

CONVOLVULUS *argenteus* Althææ folio. *Bauh. Pin. p. 295.*

CONVOLVULUS Althææ folio. *Clus. Hist. p. xlix. fig.*

PAPAVER cornutum luteum minus. *Ger. Herb. p. 294. f. 4.*

CONVOLVULUS *argenteus elegantissimus* foliis tenuiter incisis. *Tournef. Inst. 85.*



N°. 359

The name of *Althæoides* and the description of LINNÆUS accord much better with the figure of this plant, as given by CLUSIUS, than with the plant itself as cultivated in our gardens: whether the foliage of our plant becomes

smaller and more finely divided by cultivation; whether it be a distinct species, as MILLER affirms, or a permanent variety, as Mr. AITON makes it, we have not been able satisfactorily to ascertain; the former gives no description of the radical leaves of his *elegantissimus*, and we have not found in any of our gardens a variety different from the one here figured. This species of *Convolvulus*, though cultivated here by J. TRADESCANT in 1656, is far from being common, which is the more surprising, as the plant is easy of culture and productive of flowers at once large and beautiful, and peculiarly interesting from the extreme variableness of its silky foliage, expanding into the most elegant forms imaginable.

Mr. STEVENS, of Camerton-House, near Bath, informs me that it grows abundantly on the mountains near Victri, southeast of Naples, and in the Isle of Capri, mixed with *Convolvulus Cneorum*, *Lithospermum fruticosum*, &c. and in the adjacent islands and continent, forming a beautiful ornament to the shrubs it entwines: CLUSIUS observed it in Spain and Portugal.

It flowers with us in June and July, and is increased without difficulty by parting its roots, which are of the creeping kind, in spring or autumn. It is usually kept in the greenhouse, but will succeed very well in the open border, guarding it against the unusual severity of weather.

[360]

HIBISCUS SPECIOSUS. SUPERB HIBISCUS.

Class and Order.

MONADELPHIA POLYANDRIA.

Generic Character.

Cal. 2-plex: exterior polyphyllus aut multifidus. *Caps.* 5-valvis, 5-locularis: loculis polyspermis ramis 1-spermis.

Specific Character and Synonyms.

HIBISCUS *Speciosus* foliis glabris palmatis: laciniis lanceolatis serratis, caule pedunculis calicibusque lævibus. *Ait. Hort. Kew.* 2. p. 456. *Linn. Syst. Nat. ed. Gmel.* p. 1063.



N^o. 360

Those who "admire Nature's clocks more than her watches," will view, we presume, with some pleasure our representation of this very superb

species of Hibiscus, a native of Carolina, and cultivated here by Dr. JOHN FOTHERGILL, in 1778.

This species is altogether herbaceous, and from a perennial root yearly throws up a stem to the height of many feet, clothed with foliage of a beautiful form and smoothness, and supporting at its summit several flowers, distinguished for their grandeur and richness of colour; these usually blossom in August, and if the plant be kept in the stove, as it most commonly is, are followed by ripe seeds, by which it is most commonly propagated.

In the *Hort. Kew.* it is marked as a greenhouse plant; it may no doubt be preserved in the greenhouse; there is even no impediment to its growing in the open border, if placed in a warm and sheltered situation; and the only motive for keeping it in the stove is its being found to flower there more advantageously, and to ripen its seeds with more certainty.

INDEX.

In which the Latin Names of the Plants contained in the *Tenth Volume* are alphabetically arranged.

Pl.

- [338](#) Allamanda cathartica.
- [343](#) Antholyza Cunonia.
- [339](#) Arum trilobatum.
- [344](#) Aspalathus pedunculata.
- [357](#) Briza maxima.
- [348](#) Calceolaria Fothergillii.
- [333](#) Cerinthe major.
- [327](#) Chrysanthemum indicum.
- [359](#) Convolvulus althæoides.
- [358](#) Erica baccans.
- [356](#) ——— Massoni.
- [342](#) ——— persoluta.
- [350](#) ——— ventricosa.
- [360](#) Hibiscus speciosus.
- [334](#) Hypericum monogynum.
- [325](#) Justicia nasuta.
- [331](#) Kalmia angustifolia.
- [336](#) Lotus hirsutus.
- [353](#) Mahernia incisa.
- [326](#) Mesembryanthemum viridiflorum.
- [354](#) Mimulus aurantiacus.
- [332](#) Oenothera fruticosa.
- [355](#) ——— pumila.
- [352](#) ——— purpurea.

- [347](#) ——— rosea.
- [329](#) Ononis natrix.
- [335](#) ——— rotundifolia.
- [340](#) Polygala Heisteria.
- [345](#) ——— bracteolata.
- [346](#) Protea mellifera.
- [337](#) Prunella grandiflora.
- [351](#) Saxifraga mutata.
- [341](#) Scilla amæna.
- [330](#) Sida Cristata.
- [349](#) Solanum laciniatum.
- [328](#) Trifolium incarnatum.

INDEX.

In which the English Names of the Plants contained in the *Tenth Volume* are alphabetically arranged.

Pl.

- [338](#) Allamanda willow-leaved.
- [343](#) Antholyza scarlet-flowered.
- [339](#) Arum three-lobed.
- [344](#) Aspalathus small-leaved.
- [336](#) Bird's-foot Trefoil hairy.
- [327](#) Chrysanthemum indian.
- [359](#) Convolvulus silky-leaved.
- [326](#) Fig-marigold green-flowered.
- [342](#) Heath blush-flowered.
- [350](#) ——— porcelain.
- [358](#) ——— Arbutus-leaved.
- [356](#) ——— Masson's.
- [360](#) Hibiscus superb.
- [333](#) Honey-wort great.
- [325](#) Justicia dichotomous.
- [331](#) Kalmia narrow-leaved.
- [353](#) Mahernia cut-leaved.
- [340](#) Milkwort heath-leaved.
- [345](#) ——— spear-leaved.
- [354](#) Monkey-flower orange.
- [349](#) Nightshade cut-leaved.
- [355](#) Oenothera dwarf.
- [332](#) ——— shrubby.
- [347](#) ——— rose-coloured.

- [352](#) ——— purple.
 - [346](#) Protea honey-bearing.
 - [357](#) Quaking grass great.
 - [329](#) Rest-harrow yellow-flowered.
 - [335](#) ——— round-leaved.
 - [351](#) Saxifrage saffron-coloured.
 - [337](#) Self-heal great-flowered.
 - [330](#) Sida crested.
 - [348](#) Slipper-wort Fothergill's.
 - [341](#) Squill Byzantine.
 - [334](#) St. John's-wort Chinese.
 - [328](#) Trefoil crimson.
-

INDEX.

In which the *Latin Names* of the Plants contained in the first *Ten Volumes* of the BOTANICAL MAGAZINE are alphabetically arranged.

<i>Vol.</i>	<i>Pag.</i>	
4	134	<i>Adonis vernalis.</i>
9	295	<i>Agrostemma Cœli rosa.</i>
1	24	—— <i>coronaria.</i>
5	173	<i>Aitonia capensis.</i>
10	338	<i>Allamanda cathartica.</i>
7	251	<i>Allium descendens.</i>
4	125	<i>Alstrœmeria Ligtu.</i>
4	139	—— <i>Pelegrina.</i>
4	126	<i>Alyssum deltoideum.</i>
3	101	—— <i>halimifolium.</i>
5	159	—— <i>saxatile.</i>
4	130	—— <i>utriculatum.</i>
7	239	<i>Amaryllis Atamasco.</i>
9	305	—— <i>equestris.</i>
2	47	—— <i>formosissima.</i>
9	290	—— <i>lutea.</i>
9	294	—— <i>sarniensis.</i>
4	129	—— <i>vittata.</i>
5	161	<i>Amygdalus nana.</i>
9	319	<i>Anagallis Monelli.</i>
1	10	<i>Anemone Hepatica.</i>
4	123	—— <i>hortensis.</i>
2	54	—— <i>sylvestris.</i>
6	207	<i>Antirrhinum alpinum.</i>

3	99	—— purpureum.
6	200	—— sparteum.
3	74	—— triste.
9	324	—— triphyllum.
9	318	Anthericum Liliastrum.
10	343	Antholyza Cunonia.
3	108	Anthyllis tetraphylla.
8	280	Apocynum androsæmifolium.
7	246	Aquilegia canadensis.
7	226	Arabis alpina.
7	243	Argemone mexicana.
10	339	Arum trilobatum.
10	344	Aspalathus pedunculata.
6	199	Aster alpinus.
1	33	—— tenellus.
5	180	Azalea nudiflora, var. coccinea.
7	228	Bellis perennis, var. major fl. pl.
8	276	Blitum virgatum.
8	274	Borbonia crenata.
10	357	Briza maxima.
1	34	Browallia elata.
7	217	Buchnera viscosa.
5	174	Buddlea globosa.
5	153	Bulbocodium vernum.
1	17	Cactus flagelliformis.
10	348	Calceolaria Fothergillii.
2	41	—— pinnata.
4	117	Campanula carpatica.
7	252	—— grandiflora.
3	102	—— Speculum.

2	42	<i>Camellia japonica</i> .
9	291	<i>Capparis spinosa</i> .
3	107	<i>Cassia Chamæcrista</i> .
4	131	<i>Catesbæa spinosa</i> .
9	293	<i>Catananche cærulea</i> .
6	210	<i>Celsia linearis</i> .
2	62	<i>Centaurea glastifolia</i> .
3	77	—— <i>montana</i> .
10	333	<i>Cerinth major</i> .
5	166	<i>Cheiranthus maritimus</i> .
6	195	—— <i>mutabilis</i> .
7	233	<i>Chironia baccifera</i> .
2	37	—— <i>frutescens</i> .
10	327	<i>Chrysanthemum indicum</i> .
7	249	<i>Cineraria Amelloides</i> .
2	53	—— <i>lanata</i> .
8	264	<i>Cistus formosus</i> .
2	43	—— <i>incanus</i> .
4	112	<i>Cistus ladaniferus</i> .
2	65	<i>Clematis integrifolia</i> .
3	81	<i>Colutea arborescens</i> .
6	181	—— <i>frutescens</i> .
10	359	<i>Convolvulus althæoides</i> .
9	289	—— <i>linearis</i> .
6	188	—— <i>Nil</i> .
4	113	—— <i>purpureus</i> .
1	27	—— <i>tricolor</i> .
5	156	<i>Coreopsis verticillata</i> .
1	13	<i>Coronilla glauca</i> .
6	185	—— <i>valentina</i> .

8	258	—— varia.
9	321	Cotyledon orbiculata.
1	35	Crepis barbata.
2	45	Crocus vernus.
1	4	Cyclamen Coum.
2	44	—— persicum.
1	7	Cynoglossum Omphalodes.
6	192	Cypripedium acaule.
6	216	—— album.
8	271	Cyrtanthus angustifolius.
5	176	Cytisus Laburnum.
8	255	—— sessilifolius.
5	147	Dais cotinifolia.
9	313	Daphne Cneorum.
6	207	Dianthus barbatus.
2	39	—— Caryophyllus, var.
1	25	—— chinensis.
9	297	—— superbus.
8	273	Diosma uniflora.
7	218	Disandra prostrata.
1	12	Dodecatheon Meadia.
5	170	Draba aizoides.
6	214	Dracocephalum denticulatum.
5	152	Epidendrum cochleatum.
3	76	Epilobium angustissimum.
9	303	Erica ampullacea.
10	358	—— baccans.
7	220	—— cerinthoides.
6	189	—— grandiflora.
1	11	—— herbacea.

10	356	—— Massoni.
10	342	—— persoluta.
10	350	—— ventricosa.
9	310	Erinus alpinus.
1	5	Erythronium Dens Canis.
8	261	Erodium incarnatum.
7	241	Fagonia cretica.
4	144	Ferraria undulata.
2	63	Fragaria monophylla.
6	194	Fritillaria imperialis.
3	97	Fuchsia coccinea.
7	232	Fumaria cava.
5	179	—— glauca.
7	231	Fumaria solida.
9	314	Genista triquetra.
2	52	Gentiana acaulis.
6	203	Geranium angulatum.
6	206	—— anemonefolium.
2	56	—— lanceolatum.
1	20	—— peltatum.
2	55	—— striatum.
1	18	—— Reichardi.
3	95	—— Radula.
3	86	Gladiolus communis.
4	135	—— cardinalis.
8	272	—— tristis.
8	263	Glycine bimaculata.
8	270	—— coccinea.
8	268	—— rubicunda.
9	300	Gnaphalium eximium.

8	287	<i>Goodenia lævigata</i> .
3	90	<i>Gorteria rigens</i> .
8	282	<i>Hedysarum obscurum</i> .
7	227	<i>Helianthus multiflorus</i> .
4	141	<i>Heliotropium peruvianum</i> .
1	3	<i>Helleborus hyemalis</i> .
2	72	—— <i>lividus</i> .
1	8	—— <i>niger</i> .
1	19	<i>Hemerocallis flava</i> .
2	64	—— <i>fulva</i> .
9	299	<i>Hermannia alnifolia</i> .
9	307	—— <i>althæifolia</i> .
9	304	—— <i>lavandulifolia</i> .
5	158	<i>Hibiscus Rosa sinensis</i> .
10	360	—— <i>speciosus</i> .
3	83	—— <i>syriacus</i> .
6	209	—— <i>Trionum</i> .
5	157	<i>Hyacinthus botryoides</i> .
4	133	—— <i>comosus</i> .
4	122	—— <i>racemosus</i> .
3	87	<i>Hyoscyamus aureus</i> .
4	137	<i>Hypericum balearicum</i> .
5	146	—— <i>calycinum</i> .
5	178	—— <i>Coris</i> .
10	334	—— <i>monogynum</i> .
4	124	<i>Iberis gibraltarica</i> .
3	106	—— <i>umbellata</i> .
6	198	<i>Indigofera candicans</i> .
7	221	<i>Ipomœa coccinea</i> .
7	244	—— <i>Quamoclit</i> .

2	61	<i>Iris ochroleuca.</i>
1	1	—— <i>persica.</i>
1	9	—— <i>pumila.</i>
5	168	—— <i>pavonia.</i>
2	50	—— <i>sibirica.</i>
2	58	—— <i>spuria.</i>
3	91	<i>Iris susiana.</i>
6	187	—— <i>sambucina.</i>
1	16	—— <i>variegata.</i>
1	21	—— <i>versicolor.</i>
8	265	<i>Ixia Bulbocodium.</i>
5	171	—— <i>chinensis.</i>
6	184	—— <i>crocata.</i>
4	127	—— <i>flexuosa.</i>
8	256	—— <i>longiflora.</i>
5	169	<i>Ixora coccinea.</i>
1	31	<i>Jasminum officinale.</i>
8	285	—— <i>odoratissimum.</i>
10	325	<i>Justicia nasuta.</i>
10	331	<i>Kalmia angustifolia.</i>
5	177	—— <i>glauca.</i>
4	138	—— <i>hirsuta.</i>
5	175	—— <i>latifolia.</i>
3	82	<i>Lachenalia tricolor.</i>
5	172	<i>Lamium Orvala.</i>
3	96	<i>Lantana aculeata.</i>
8	253	<i>Lathyrus articulatus.</i>
2	60	—— <i>odoratus.</i>
4	115	—— <i>sativus.</i>
3	100	—— <i>tingitanus.</i>

4	111	—— tuberosus.
4	109	Lavatera trimestris.
2	46	Leucojum vernal.
1	36	Lilium bulbiferum.
1	30	—— chalcedonicum.
8	259	—— Catesbæi.
8	278	—— candidum.
4	116	Limodorum tuberosum.
7	234	Linum arboreum.
9	312	—— flavum.
8	275	Liriodendron Tulipifera.
9	325	Lobelia Cardinalis.
7	225	—— surinamensis.
8	254	Lopezia racemosa.
10	336	Lotus hirsutus.
3	79	—— jacobæus.
5	151	—— tetragonolobus.
4	140	Lupinus luteus.
6	202	—— perennis.
7	223	Lychnis coronata.
8	257	—— chalcedonica.
3	104	Lysimachia bulbifera.
10	353	Mahernia incisa.
8	277	—— pinnata.
9	322	Manulea tomentosa.
9	301	Melianthus minor.
6	208	Melissa grandiflora.
8	262	Mesembry ^m . aureum.
2	70	—— barbatum.
2	59	—— bicolorum.

1	32	—— dolabriforme.
2	67	—— pinnatifidum.
10	326	—— viridiflorum.
8	260	<i>Metrosideros citrina</i> .
7	219	<i>Michauxia campanuloides</i> .
9	302	<i>Mimosa myrtifolia</i> .
4	110	—— verticillata.
10	354	<i>Mimulus aurantiacus</i> .
8	283	—— ringens.
5	145	<i>Monarda fistulosa</i> , var.
3	73	<i>Monsonia speciosa</i> .
7	250	<i>Myrtus tomentosa</i> .
6	193	<i>Narcissus angustifolius</i> .
6	197	—— biflorus.
3	88	—— <i>Bulbocodium</i> .
4	121	—— incomparabilis.
1	15	—— Jonquilla.
2	51	—— major.
1	6	—— minor.
3	78	—— odorus.
2	48	—— triandrus.
1	22	<i>Nigella damascena</i> .
10	332	<i>Oenothera fruticosa</i> .
10	347	—— rosea.
10	352	—— purpurea.
10	355	—— pumila.
9	317	<i>Ononis fruticosa</i> .
10	329	—— <i>Natrix</i> .
10	335	—— rotundifolia.
9	298	<i>Origanum Dictamnus</i> .

6	190	<i>Ornithogalum aureum</i> .
8	269	—— <i>nutans</i> .
9	306	<i>Othonna pectinata</i> .
7	237	<i>Oxalis caprina</i> .
5	155	—— <i>versicolor</i> .
2	57	<i>Papaver orientale</i> .
9	292	<i>Passerina grandiflora</i> .
2	66	<i>Passiflora alata</i> .
1	28	—— <i>cærulea</i> .
8	288	—— <i>ciliata</i> .
3	103	<i>Pelargonium acetosum</i> .
5	148	—— <i>betulinum</i> .
6	201	—— <i>bicolor</i> .
5	165	—— <i>cordifolium</i> .
9	315	—— <i>ceratophyllum</i> .
9	309	—— <i>echinatum</i> .
4	143	—— <i>glutinosum</i> .
4	136	—— <i>tetragonum</i> .
7	240	—— <i>tricolor</i> .
5	163	<i>Phlox divaricata</i> .
7	224	<i>Phylla ericoides</i> .
7	230	<i>Plumbago rosea</i> .
8	279	<i>Plumeria rubra</i> .
10	345	<i>Polygala bracteolata</i> .
9	316	—— <i>chamæbuxus</i> .
10	340	—— <i>Heisteria</i> .
6	213	<i>Polygonum orientale</i> .
8	286	<i>Portlandia grandiflora</i> .
3	75	<i>Potentilla grandiflora</i> .
7	229	<i>Primula acaulis</i> , fl. pl. carn.

6	191	—— marginata.
1	14	—— villosa.
10	346	<i>Protea mellifera</i> .
10	337	<i>Prunella grandiflora</i> .
5	160	<i>Pulmonaria virginica</i> .
8	267	<i>Pyrus spectabilis</i> .
6	204	<i>Ranunculus aconitifolius</i> .
6	215	—— acris, flore pleno.
8	266	—— amplexicaulis.
5	164	—— gramineus.
1	29	<i>Reseda odorata</i> .
9	311	<i>Robinia hispida</i> .
2	69	—— <i>Rosa muscosa</i> .
8	284	—— <i>semperflorens</i> .
4	132	<i>Rubus arcticus</i> .
9	323	—— odoratus.
1	2	<i>Rudbeckia purpurea</i> .
6	182	<i>Salvia aurea</i> .
5	162	<i>Sanguinaria canadensis</i> .
5	154	<i>Saponaria Ocymoides</i> .
6	196	<i>Saxifraga crassifolia</i> .
10	351	—— mutata.
3	92	—— sarmentosa.
7	247	<i>Scabiosa atropurpurea</i> .
10	341	<i>Scilla amæna</i> .
4	128	—— campanulata.
4	142	<i>Scorzonera tingitana</i> .
4	118	<i>Sedum Anacampseros</i> .
6	211	—— populifolium.
6	186	<i>Selago ovata</i> .

2	68	<i>Sempervivum arachnoideum.</i>
3	93	—— <i>monanthes.</i>
9	296	<i>Sempervivum tortuosum.</i>
7	238	<i>Senecio elegans.</i>
10	230	<i>Sida cristata.</i>
4	114	<i>Silene pendula.</i>
3	94	<i>Sisyrinchium iridioides.</i>
5	167	<i>Sophora tetraptera.</i>
10	349	<i>Solanum laciniatum.</i>
2	49	<i>Soldanella alpina.</i>
3	85	<i>Spartium junceum.</i>
3	80	<i>Spigelia marilandica.</i>
1	26	<i>Stapelia variegata.</i>
2	71	<i>Statice sinuata.</i>
4	119	<i>Strelitzia Reginae.</i>
7	222	<i>Struthiola erecta.</i>
6	183	<i>Syringa vulgaris.</i>
5	150	<i>Tagetes patula.</i>
6	212	<i>Tanacetum flabelliforme.</i>
7	245	<i>Teucrium latifolium.</i>
3	105	<i>Tradescantia virginica.</i>
10	328	<i>Trifolium incarnatum.</i>
2	40	<i>Trillium sessile.</i>
7	235	<i>Trollius asiaticus.</i>
1	23	<i>Tropæolum majus.</i>
3	98	—— <i>minus.</i>
8	281	<i>Turnera angustifolia.</i>
3	84	<i>Tussilago alpina.</i>
7	236	<i>Verbascum Myconi.</i>
9	308	<i>Verbena Aubletia.</i>

7	242	Veronica decussata.
2	38	Viburnum Tinus.
7	248	Vinca rosea.
3	89	Viola pedata.
5	149	Zinnia multiflora.

INDEX.

In which the *English Names* of the Plants contained in the first *Ten Volumes* of the BOTANICAL MAGAZINE are alphabetically arranged.

<i>Vol.</i>	<i>Pag.</i>	
4	134	Adonis spring.
5	173	Aitonia cape.
10	338	Allamanda willow-leaved.
5	161	Almond dwarf.
4	125	Alstrœmeria striped-flowered.
4	139	—— spotted flowered.
4	130	Alyssum bladder-podded.
4	126	—— purple.
3	101	—— sweet.
5	159	—— yellow.
9	305	Amaryllis Barbadoes.
9	294	—— Guernsey.
2	47	—— Jacobean.
4	129	—— superb.
9	290	—— yellow.
2	54	Anemone Snow-drop.
4	123	—— star.
9	318	Anthericum Savoy.
10	343	Antholyza scarlet-flowered.
8	267	Apple tree Chinese.
5	174	Archangel balm-leav'd.
10	339	Arum three-lobed
10	344	Aspalathus small-leaved.
6	199	Aster alpine.

1	33	—— bristly-leaved.
5	180	Azalea scarlet.
6	208	Balm great-flowered.
4	117	Bell-flower Carpatian.
7	252	—— great-flowered.
4	113	Bindweed purple.
6	181	Bladder-senna scarlet.
3	81	—— common.
8	276	Blite strawberry.
3	77	Blue-bottle greater.
8	274	Borbonia heart-leaved.
4	132	Bramble dwarf.
3	85	Broom Spanish.
1	34	Browallia tall.
7	217	Buchnera clammy.
5	174	Buddlea round-headed.
5	153	Bulbocodium vernal.
2	42	Camellia rose.
4	124	Candy-tuft Gibraltar.
3	106	—— purple.
9	291	Caper Shrub.
3	107	Cassia dwarf.
9	293	Catananche blue.
4	114	Catchfly pendulous.
4	131	Catesbæa thorny.
6	210	Celsia linear-leaved.
2	43	Centaurea woad-leaved.
1	17	Cereus creeping.
7	233	Chironia berry-bearing.
2	37	—— shrubby.

10	327	Chrysanthemum indian.
7	249	Cineraria blue-flowered.
2	53	—— woolly.
8	264	Cistus beautiful.
4	112	—— gum.
2	43	—— hoary, or rose.
2	65	Clematis, or Virgin's-bower, entire-leaved.
1	24	Cockle rose.
9	295	—— smooth-leaved.
3	84	Colts-foot alpine.
7	246	Columbine Canadian.
6	188	Convolvulus azure.
9	289	—— narrow-leaved.
1	27	—— small.
10	359	—— silky-leaved.
5	156	Coreopsis whorled.
3	86	Corn-flag common.
8	272	—— square-leaved.
4	135	—— superb.
8	258	Coronilla purple.
6	185	—— rue-leaved.
1	13	—— sea-green.
6	203	Crane's-bill angular-stalked.
5	148	—— birch-leaved.
4	143	—— clammy.
8	261	—— flesh-coloured.
5	165	—— heart-leaved.
9	315	—— horn leaved.
3	103	—— sorrel.
6	201	—— two-coloured.

7	240	—— three-coloured.
1	35	Crepis bearded.
2	45	Crocus spring.
5	164	Crow-foot grass-leaved.
6	204	—— mountain.
8	266	—— plantain-leaved.
6	215	—— upright double.
6	194	Crown imperial.
9	300	Cudweed giant.
1	4	Cyclamen round-leaved.
2	44	—— Persian.
8	271	Cyrtanthus narrow-leaved.
8	255	Cytisus common.
2	51	Daffodil great.
1	6	—— lesser.
4	121	—— peerless.
2	48	—— reflexed.
5	147	Dais cotinus-leaved.
7	228	Daisy great double.
9	313	Daphne trailing.
2	64	Day-lily tawny.
1	19	—— yellow.
8	273	Diosma one-flowered.
7	218	Disandra trailing.
9	298	Dittany of Crete.
1	12	Dodecatheon Mead's.
1	5	Dog's tooth.
8	280	Dog's-bane tutsan-leav'd.
5	170	Draba sengreen.
6	214	Dragon's-head toothed.

5	152	Epidendrum two-leaved.
9	310	Erinus alpine.
7	241	Fagonia cretan.
1	22	Fennel-flower garden.
4	144	Ferraria curled.
2	70	Fig-marigold bearded.
8	262	—— golden.
10	326	—— green-flowered.
1	32	—— hatchet-leaved.
2	67	—— jagged-leaved.
2	59	—— two-coloured.
7	234	Flax tree.
9	312	—— yellow.
2	39	Franklin's Tartar.
3	97	Fuchsia scarlet.
5	179	Fumitory glaucous.
7	232	—— hollow-rooted.
7	231	—— solid-rooted.
7	251	Garlick purple-headed.
9	314	Genista triangular-stalked.
2	52	Gentian large-flowered.
6	200	Geranium anemone-leaved.
1	18	—— dwarf.
1	20	—— ivy-leaved.
9	309	—— prickly-stalked.
3	95	—— rasp-leaved.
2	55	—— striped.
2	56	—— spear-leaved.
4	136	—— square-stalked.
7	245	Germander broad-leaved shrubby.

7	235	Globe-flower Asiatic.
8	268	Glycine dingy-flowered.
8	263	—— purple.
8	270	—— scarlet.
8	287	Goodenia smooth.
3	90	Gorteria rigid-leaved.
10	358	Heath Arbutus-leaved.
10	342	—— blush-flowered.
9	303	—— flask.
6	189	—— great-flowered.
1	11	—— herbaceous.
7	220	—— honeywort-flowered.
10	356	—— Masson's.
10	350	—— porcelain.
8	282	Hedysarum creeping-rooted.
1	8	Hellebore black.
2	72	—— livid or purple.
1	3	—— winter.
3	87	Henbane golden-flowered.
1	10	Hepatica.
9	299	Hermannia alder-leaved.
9	304	—— lavender-leaved.
9	307	—— marshmallow-leaved.
6	209	Hibiscus bladder.
5	158	—— China rose.
3	83	—— Syrian.
10	360	—— superb.
10	333	Honey-wort great.
2	68	House-leek cobweb.
3	93	—— dwarf.

9	296	—— gouty.
5	157	Hyacinth grape.
4	122	—— starch.
4	133	—— two-coloured.
1	23	Indian-cress greater.
3	98	—— small.
6	198	Indigo white-leaved.
1	15	—— Jonquil common.
3	78	—— great.
7	221	Ipomœa scarlet.
7	244	—— winged-leaved.
3	91	Iris chalcedonian.
1	9	—— dwarf.
6	187	—— elder-scented.
1	1	—— persian.
1	21	—— particoloured.
5	168	—— peacock.
2	50	—— Siberian.
2	58	—— spurious.
2	61	—— tall.
1	16	—— variegated.
4	127	Ixia bending-stalked.
5	171	—— Chinese.
8	265	—— crocus-leaved.
8	256	—— long-flowered.
6	184	—— saffron-coloured.
5	169	Ixora scarlet.
1	31	Jasmine common.
8	285	—— sweet.
10	325	Justicia dichotomous.

5	175	Kalmia broad-leaved.
5	177	—— glaucous.
4	138	—— hairy.
10	331	—— narrow-leaved.
5	176	Laburnum common.
3	82	Lachenalia three-coloured.
3	108	Ladies-finger four-leaved.
6	192	Ladies-slipper two-leaved.
6	216	—— white-petal'd.
3	96	Lantana prickly.
4	115	Lathyrus blue-flowered.
8	253	—— jointed-podded.
4	111	—— tuberous.
4	109	Lavatera annual.
2	38	Laurustinus common.
7	230	Lead-wort rose-coloured.
6	183	Lilac common.
7	239	Lily Atamasco.
8	259	—— Catesby's.
1	30	—— chalcedonian.
1	36	—— orange.
8	278	—— white.
4	116	Limodorum tuberous-rooted.
7	225	Lobelia shrubby.
9	320	—— scarlet.
8	254	Lopezia Mexican.
3	104	Loosestrife bulb-bearing.
3	79	Lotus black-flowered.
5	151	—— winged.
10	336	—— hairy.

5	160	Lungwort Virginian.
6	202	Lupine perennial.
4	140	—— yellow.
5	163	Lychnidea early-flowering.
7	223	Lychnis Chinese.
8	257	—— scarlet.
10	353	Mahernia cut-leaved.
8	277	—— winged.
9	322	Manulea woolly.
5	150	Marigold French.
9	301	Melianthus small.
8	260	Metrosideros harsh-leaved.
7	219	Michauxia rough-leaved.
1	29	Mignonet.
9	316	Milkwort box-leaved.
10	340	—— heath-leaved.
10	345	—— spear-leaved.
9	302	Mimosa myrtle-leaved.
4	110	—— whorled-leaved.
5	145	Monarda crimson.
8	283	Monkey-flower narr. leaved.
10	354	—— orange.
3	73	Monsonia large-flowered.
7	236	Mullein borage-leaved.
7	250	Myrtle woolly-leaved.
3	88	Narcissus hoop-petticoat.
6	193	—— narrow-leaved.
6	197	—— two-flowered.
1	7	Navel-wort blue.
9	321	—— round-leaved.

10	349	Nightshade cut-leaved.
10	355	Oenothera dwarf.
10	352	—— purple.
10	347	—— rose-coloured.
10	332	—— shrubby.
6	190	Ornithogalum golden.
4	118	Orpine evergreen.
9	306	Othonna wormwood-leaved.
9	292	Passerina great-flowered.
1	28	Passion-flower common.
8	288	—— fringed-leaved.
2	66	—— winged.
2	60	Pea sweet.
3	100	—— Tangier.
7	248	Periwinkle Madagascar.
6	213	Persicaria tall.
7	224	Phylica heath-leaved.
9	319	Pimpernel Italian.
1	25	Pink China, or Indian.
9	297	—— superb.
8	279	Plumeria red.
2	57	Poppy eastern.
7	243	—— prickly.
8	286	Portlandia great-flowered.
3	75	Potentilla large-flowered.
7	229	Primrose lilac double.
1	14	Primula mountain.
6	191	—— silver-edged.
10	346	Protea honey-bearing.
5	162	Puccoon Canada.

10	357	Quaking-grass great.
7	238	Rag-wort purple.
9	323	Raspberry flowering.
10	335	Rest-harrow round-leaved.
9	317	—— shrubby.
10	329	—— yellow-flowered.
1	29	Reseda sweet-scented.
8	284	Rose ever-blowing.
2	69	—— moss.
9	311	Robinia rough-stalked.
1	2	Rudbeckia purple.
6	182	Sage golden.
6	196	Saxifrage oval-leaved.
10	351	—— saffron-coloured.
3	92	—— strawberry.
7	247	Scabious sweet.
4	142	Scorzonera Tangier.
6	186	Selago oval-headed.
10	337	Self-heal great-flowered.
10	330	Sida crested.
3	94	Sisyrinchium Iris-leaved.
10	334	St. John's-wort Chinese.
5	178	—— heath-leaved.
5	146	—— large-flowered.
4	137	—— warty.
10	348	Slipper-wort Fothergill's.
2	41	—— pinnated.
2	46	Snow-flake spring.
5	154	Soap-wort basil.
2	49	Soldanella alpine.

5	167	Sophora winged-podded.
7	242	Speedwell cross-leaved.
3	80	Spigelia Maryland.
4	128	Squill bell-flowered.
10	341	—— Byzantine.
1	26	Stapelia variegated.
8	269	Star of Bethlehem Neapolit ⁿ .
5	166	Stock Mediterranean.
6	211	Stone-crop poplar-leaved.
2	63	Strawberry one-leaved.
4	119	Strelitzia Canna-leaved.
7	222	Struthiola smooth.
7	227	Sun-flower perennial.
6	207	Sweet william.
6	212	Tansey fan-leaved.
2	71	Thrift purple-cup'd.
6	205	Toad-flax alpine.
6	200	—— branching.
3	74	—— black-flowered.
3	99	—— purple.
9	324	—— three-leaved.
3	105	Tradescantia virginian.
10	328	Trefoil crimson.
2	40	Trillium sessile.
8	275	Tulip-tree common.
8	281	Turnera narrow-leaved.
4	141	Turnsole peruvian.
9	308	Vervain rose.
3	89	Violet cut-leaved.
7	226	Wall-cress alpine.

6	195	Wall-flower changeable.
3	76	Willow-herb narrowest-leav ^d .
5	155	Wood-Sorrel striped-flowered
7	237	—— goat's-foot.
5	149	Zinnia many-flowered.

INDEX

To the *Hardy Trees* contained in the first *Ten Volumes* of the *Botanical Magazine*.

<i>Vol.</i>	<i>Pag.</i>	
5	176	Cytisus Laburnum.
1	31	Jasminum officinale.
8	275	Liriodendron Tulipifera.
1	28	Passiflora cærulea.
8	267	Pyrus spectabilis.
6	183	Syringa vulgaris.

INDEX

To the *Hardy Shrubs* contained in the first *Ten Volumes* of the *Botanical Magazine*.

<i>Vol.</i>	<i>Pag.</i>	
5	161	<i>Amygdalus nana</i> .
5	180	<i>Azalea nudiflora</i> , var. <i>coccin</i> .
4	112	<i>Cistus ladaniferus</i> .
3	81	<i>Colutea arborescens</i> .
8	255	<i>Cytisus sessilifolius</i> .
9	313	<i>Daphne Cneorum</i> .
1	11	<i>Erica herbacea</i> .
9	314	<i>Genista triquetra</i> .
3	83	<i>Hibiscus syriacus</i> .
10	331	<i>Kalmia angustifolia</i> .
5	177	—— <i>glauca</i> .
4	138	—— <i>hirsuta</i> .
5	175	—— <i>latifolia</i> .
9	317	<i>Ononis fruticosa</i> .
10	335	—— <i>rotundifolia</i> .
9	316	<i>Polygala Chamæbuxus</i> .
9	311	<i>Robinia hispida</i> .
2	69	<i>Rosa muscosa</i> .
9	323	<i>Rubus odoratus</i> .
3	85	<i>Spartium junceum</i> .
2	38	<i>Viburnum Tinus</i> .

INDEX

To the *Hardy Herbaceous Perennial Plants* contained in the first *Ten Volumes* of the *Botanical Magazine*.

<i>Vol.</i>	<i>Pag.</i>	
4	134	<i>Adonis vernalis</i> .
1	24	<i>Agrostemma coronaria</i> .
7	251	<i>Allium descendens</i> .
4	126	<i>Alyssum deltoideum</i> .
5	159	—— <i>saxatile</i> .
4	130	—— <i>utriculatum</i> .
9	290	<i>Amaryllis lutea</i> .
1	10	<i>Anemone Hepatica</i> .
4	123	—— <i>hortensis</i> .
2	54	—— <i>sylvestris</i> .
9	318	<i>Anthericum Liliastrum</i> .
6	207	<i>Antirrhinum alpinum</i> .
3	99	—— <i>purpureum</i> .
8	280	<i>Apocynum androsæmifolium</i> .
7	246	<i>Aquilegia canadensis</i> .
7	226	<i>Arabis alpina</i> .
6	199	<i>Aster alpinus</i> .
7	228	<i>Bellis perennis</i> , var. <i>maj</i> .
5	153	<i>Bulbocodium verum</i> .
4	117	<i>Campanula carpatica</i> .
7	252	—— <i>grandiflora</i> .
9	293	<i>Catananche cærulea</i> .
2	62	<i>Centaurea glastifolia</i> .
3	77	—— <i>montana</i> .

2	65	<i>Clematis integrifolia</i> .
5	156	<i>Coreopsis verticillata</i> .
8	258	<i>Coronilla varia</i> .
2	45	<i>Crocus vernus</i> .
1	4	<i>Cyclamen Coum</i> .
1	7	<i>Cynoglossum Omphalodes</i> .
6	216	<i>Cypripedium album</i> .
6	192	—— <i>acaule</i> .
6	207	<i>Dianthus barbatus</i> .
2	39	—— <i>Caryophyllus</i> , var.
1	12	<i>Dodecatheon Meadia</i> .
5	170	<i>Draba aizoides</i> .
6	214	<i>Dracocephalum denticulatum</i> .
3	76	<i>Epilobium angustissimum</i> .
9	310	<i>Erinus alpinus</i> .
1	5	<i>Erythronium Dens Canis</i> .
2	63	<i>Fragaria monophylla</i> .
6	194	<i>Fritillaria imperialis</i> .
7	232	<i>Fumaria cava</i> .
7	231	—— <i>solida</i> .
2	52	—— <i>Gentiana acaulis</i> .
6	203	<i>Geranium angulatum</i> .
2	55	—— <i>striatum</i> .
3	86	<i>Gladiolus communis</i> .
8	282	<i>Hedysarum obscurum</i> .
7	227	<i>Helianthus multiflorus</i> .
1	3	<i>Helleborus hyemalis</i> .
2	72	—— <i>lividus</i> .
1	8	—— <i>niger</i> .
1	19	<i>Hemerocallis flava</i> .

2	64	—— fulva.
5	157	Hyacinthus botryoides.
4	133	—— comosus.
4	122	—— racemosus.
5	146	Hypericum calycinum.
2	61	Iris ochroleuca.
1	1	—— persica.
1	9	—— pumila.
6	187	—— sambucina.
3	91	—— susiana.
2	50	—— sibirica.
2	58	—— spuria.
1	16	—— variegata.
1	21	—— versicolor.
8	265	Ixia Bulbocodium.
5	171	—— chinensis.
5	172	Lamium Orvala.
4	111	Lathyrus tuberosus.
2	46	Leucojum vernal.
1	36	Lilium bulbiferum.
8	278	—— candidum.
1	30	—— chalcedonicum.
9	312	Linum flavum.
9	320	Lobelia cardinalis.
6	202	Lupinus perennis.
8	257	Lychnis chalcedonica.
3	104	Lysimachia bulbifera.
6	208	Melissa grandiflora.
8	283	Mimulus ringens.
5	145	Monarda fistulosa, var.

6	193	<i>Narcissus angustifolius</i> .
6	197	—— <i>biflorus</i> .
3	88	—— <i>Bulbocodium</i> .
4	121	—— <i>incomparabilis</i> .
1	15	—— <i>Jonquilla</i> .
2	51	—— <i>major</i> .
1	6	—— <i>minor</i> .
3	78	—— <i>odorus</i> .
2	48	—— <i>triandrus</i> .
10	332	<i>Oenothera fruticosa</i> .
10	355	—— <i>pumila</i> .
8	269	<i>Ornithogalum nutans</i> .
2	57	<i>Papaver orientale</i> .
5	163	<i>Phlox divaricata</i> .
3	75	<i>Potentilla grandiflora</i> .
7	229	<i>Primula acaulis</i> , v. <i>carnea</i> .
6	191	—— <i>marginata</i> .
1	14	—— <i>villosa</i> .
10	337	<i>Prunella grandiflora</i> .
5	160	<i>Pulmonaria virginica</i> .
6	215	<i>Ranunculus acris</i> , v. <i>pl</i> .
6	204	—— <i>aconitifol. pl</i> .
8	266	—— <i>amplexicaulis</i> .
5	164	—— <i>gramineus</i> .
4	132	<i>Rubus arcticus</i> .
1	2	<i>Rudbeckia purpurea</i> .
5	162	<i>Sanguinaria canadensis</i> .
5	154	<i>Saponaria Ocymoides</i> .
6	190	<i>Saxifraga crassifolia</i> .
10	351	—— <i>mutata</i> .

3	92	—— sarmentosa.
10	341	Scilla amæna.
4	128	—— campanulata.
4	118	Sedum Anacampseros.
6	211	—— populifolium.
2	68	Sempervivum arachnoideum.
2	49	Soldanella alpina.
3	80	Spigelia marilandica.
3	105	Tradescantia virginica.
2	40	Trillium sessile.
7	235	Trollius asiaticus.
3	84	Tussilago alpina.
7	236	Verbascum Myconi.
3	89	Viola pedata.

INDEX

To the *Annual* and *Biennial Plants* (or, if not strictly such, requiring to be frequently renewed from Seed) usually cultivated in the open Borders, contained in the first *Ten Volumes* of the *Botanical Magazine*.

<i>Vol.</i>	<i>Pag.</i>	
9	295	<i>Agrostemma Cœli rosa.</i>
3	101	<i>Alyssum halimifolium.</i>
3	108	<i>Anthyllis tetraphylla.</i>
6	200	<i>Antirrhinum sparteum.</i>
9	324	<i>Antirrhinum triphyllum.</i>
7	243	<i>Argemone mexicana.</i>
8	276	<i>Blitum virgatum.</i>
10	357	<i>Briza maxima.</i>
3	102	<i>Campanula Speculum.</i>
10	333	<i>Cerinthe major.</i>
5	166	<i>Cheiranthus maritimus.</i>
6	188	<i>Convolvulus Nil.</i>
4	113	—— <i>purpureus.</i>
1	27	—— <i>tricolor.</i>
1	35	<i>Crepis barbata.</i>
1	25	<i>Dianthus chinensis.</i>
9	297	—— <i>superbus.</i>
5	179	<i>Fumaria glauca.</i>
6	209	<i>Hibiscus Trionum.</i>
3	106	<i>Iberis umbellata.</i>
7	221	<i>Ipomœa coccinea.</i>
8	253	<i>Lathyrus articulatus.</i>
2	60	—— <i>odoratus.</i>

4	115	—— sativus.
3	100	—— tingitanus.
4	109	Lavatera trimestris.
5	151	Lotus tetragonolobus.
4	140	Lupinus luteus.
1	22	Nigella damascena.
10	352	Oenothera purpurea.
10	347	—— rosea.
6	213	Polygonum orientale.
1	29	Reseda odorata.
7	247	Scabiosa atropurpurea.
4	142	Scorzonera tingitana.
7	238	Senecio elegans.
4	114	Silene pendula.
5	150	Tagetes patula.
10	328	Trifolium incarnatum.
1	23	Tropæolum majus.
3	98	—— minus.
5	149	Zinnia multiflora.

INDEX

To the *Greenhouse Plants* contained in the first *Ten Volumes* of the *Botanical Magazine*.

<i>Vol.</i>	<i>Pag.</i>	
5	173	Aitonia capensis.
4	139	Alstrœmeria Pelegrina.
7	239	Amaryllis Atamasco.
9	294	—— sarniensis.
9	319	Anagellis Monelli.
3	74	Antirrhinum triste.
10	343	Antholyza Cunonia.
10	344	Aspalathus pedunculata.
1	33	Aster tenellus.
8	274	Borbonia crenata.
7	217	Buchnera capensis.
5	174	Buddleja globosa.
1	17	Cactus flagelliformis.
10	348	Calceolaria Fothergillii.
2	42	Camellia japonica.
9	291	Capparis spinosa.
6	210	Celsia linearis.
6	195	Cheiranthus mutabilis.
7	233	Chironia baccifera.
2	37	—— frutescens.
10	327	Chrysanthemum indicum.
7	249	Cineraria Amelloides.
2	53	—— lanata.
8	264	Cistus formosus.

2	43	—— incanus.
6	181	<i>Colutea frutescens</i> .
10	359	<i>Convolvulus althæoides</i> .
9	289	—— <i>linearis</i> .
1	13	<i>Coronilla glauca</i> .
6	185	—— <i>valentina</i> .
9	321	<i>Cotyledon orbiculata</i> .
2	44	<i>Cyclamen persicum</i> .
8	271	<i>Cyrtanthus angustifolius</i> .
5	147	<i>Dais cotinifolia</i> .
8	273	<i>Diosma uniflora</i> .
7	218	<i>Disandra preferata</i> .
9	303	<i>Erica ampullacca</i> .
10	358	—— <i>baccans</i> .
7	220	—— <i>cerinthoides</i> .
6	189	—— <i>grandiflora</i> .
10	356	—— <i>Massoni</i> .
10	342	—— <i>persoluta</i> .
10	350	—— <i>ventricosa</i> .
8	261	<i>Erodium incarnatum</i> .
7	241	<i>Fagonia cretica</i> .
3	97	<i>Fuchsia coccinea</i> .
6	206	<i>Geranium anemonefolium</i> .
2	56	—— <i>lanceolatum</i> .
1	20	—— <i>peltatum</i> .
1	18	—— <i>Reichardi</i> .
3	95	—— <i>Radula</i> .
4	135	<i>Gladiolus cardinalis</i> .
8	272	—— <i>tristis</i> .
8	263	<i>Glycine bimaculata</i> .

8	270	—— coccinea.
8	268	—— rubicunda.
9	300	Gnaphalium eximium.
8	287	Goodenia lævigata.
3	90	Gorteria rigens.
9	299	Hermannia alnifolia.
9	307	—— althæifolia.
9	304	—— lavandulifolia.
3	87	Hyoscyamus aureus.
4	137	Hypericum balearicum.
5	178	—— Coris.
10	334	—— monogynum.
8	285	Jasminum odoratissimum.
4	124	Iberis gibraltarica.
6	198	Indigofera candicans.
5	168	Iris pavonia.
6	184	Ixia crocata.
4	127	—— flexuosa.
8	256	—— longiflora.
3	82	Lachenalia tricolor.
8	259	Lilium Catesbæi.
7	234	Linum arboreum.
8	254	Lopezia racemosa.
10	336	Lotus hirsutus.
3	79	—— Jacobæus.
7	223	Lychnis coronata.
10	353	Mahernia incisa.
8	277	—— pinnata.
9	322	Manulea tomentosa.
9	301	Melianthus minor.

8	262	<i>Mesembrym aureum.</i>
2	70	—— <i>barbatum.</i>
2	59	—— <i>bicolorum.</i>
1	32	—— <i>dolabriforme.</i>
2	67	—— <i>pinnatifidum.</i>
10	326	—— <i>viridiflorum.</i>
8	260	<i>Metrosideros citrina.</i>
7	219	<i>Michauxia campanuloides.</i>
9	302	<i>Mimosa myrtifolia.</i>
4	110	—— <i>verticillata.</i>
10	354	<i>Mimulus aurantiacus.</i>
3	73	<i>Monsonia speciosa.</i>
10	329	<i>Ononis Natrux.</i>
6	190	<i>Ornithogalum aureum.</i>
9	298	<i>Origanum Dictamnus.</i>
9	306	<i>Othonna pectinata.</i>
7	237	<i>Oxalis caprina.</i>
5	155	—— <i>versicolor.</i>
9	292	<i>Passerina grandiflora.</i>
3	103	<i>Pelargonium acetosum.</i>
5	148	—— <i>betulinum.</i>
6	201	—— <i>bicolor.</i>
5	165	—— <i>cordifolium.</i>
9	315	—— <i>ceratophyllum.</i>
9	309	—— <i>echinatum.</i>
4	143	—— <i>glutinosum.</i>
4	136	—— <i>tetragonum.</i>
7	240	—— <i>tricolor.</i>
7	224	<i>Phylla ericoides.</i>
10	345	<i>Polygala bractæolata.</i>

10	340	—— Heisteria.
10	346	Protea mellifera.
8	284	Rosa semperflorens.
6	182	Salvia aurea.
6	186	Selago ovata.
3	93	Sempervivum monanthes.
9	296	—— tortuosum.
10	330	Sida cristata.
3	94	Sisyrinchium Iridioides.
10	349	Solanum laciniatum.
5	167	Sophora tetraptera.
2	71	Statice sinuata.
7	222	Struthiola erecta.
6	212	Tanacetum flabelliforme.
7	245	Teucrium latifolium.
9	308	Verbena Aubletia.
7	242	Veronica decussata.

INDEX

To the *Stove Plants* contained in the first *Ten Volumes* of the *Botanical Magazine*.

<i>Vol.</i>	<i>Pag.</i>	
10	338	Allamanda cathartica.
4	125	Alstrœmeria Ligtu.
9	305	Amaryllis equestris.
2	47	—— formosissima.
4	129	—— vittata.
10	339	Arum trilobatum.
1	34	Browallia elata.
2	41	Calceolaria pinnata.
3	107	Cassia Chamæcrista.
4	131	Catesbæa spinosa.
5	152	Epidendrum cochleatum.
4	144	Ferraria undulata.
4	141	Heliotropium peruvianum.
5	158	Hibiscus Rosa sinensis.
10	360	—— speciosus.
7	244	Ipomœa Quamoclit.
5	169	Ixora coccinea.
10	325	Justicia nasuta.
3	96	Lantana aculeata.
4	116	Limodorum tuberosum.
7	225	Lobelia surinamensis.
7	250	Myrtus tomentosa.
2	66	Passiflora alata.
8	288	—— ciliata.

7	230	Plumbago rosea.
8	279	Plumeria rubra.
8	286	Portlandia grandiflora.
1	26	Stapelia variegata.
4	119	Strelitzia Reginae.
8	281	Turnera angustifolia.
7	248	Vinca rosea.

INDEX.

In which the Plants contained in the first *Ten Volumes* of the BOTANICAL MAGAZINE are arranged according to the System of LINNÆUS, as published by himself.

MONANDRIA.

Monandria Monogynia.

LOPEZIA RACEMOSA.

Monandria Digynia.

Blitum virgatum.

DIANDRIA.

Diandria Monogynia.

Calceolaria pinnata.

—— Fothergillii.

Jasminum officinale.

—— odoratissimum.

Justicia nasuta.

Monarda fistulosa, var.

Salvia aurea.

Syringa vulgaris.

Verbena Aubletia^[6].

Veronica decussata.

TRIANDRIA.

Triandria Monogynia.

Antholyza Cunonia.

Crocus vernus.

Gladiolus communis.

—— cardinalis.

—— tristis.
Iris ochroleuca.
—— persica.
—— pumila.
—— pavonia.
—— sambucina.
—— sibirica.
—— spuria.
—— susiana.
—— variegata.
—— versicolor.
Ixia Bulbocodium.
—— chinensis.
—— crocata.
—— flexuosa.
—— longiflora.

Triandria Digynia.

Briza maxima.

TETRANDRIA.

Tetrandria Monogynia.

Buddlea globosa.
Catesbæa spinosa.
Ixora coccinea.
Protea mellifera.
Scabiosa atropurpurea.
Struthiola erecta.

PENTANDRIA.

Pentandria Monogynia.

Allamanda cathartica.
Anagallis Monelli.
Azalea nudiflora, var.

Campanula carpatica.
—— grandiflora.
—— Speculum.
Cerinthe major.
Chironia baccifera.
—— frutescens.
Convolvulus althæoides.
Convolvulus linearis.
—— Nil.
—— purpureus.
—— tricolor.
Cyclamen Coum.
—— persicum.
Cynoglossum Omphalodes.
Diosma uniflora.
Dodecatheon Meadia.
Goodenia lævigata.
Heliotropium peruvianum.
Hyoscyamus aureus.
Ipomœa coccinea.
—— Quamoclit.
Lysimachia bulbifera.
Phlox divaricata.
Phylica ericoides.
Plumbago rosea.
Plumeria rubra.
Portlandia grandiflora.
Primula acaulis, pl.
—— marginata.
—— villosa.
Pulmonaria virginica.
Solanum laciniatum.
Soldanella alpina.
Spigelia marilandica.
Strelitzia Reginæ.
Verbascum Myconi.
Vinca rosea.

Pentandria Digynia.

Apocynum androsæmifolium.

Gentiana acaulis.

Stapelia variegata.

Pentandria Trigynia.

Turnera angustifolia.

Viburnum Tinus.

Pentandria Pentagynia.

Linum arboreum.

—— flavum.

Mahernia incisa.

—— pinnata.

Statice sinuata.

HEXANDRIA.

Hexandria Monogynia.

Allium descendens.

Alstrœmeria Pelegrina.

—— Ligtu.

Amaryllis Atamasco.

Amaryllis equestris.

—— formosissima.

—— lutea.

—— sarniensis.

—— vittata.

Anthericum Liliastrum.

Bulbocodium vernal.

Cyrtanthus angustifolius.

Erythronium Dens Canis.

Fritillaria imperialis.

Hemerocallis flava.

—— fulva.

Hyacinthus botryoides.
—— comosus.
—— racemosus.
Lachenalia tricolor.
Leucojum vernum.
Lilium bulbiferum.
—— candidum.
—— Catesbæi.
—— chalcedonicum.
Narcissus angustifolius.
—— Bulbocodium.
—— biflorus.
—— Jonquilla.
—— incomparabilis.
—— major.
—— minor.
—— odorus.
—— triandrus.
Ornithogalum aureum.
—— nutans.
Scilla amæna.
—— campanulata.
Tradescantia virginica.

Hexandria Trigynia.

Trillium sessile.

HEPTANDRIA.

Heptandria Monogynia.

Disandra prostrata.

OCTANDRIA.

Octandria Monogynia.

Daphne Cneorum.

Epilobium angustissimum.

Erica ampullacea.

—— baccans.

—— cerinthoides.

—— grandiflora.

—— herbacea.

—— Massoni.

—— persoluta.

—— ventricosa.

Fuchsia coccinea.

Michauxia campanuloides.

Oenothera fruticosa.

—— pumila.

—— purpurea.

—— rosea.

Passerina grandiflora.

Tropæolum majus.

—— minus.

Octandria Trigynia.

Polygonum orientale.

Decandria Monogynia.

Cassia Chamæcrista.

Dais cotinifolia.

Fagonia cretica.

Kalmia angustifolia.

—— glauca.

—— hirsuta.

—— latifolia.

Sophora tetraptera.

Decandria Digynia.

Dianthus barbatus.

—— Caryophyllus, var.

—— chinensis.
—— superbus.
Saponaria Ocymoides.
Saxifraga crassifolia.
—— mutata.
—— sarmentosa.

Decandria Trigynia.

Silene pendula.

Decandria Pentagynia.

Agrostemma Coeli rosa.
—— coronaria.
Cotyledon orbiculata.
Lychnis chalcedonica.
—— coronata.
Oxalis caprina.
—— versicolor.
Sedum Anacampseros.
—— populifolium.

DODECANDRIA.

Dodecandria Trigynia.

Reseda odorata.

Dodecandria Dodecagynia.

Sempervivum arachnoideum.
—— monanthes.
—— tortuosum.

ICOSANDRIA.

Icosandria Monogynia.

Amygdalus nana.
Cactus flagelliformis.
Myrtus tomentosa.

Icosandria Pentagynia.

Mesembryanthemum aureum.
—— barbatum.
—— bicolorum.
—— dolabriforme.
—— pinnatifidum.
—— viridiflorum.
Pyrus spectabilis.

Icosandria Polygynia.

Fragaria monophylla.
Potentilla grandiflora.
Rubus arcticus.
—— odoratus.
Rosa muscosa.
—— semperflorens.

POLYANDRIA.

Polyandria Monogynia.

Argemone mexicana.
Capparis spinosa.
Cistus formosus.
—— incanus.
—— ladaniferus.
Papaver orientale.
Sanguinaria canadensis.

Polyandria Pentagynia.

Aquilegia canadensis.
Nigella damascena.

Polyandria Polygynia.

Adonis vernalis.
Anemone Hepatica.
—— hortensis.
—— sylvestris.
Clematis integrifolia.
Helleborus hyemalis.
—— lividus.
—— niger.
Liriodendron Tulipifera.
Ranunculus aconitifolius, pl.
—— acris, pl.
—— amplexicaulis.
—— gramineus.
Trollius asiaticus.

DIDYNAMIA.

Didynamia Gymnospermia.

Dracocephalum denticulatum.
Lamium Orvala.
Melissa grandiflora.
Origanum Dictamnus.
Prunella grandiflora.
Teucrium latifolium.

Didynamia Angiospermia.

Antirrhinum alpinum.
—— purpureum.
—— sparteum.
—— triphyllum.
—— triste.
Browallia elata.
Buchnera viscosa.
Celsia linearis.

Erinus alpinus.
Lantana aculeata.
Manulea tomentosa.
Melianthus minor.
Mimulus ringens.
—— aurantiacus.
Selago ovata.

TETRADYNAMIA.

Tetradynamia Siliculosa.

Alyssum deltoideum.
—— halimifolium.
—— saxatile.
—— utriculatum.
Draba aizoides.
Iberis gibraltarica.
—— umbellata.

Tetradynamia Siliquosa.

Arabis alpina.
Cheiranthus maritimus.
—— mutabilis.

MONADELPHIA.

Monadelphias Pentandria.

Erodium incarnatum.
Hermannia alnifolia.
—— althæifolia.
—— lavandulifolia.

Monadelphias Heptandria.

Pelargonium acetosum.
—— betulinum.

- bicolor.
- ceratophyllum.
- cordifolium.
- echinatum.
- glutinosum.
- tetragonum.
- tricolor.

Monadelphia Octandria.

Aitonia capensis.

Monadelphia Decandria.

Geranium anemonefolium.

- angulatum.
- lanceolatum.
- peltatum.
- Radula.
- Reichardi.
- striatum.

Monadelphia Polyandria.

Camellia japonica.

Hibiscus Rosa sinensis.

- speciosus.
- syriacus.
- Trionum.

Lavatera trimestris.

Sida cristata.

DIADELPHIA.

Diadelphia Hexandria.

Fumaria cava.

- glauca
- solida.

Diadelphia Octandria

Polygala bracteolata.
—— Chamæbuxus.
—— Heisteria.

Diadelphia Decandria.

Anthyllis tetraphylla.
Aspalathus pedunculata.
Borbonia crenata.
Colutea arborescens.
—— frutescens.
Coronilla glauca.
—— valentina.
—— varia.
Cytisus Laburnum.
—— sessilifolius.
Genista triquetra.
Glycine bimaculata.
—— coccinea.
—— rubicunda.
Hedysarum obscurum.
Indigofera candicans.
Lathyrus articulatus.
—— odoratus.
—— sativus.
—— tingitanus.
—— tuberosus.
Lotus hirsutus.
—— jacobæus.
—— tetragonolobus.
Lupinus luteus.
—— perennis.
Ononis fruticosa.
—— Natrix.
—— rotundifolia.

Robinia hispida.
Spartium junceum.
Trifolium incarnatum.

POLYADELPHIA.

Polyadelphia Dodecandria.

Monsonia speciosa.

Polyadelphia Polyandria.

Hypericum balearicum.
—— calycinum.
—— Coris.
—— monogynum.
Metrosideros citrina.

SYNGENESIA.

Syngenesia Polygamia Æqualis.

Catananche cærulea.
Crepis barbata.
Scorzonera tingitana.

Syngenesia Polygamia Superflua.

Aster alpinus.
—— tenellus.
Bellis perennis, fl. pl.
Chrysanthemum indicum.
Cineraria Amelloides.
—— lanata.
Gnaphalium eximium.
Senecio elegans.
Tagetes patula.
Tussilago alpina.
Tanacetum flabelliforme.

Zinnia multiflora.

Syngenesia Polygamia Frustranea.

Centaurea glastifolia.

—— montana.

Coreopsis verticillata.

Gorteria rigens.

Helianthus multiflorus.

Rudbeckia purpurea.

Syngenesia Polygamia Necessaria.

Othonna pectinata.

Syngenesia Monogamia.

Lobelia Cardinalis.

—— surinamensis.

Viola pedata.

GYNANDRIA.

Gynandria Diandria.

Cypripedium acaule.

—— album.

Epidendrum cochleatum.

Limodorum tuberosum.

Gynandria Triandria.

Ferraria undulata.

Sisyrinchium Iridioides.

Gynandria Pentandria.

Passiflora alata.

—— cærulea.

—— ciliata.

Gynandria Polyandria.

Arum trilobatum.

POLYGAMIA.

Polygamia Monoccia.

Mimosa myrtifolia.

—— verticillata.

FOOTNOTES:

[1] Ait. Kew.

[2] The flowers of *rotundifolia* are yellow, and therefore cannot be our plant.

[3] Ait. Kew.

[4] *Donn. Hort. Cantab.* 1796.

[5] Ait. Kew.

[6] Some Plants of this genus have two, others four stamina;—the present one has four.

Transcriber's Note:

1. Original spelling has been retained.
2. Where necessary, punctuation has been corrected.
3. The Table of Contents was created by the transcriber.

*** END OF THE PROJECT GUTENBERG EBOOK THE BOTANICAL
MAGAZINE, VOL. 10 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the

collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain

Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, “Information about donations to the Project Gutenberg Literary Archive Foundation.”
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration,

modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.