

THE NAUTILUS.

VOL. XXXI. OCTOBER, 1917. NO. 2

The cover image was created by the transcriber and is placed in the public domain.

<https://www.pgdp.net>

THE NAUTILUS.

VOL. XXXI. OCTOBER, 1917. NO. 2

The cover image was created by the transcriber and is placed in the public domain.

<https://www.pgdp.net>

The Project Gutenberg eBook of The Nautilus. Vol. XXXI, No. 2, October 1917

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The Nautilus. Vol. XXXI, No. 2, October 1917

Author: Various

Editor: Charles Willison Johnson
Henry Augustus Pilsbry

Release date: April 11, 2019 [eBook #59245]

Language: English

Other information and formats: www.gutenberg.org/ebooks/59245

Credits: Produced by The Online Distributed Proofreading Team at <http://www.pgdp.net> (This file was produced from images generously made available by The Internet Archive)

*** START OF THE PROJECT GUTENBERG EBOOK THE
NAUTILUS. VOL. XXXI, NO. 2, OCTOBER 1917 ***

\$2.00 per Year. (\$2.20 to Foreign Countries.) 50 cts. a copy

**THE
NAUTILUS
A QUARTERLY
DEVOTED TO THE INTERESTS
OF CONCHOLOGISTS**

EDITORS AND PUBLISHERS:

H. A. PILSBRY, Special Curator of the Department of Mollusca, Academy of Natural Sciences,
Philadelphia.

C. W. JOHNSON, Curator of the Boston Society of Natural History.

Vol. XXXI.

OCTOBER, 1917.

No. 2

CONTENTS:

Notes on the variation of <i>Ischnochiton conspicuus</i> . By <i>E. P. Chace</i>	<u>37</u>
<i>Lampsilis ventricosa cohongoronta</i> in the Potomac River. By <i>Wm. A. Marshall</i>	<u>40</u>
Collecting about Guantanamo Bay, Cuba. By <i>John B. Henderson</i>	<u>41</u>
Amnicolidæ from Oneida Lake, N. Y. By <i>H. A. Pilsbry</i>	<u>44</u>
New Genera and Species of Central American Naiades. By <i>L. S. Frierson</i>	<u>47</u>
On the rate of growth of pond Unios. By <i>L. S. Frierson</i>	<u>49</u>
A new South African Nesopupa. By <i>H. A. Pilsbry</i>	<u>50</u>
A new Gundlachia from Guatemala. By <i>Bryant Walker</i>	<u>51</u>
A list of shells from the East Coast of Florida. By <i>Bryant Walker</i>	<u>53</u>
Collecting in Digby, Nova Scotia. By <i>Lilian Dyer Thompson</i>	<u>57</u>
A new type of the naiad-genus <i>Fusconaia</i> . By <i>A. E. Ortmann</i>	<u>58</u>
The Relation of snail fauna to floods. By <i>A. Richards</i>	<u>64</u>
William Bullock Clark	<u>68</u>
Publications received	<u>69</u>
Notes	<u>71</u>

C. W. JOHNSON, Business Manager,

Boston Society of Natural History,
Berkeley Street,
Boston, Mass.

Entered as Second-Class matter at the Boston Post-Office.

The Cruise of the “Tomas Barrera”

By JOHN B. HENDERSON

32 full-page illustrations in black and white. 5 color plates. 5 maps. Octavo.
\$2.50 net.

This is the narrative of a scientific expedition to Western Cuba and the Colorados Reefs, with observations on the Geology, Fauna, and Flora of the region, undertaken in May and June, 1914, under the joint auspices of the Smithsonian Institution and the Cuban Government.

“A genuine feast for the reader. It is a narrative replete with interest concerning the roads, mountains, trees, dwellings, the inland lakes and ponds (alive with queer fish), many species of mollusca, found along the coast; the land crabs and their curious habits,” etc.—*Louisville Courier Journal*.

ALL BOOKSELLERS

NEW YORK
2 West 45th St.

G. P. PUTNAM'S SONS

LONDON
24 Bedford St.

NEW SHELL LISTS FREE.

A new list of Philippine Land Shells covering many of the most beautiful forms at greatly reduced prices. A list of 2,000 species of shells, priced at 5 to 10 cents per species. Two lists of the finer cabinet shells, the more aristocratic forms. List of American Land Shells. Illustrated list of Philippine Shells. Further lists in preparation. I desire correspondence with collectors who wish to build up large and extensive cabinets. I have in stock over three times as many species as are covered by my lists. Collections of a strictly scientific nature purchased for cash.

WALTER F. WEBB,

202 Westminster Road,

ROCHESTER, N. Y.

Exchange notices not exceeding three lines will be free to subscribers as long as space will allow.

FOR EXCHANGE: Marine shells from various parts of the world, for others. Send lists.

J. R. LEB. TOMLIN, 120 Hamilton Road, Reading, England.

THE NAUTILUS.

VOL. XXXI.	OCTOBER, 1917.	No. 2
------------	----------------	-------

NOTES ON THE VARIATION OF ISCHNOCHITON CONSPICUUS CPR.

BY E. P. CHACE.

To the naturalist the study of variation and environment and their relation to each other is always interesting and to the conchologist who studies his shells in their natural surroundings as well as in the cabinet many things are revealed. It is not, often, however, that variation in form may be so easily traced to qualities of environment as in the following instance.

Ischnochiton conspicuus, Cpr. is common at San Pedro and is usually found on the under side of rocks in sandy tidepools. In this situation they grow quite regularly, showing but slight variation in form. They are active fellows and evidently sensitive to light; for if the rocks to which they are clinging are turned over, they soon glide away, always taking the shortest route to the under side of the rock.

While cleaning a lot of this species taken at San Pedro last fall two specimens were noticed which differed so widely from the others that they might easily have been mistaken for another species. They were much wider and lower-arched than the typical form and the posterior corners of the valves were rounded off, making the lateral areas very narrow.

Hoping to find more specimens of this odd form, a trip to Point Firmen was devoted wholly to the collection of chitons, with interesting results. The usual species were found in the tidepools including numerous specimens of *Ischnochiton conspicuus* of the ordinary form. Ledges of soft rock beyond the tidepools contained many old pholad holes and in these were found the form for which I was searching. Twenty specimens of various sizes were collected from as many pholad holes. In the larger specimens the foot had become so greatly enlarged to fit the concave

bottoms of the holes that it was impossible for them to curl up in the usual manner. Some of these specimens were so badly eroded by the sand and gravel which wash in and out of the holes that the anterior valve was reduced to two thirds of its normal height.

In color pattern, sculpture, and mantle characters these specimens were identical with those from the tidepools, and, as will be seen by referring to the table of measurements, the smaller specimens approach quite closely to the proportions of the typical or tidepool forms.

TYPICAL OR TIDEPOOL FORMS. (Lot 1)				SPECIMENS FROM PHOLAD HOLES. (Lot 2)			
<i>Lon.</i>	<i>Lat.</i>	<i>Index.</i>	<i>Div.</i>	<i>Lon.</i>	<i>Lat.</i>	<i>Index.</i>	<i>Div.</i>
99	35	2.52	130°	65	33	1.97	135°
83	33	2.51	130°	63	32	1.97	155°
83	30	2.76	125°	62	27	2.29	140°
62	22	2.80	125°	60	30	2.00	130°
59	23	2.56	130°	56	29	1.93	130°
38	14	2.71	130°	42	17	2.47	130°

In brief, Lot 1 shows an angle of divergence constant at 125° to 130°, where Lot 2 shows an angle varying from 130° to 155°, and a proportion of length to width 2.51 to 2.80 as against a proportion ranging from 2.47 to 1.93.

The noticeable differences to the eye are first, the narrow and sharply raised lateral areas, and second, the shape of the posterior edge of the median valves. In the tidepool specimens the posterior or exposed edge of each valve is a straight line, while in specimens from the pholad holes this line becomes a double convex curve, the most posterior portion of the valves being about midway between the beaks and the girdle.

These differences seem to be explained by the following facts. In collecting, the tidepool specimens are usually found on the under side of large rocks and well back from the edge. This situs protects them from the light which they evidently find objectionable, but it makes necessary a nightly journey of about two feet to the nearest growth of algae on which they feed. This activity stretches the girdle downward from the edges of the valves and permits a free play of all the valves so that the mantle deposits its shelly secretions according to the normal habit of the species. The specimens living in the pholad holes, however, apparently never leave them as they are frequently found feeding on the fucus which overhangs them. It

protects them from the light, so they have no occasion to move about, and the sand which is washed down into these burrows would make re-entrance almost impossible. A series of these specimens shows a gradual change of form. The young specimens are very similar to young specimens from the tidepools, but as they increase in size they become crowded so that the valves press against each other, especially at the posterior end where the valves are bent back across the bottom of the hole. This crowding of the valves upon each other and the crowding of the girdle against the outer edges of the valves so displaces portions of the mantle as to cause the changes noted above.

Several specimens from each situs were disjointed and a study of the individual valves showed that those from pholad-hole specimens were thicker and had shorter sutural plates and a wider sinus, this last being especially noticeable in the valves from the posterior end. Apparently this change in the sinus is the result of the broadening of the connecting ligaments due to compression by the crowding valves.

A count of the insertion plates of these disjointed specimens was made and considerable variation noticed. So much, in fact, that more specimens were pulled apart for the express purpose of counting these plates. Representative counts were as follows: 9 slits on the anterior valve, 2-3 on the median valves, and 10 on the posterior. Others show 12, 2-3, 8; 11, 2-3; 14, 3-4, 11. Absolutely no difference in this character could be found between specimens from the tidepools and those from the pholad holes.

On page 64 of vol. xiv of the Manual of Conchology, Dr. Pilsbry says, "Carpenter has given a varietal name to a broad, worn specimen which he thus describes:

"Var. *solidus*. Very solid, wide, ashen; inside whitish, the posterior valve with 10, central valve 2-3, anterior valve 12 slits. Length 72, breadth 40, divergence 130°. Carpenteria, near Sta. Barbara, Cal. This is scarcely more than an individual variation. The mantle (girdle) is normal. The sculpture is worn away except at the edge. It has evidently lived in a very exposed situation."

From this description and the figure which he gives of the valves it would seem that this is the same form as my specimens from the pholad holes. I fully agree with Dr. Pilsbry that it is hardly worthy of varietal rank. It is, however, too distinct in appearance to be labeled simply *Ischnochiton*

conspicuus, Cpr., and I have therefore marked these specimens from the pholad holes *I. conspicuus*, Cpr., *form solida*, Cpr., using the term *form* as advocated by Dr. Cockerell, “to designate variations plainly due to environment.”^[1]

LAMPSILIS VENTRICOSA COHONGORONTA IN THE POTOMAC RIVER.

BY WILLIAM A. MARSHALL.

In 1912 Dr. A. E. Ortmann recorded^[2] finding in the Potomac River a variety of *Lampsilis ventricosa*^[3] to which he gave the name *cohongoronta*. His records were:

September 4, 1909. Potomac River, Hancock, Washington, Co., Md. (about two dozen).

May 9, 1911. South Branch, Potomac River, Southbranch, Hampshire Co., W. Va. (about a dozen).

August 16, 1911. Shenandoah River, Harper's Ferry, Jefferson Co., W. Va. (a single male, below medium size).

May 6, 1912. South Branch, Potomac River, Romney, Hampshire Co., W. Va. (about a dozen).

Dr. Ortmann remarked "It is probable that this species will turn up elsewhere in the Potomac. The localities known at present are all to the west of the Blue Ridge Mountain, that is to say, within the Great Alleghany Valley and the Alleghany Mountains."

Perhaps the above prediction has been realized in a specimen found at Great Falls, Md., by Mr. Manly D. Barber of Knoxville, Tennessee, in Sept. 1915. At that time Mr. Barber brought to the National Museum a basketful of naiades which he had collected the same day at Great Falls, about 18 miles above Washington. Among the shells, which were mostly dead ones, was a specimen of *cohongoronta*, dead, but in a fine state of preservation and with the periostracum nearly unblemished except for the usual erosion at the beaks. Its appearance indicated that it had been recently alive and that

its home had been in the immediate vicinity of the place in which it was found. Had it been washed down from Harper's Ferry, some 50 or more miles above Great Falls it probably would have shown ill effects from so long a journey.

When found the two valves were separated, but so accurately do they fit together that it is evident they belong to the same individual. The fact that the valves were separated and yet were found near each other is additional (though not conclusive) evidence that they had not been transported any great distance by currents. At any rate this is the first recorded finding of the species in the Potomac River so far south as Great Falls.

The specimen is rather a small one. It measures, length 71 mm.; height 47 mm.; diameter 28 mm. It is in the collection of the U. S. National Museum, catalogue number 273834.

COLLECTING DAYS ABOUT THE NAVAL STATION, GUANTANAMO BAY, CUBA.

BY JOHN B. HENDERSON.

In March last, while waiting for a boat to take us to Haiti, Dr. Bartsch and I spent nearly three weeks at the U. S. Naval Station at the entrance to Guantanamo Bay. We employed our time in exploring the country about and subjecting it to a high degree of intensive collecting. In this eastern corner of Cuba the coastal strip of some ten miles in width is a semi-arid region with a complex of mountains that are either quite bare of trees or, at most, covered with a scrub forest and low-growing spiny shrubs, with, here and there, a wealth of cacti that almost suggests Lower California. The rock foundation of all this region,—barring some shore strips of very recently elevated coral, is everywhere composed of about everything in the line of rocks except limestone. This is a condition that in the Antilles usually spells disappointment and failure to the snail hunter. North of the big bay and then across several miles of low flat country, just where the foothills of the sierras begin, lies the city of Guantanamo, interesting to us as the home of Charles Ramsden, the naturalist. Just north of Guantanamo is a great rampart of high limestone mountains which beckon most alluringly to the collector. Sections of this rampart, somewhat arbitrarily marked off, are the “Monte Verde,” the “Monte Toro” and the “Monte Libano” of classic fame in Cuban Natural History.

In company with Ramsden we spent a wonderful day on nearby Monte Libano but a revolution that was then devastating the province and filling the land with incendiaries and bandits drove us out of this richer field and obliged us to confine our attentions thereafter to the arid country lying within the safer limits of the Naval Station,—some fifty square miles upon which Uncle Sam holds a long lease.

It seems to be a natural law that arid or desert lands support but few species of snails, but that these few species exist in great numbers and that they take on a very considerable range of variation. All this is perfectly true of this region. We were constantly amazed by the great number of specimens to be found; and each day of exploration in some new valley or over some range of hills added even greater figures of abundance to our already astonishing records.

The “prevailing” snail of this region is *Cepolis ovumreguli* Lea. Its shell is very suggestive of the true helix of Spain or Algeria of the *lactea* group. The variation is exceedingly great in color, size and shape, and it would make a dozen excellent species if the intermediates were left out of account. Those living near the coast and among the cacti of the most arid parts of the district are of whiter and more dull color, are more banded and show a decided tendency to abnormalities, especially about the apertures. Specimens from further inland are more polished and shining, even as though varnished, and are much more given to a dotted or fly-specked type of ornamentation than to bands. A fence-post or a dead tree-limb with a hundred specimens closely assembled in aestivation was no unusual sight. We learned finally to pay no attention to them. Upon the low bushes in certain localities the lovely little *Cepolis lucipeta* Poey cling like berries. These are the largest and finest of the species I have ever seen. The range of color variation in this delightful little snail is also very great, but the colors never blaze out in the vivid flash of the *Polymitas*. The blues and purples and chestnut browns are subdued but very rich and splendid. One very noticeable color form is the subsp. *velasqueziana* of Poey where the many broken bands of the type coalesce into two broad bluish-black zones of solid color.

As nearly all the vegetation of this dry region bears thorns we did not at first discover that many of these thorns were in reality *Macrocerami*. When we did find this out we could see nothing else. Bartsch and I finally agreed, and shook hands upon it, that we would gather no more of them, and a stiff penalty was placed upon any violation of the compact. Two hundred and more from one bush is an earlier record before we really got started. This is the *Macroceramus festus* (Gundl.) Pfr., blue and yellow and buff in color. Another arboreal snail of this section is *Polymita versicolor* Born and it is probably very abundant in places although we never saw more than fifteen or twenty on any one tree. This is to me the least attractive species of that

wonderful genus of richly painted snails. The brilliant yellow and pink are too primitive and the two colors do not seem to harmonize very well. It always impresses me as an experimental species that was laid aside in nature's laboratory as not wholly a success.

There were some ground snails too, but to secure living ones required much grubbing up of tufts of tall grass and shaking out their roots, like digging up miniature potatoes. These are the *Annularia putris* (Gundl.) Pfr. and the *Chondropoma marginalbum* (Gundl.) Pfr., the latter apparently quite rare. There are no minute things beyond some few *Thysanophora inaguensis* Weinland.

Some days we spent gathering marines on the little pebbly beaches hidden far down under the lofty cliffs that mark this rugged shore line, and we obtained some unusual species washed up from the exceedingly narrow island-shelf; blue water is but a few hundred yards out. Among these are some *Conus cedonulli* Lam. Beach collecting is, however, an aggravation; and too much of it becomes a misdemeanor in the collector's ethical code, for it obliges an acceptance of something short of the best.

AMNICOLIDÆ FROM ONEIDA LAKE, N. Y.

BY HENRY A. PILSBRY.

The New York College of Forestry, under the direction of Professor Hugh P. Baker, is carrying on a biological survey of Oneida Lake and has issued an interesting bulletin^[4] upon the relations of mollusks to fish, by Frank C. Baker. Some *Amnicolidæ* obtained during this work, and subsequent to the preparation of the bulletin were submitted to the writer. The collection proves to be of considerable interest, including some species not before noticed.

AMNICOLA BAKERIANA, n. sp.

The shell is umbilicate, turritid-conic, thin, whitish-corneous, somewhat translucent, with unevenly developed striation, distinct, and close in places, weaker and sparse elsewhere. The summit is decidedly obtuse, as in *A. limosa*, the first whorl being nearly planorboid; subsequent whorls are evenly, strongly convex. The aperture is very shortly ovate, almost round, its length contained almost $2\frac{1}{2}$ times in that of the shell. Peristome thin, in contrast with the preceding whorl for a short distance.

Length 4.3, diam. 2.7 mm.; 5 whorls (type).

Length 3.75, diam. 2.3, length of aperture 1.35 mm.; $4\frac{2}{3}$ whorls.

Length 4.1, diam. 2.75, length of aperture 1.65 mm.; $4\frac{2}{3}$ whorls.

Oneida Lake; off Short Point in $8\frac{1}{2}$ ft., mud bottom. Lower South Bay, in 18 ft., on mud bottom.

This species resembles *A. limosa* in the conspicuously obtuse apex, but differs by the more elevated, turritid spire and the smaller calibre of the whorls, hence smaller aperture. It is also a weaker shell, with more whorls in specimens of the same length.

There is also an abundant smaller form, resembling the typical form in texture, apex and shape of the whorls, varying in form, but relatively broader than the type. There are some intermediate examples, but as Mr. Baker considers it desirable to have a designation for this form, it may be called *A. bakeriana* form *nimia*. The type measures: length 3, diam. 2.5, length of aperture 1.4 mm.; 4 whorls.

AMNICOLA CLARKEI, n. sp.

The shell is narrowly umbilicate, conic, a little obtuse at the apex, corneous, nearly smooth. The whorls are very convex, separated by a deep suture, the last whorl tubular. The aperture is distinctly oblique, almost circular, the upper end rounded, but a trifle more narrowly so than the base. It projects but little beyond the preceding whorl laterally. The peristome is thin, continuous, scarcely or barely in contact with the preceding whorl above.

Length 3.1, diam. 1.9, length aperture 1.1 mm.; 5 whorls (type).

Length 2.8, diam. 1.6, length aperture 0.85 mm.

Operculum having the spiral rather large, the nucleus being above the lower third.

This little species resembles *Lyogyrus* by its tubular whorls of small calibre. The whorls are more convex and increase less rapidly than in *Amnicola walkeriana*, which is also less slender. *A. schrockingeri* Ffld. has less deeply convex whorls, and the apex is more acute. *A. bakeriana* is much larger, with a more obtuse apex.

Found in Short Point Bay, Oneida Lake, near shore, in 3 feet of water, bottom of sand with algae; also in Lower South Bay, etc. Collected by Mr. F. C. Baker.

It is named for Dr. John M. Clarke, the distinguished Director of the Museum of the State of New York.

AMNICOLA ONEIDA, n. sp.

The shell is typically more slender than *A. lustrica*, turritid-conic, narrowly umbilicate, corneous, minutely striate. The apex is slightly obtuse, but the first whorl projects visibly, as in *lustrica*, whorls *very convex*, parted by a *deep suture*. The aperture is ovate, small, its length contained more than 3 times in that of the shell; upper extremity narrowly rounded. The peristome is continuous, thin, very briefly in contact with the preceding whorl above.

Length 4, diam. 2, length of aperture 1.25 mm.; 6 whorls.

Lower South Bay, Oneida Lake, N. Y., collected by F. C. Baker, 1916.

This species is typically narrower than *A. lustrica* Pils., with a smaller aperture and shorter whorls; but it is chiefly distinguished by the more convex whorls (deeper suture), and the rounded instead of angular posterior end of the aperture. In *Paludestrina nickliniana* the last whorl is much longer. Possibly it may be a subspecies of *lustrica*, yet it has so distinct an appearance that a special name seems desirable. There are also wider examples, which still differ from *lustrica* by the deeper suture and aperture.

NEW GENERA AND SPECIES OF CENTRAL AMERICAN NAIADES.

BY L. S. FRIERSON.

In 1893 Messrs Crosse and Fischer divided the Mexican Naiades into quite a number of sections, to which they assigned names. Almost simultaneously (in 1900) von Martens and C. T. Simpson, in treating the Central American Naiades, accepted some of these sections of Crosse and Fischer, raising them to generic or subgeneric rank. Because of paucity of material, considerable diversity of opinion concerning the specific identity of several species may be noted in the works of these authors. Furthermore, their work of classification being done independently and from different points of view, the same species was sometimes placed by them in different genera.

Thanks to the arduous labors of A. A. Hinkley, who has again and again enriched our cabinets with material and data from these tropical countries, we are enabled to offer the following suggestions concerning some of the genera of these shells, and also the description of an unpublished species.

NEPHRONAIAS. This genus has for its type the *Unio plicatulus*, Küster, a species identified by von Martens as belonging to the Lampsiline shells, as *aztecorum*. Mr. Simpson however believed it to be nearly allied to the *persulcatus*, a markedly Unioidean shell. In this the writer follows Mr. Simpson.

The genus *Nephronaias* as constituted by Mr. Simpson embraces two quite distinct groups, divisible as follows.

Nephronaias (s. s.) embraces *plicatulus*, *persulcatus*, *melleus*, *dysoni*, *ortmanni*, *ravistellus*, etc. Ample material of these two latter species show that they are anatomically very closely allied to *Elliptio*. There is no sexual

difference of shape, and the gill is gravid in its whole length. *Nephronaias* differs from *Elliptio* in its sulcated disc, in its beak sculpturing, etc.

Included in *Nephronaias* by Simpson are, however, shells of a totally different type, such as *medellinus*, *gundlachi*, *sapotalensis*, etc. These latter are sexually dimorphic, smoother, more generally rayed, and the gravid uterus is of Lampsiline type.

The position of the dorsal scars within the beak cavities is different, in the examples of the pseudo *Nephronaias* seen by the writer. *Nephronaias* (s. s.) possesses an (accessory?) adductor scar attached to the frontal portion of the cardinal teeth, which is either absent or obsoletely marked in the second assemblage. For this latter group the writer, therefore, proposes to use the generic term of *Actinonaias* Crosse and Fischer, 1893, type *U. sapotalensis* Lea. The female of this species has been described by Dr. Ortmann (1912). *Actinonaias* embraces, besides the type, *medellinus*, *gundlachi*, (accepting Simpson's interpretation of this latter species), and others.

PSORONAIAS, Crosse and Fischer (1893). This group of remarkable shells, embracing *crocodilarum*, *psoricus*, *semigranosus*, etc., was provisionally treated by Simpson as a group of *Elliptio*, but their remarkable sculpturing, and the deep beak cavities of some of their species, led him to observe that it was possible that the group should, after all, be placed in *Quadrula*.

I follow von Martens, in giving generic rank as above to the group. The type is *Unio psoricus*. To this genus we are enabled to add a species hitherto undescribed, under the name of

PSORONAIAS KUXENSIS, n. sp., Pl. VII, figs. 1, 2.

Shell small, compressed, rough, brown, biangular. Length 50, height 30, diam. 17 mm.

Shell hyperbolically rounded before, the extreme frontal point below the centre. Dorsum slightly arched, descending behind the ligament to the widely biangular posterior; the upper angle of which is midway the height, the lower angle very little above the base, which is nearly straight. The beaks are small, low, acute, approximate; and *apparently*, concentrically ridged. Epidermis dark brown (olivaceous and obsoletely rayed in the young), rough, the lines of growth numerous and well impressed. The discs are covered with fine pustulations, more pustular in front, biradially linear behind. The post ridge is low, but distinctly double, making the shell biangulate behind. The teeth are double in the left valve, single in the right. The cardinals are deeply sulcate and stout. Laterals slightly curved or nearly straight, separated by an interdentum. Nacre purple, beak cavities rather deep. Dorsal scars numerous, extending in a row from above the centre of the cavity down and forward upon the base of the cardinal teeth. Three well impressed muscle scars in front, two behind, the latter almost confluent. Habitat, Kux Creek, Chama, Guatemala. Collected by Mr. A. A. Hinkley, Feb. 6, 1917. A few dead specimens were obtained on the bank of the Isaibha River (Chama) of which the Kux Creek is a tributary. Type in Academy Natural Sciences. Cotypes in collection of A. A. Hinkley, the author and U. S. N. Museum.

I place this species in *Psoronaias* Crosse and Fischer, type *U. psoricus*, because of its evident relationship to *crocodilarum*, and *distinctus*, differing mainly from the latter in size and degree of inflation, being much inferior in both respects to *distinctus*.

ON THE RATE OF GROWTH OF POND UNIOS.

BY L. S. FRIERSON.

During the latter part of March 1916, the writer, for the purpose of constructing a fish pond, excavated a barrow-pit near the bank of a small creek, about ten feet wide, and at the time nearly dry. The barrow-pit was perhaps one hundred feet long, fifty feet wide and three feet deep. Early in April, 1916, the pit became full of water, overflowing from the adjacent creek, and together with two subsequent overflows, supplemented with seepage from the newly constructed fish pond, the pit remained more or less full of water, until May 25, 1917, when it was drained by a ditch into the nearby creek. From the dried bottom of this pit some thirty Unios were picked up by the writer. Ten of these were *Unio tetralasmus* Say, and the rest were *T. texasensis* Lea. All the specimens were of remarkably uniform size and appearance. The *texasensis* being about one and a half inches, and the *tetralasmus* two and a half inches long. Exact dimensions of a *texasensis*: length 43, height 24, diam. 16 mm.; of *tetralasmus* 75, 40, and 25 mm.

Both of these species had attained puberty. A female *texasensis* had its gills fairly full of young glochidia. A *tetralasmus* had several (three or four) ovisacs with a few (remaining) glochidia. In assigning an age to these shells it is quite sure that the *tetralasmus* discharges its glochidia in March and early April, so that when picked up on May 25, these shells were just about fourteen months old, from the date of discharge from their mother's gills.

In the case of the *texasensis* (which spawns somewhat later) it is possible that these were dropped by fish of which, at least six species obtained access to the pit on May 7, 1916 (on which date an overflow occurred), thus making about thirteen months. At any rate the maximum age of either

species is fourteen months from their mother's ovisacs. One of the *U. tetralasmus* is shown of natural size in Pl. VII, fig. 4.

Another observation concerning pond mussels might here prove of interest. A large pond was cut into two by a railroad embankment, a culvert preserving the level and providing communication between the two. In the lower and larger pond a half-bushel of Yonkapin (*Nelumbium luteum*) seed was sown. It was six years before these seeds germinated. These plants, during the summer, cover the entire surface of the pond with their broad peltate leaves. In this pond the writer planted a colony of a dozen *Anodonta grandis*. Several years after, taking advantage of extreme low water, the writer made a careful survey of these twin ponds, with the result that hundreds of Anodons could be found in the upper pond, but not a single one was found in the lower pond. Either the shade killed the young shells, or else the glochidia-laden fish avoided the shade of the lotus plants and congregated in the upper pond (there are no *Nelumbii* in the upper pond). Is not this avoidance of shade a reason for the paucity of unios in the tropics?

A NEW SOUTH AFRICAN NESOPUPA.

BY H. A. PILSBRY.

NESOPUPA FARQUHARI, n. sp.

Among Pupillidae sent by Mr. J. Farquhar there is a new species from Grahamstown which may be defined by comparison with *Nesopupa griqualandica* (Melv. and Pons.).^[5] The new form is ovate, of about the size of the other species, which it resembles in sculpture and in the lamellae of the parietal wall and columella. The two palatal plicae are subequal, the upper emerging to the lip, the lower one also long, reaching to the inner edge of the peristome. There is a very small nodule on the base of the columella. In *griqualandica* the lower palatal plica is short and very deeply immersed and there is a distinct though small basal plica within the base, in front of the lower palatal plica. In *griqualandica* there is a deep sulcus outside, over the upper palatal plica, and a flattening or short groove over the lower palatal; but in *farquhari* the sulcus is far less impressed except quite close to the lip. The color is reddish brown. Length 1.65, diam. 0.9 mm.

Mr. Burnup's figure 9, in Melvill and Ponsonby's Revision,^[6] may perhaps represent this species, while their description in the same paper appears to comprise both *griqualandica* and *farquhari*, though chiefly relating to the former. Their pl. I, figs. 8 and 10 represent *griqualandica*. The new form is named in honor of one of the most successful South-African collectors. It will be figured in the Manual of Conchology.

A NEW GUNDLACHIA FROM GUATEMALA.

BY BRYANT WALKER.

GUNDLACHIA HINKLEYI, n. sp., Pl. I, figs. 10–16; Pl. III, fig. 1.

Shell subovate, being much wider posteriorly, the anterior margin rather shortly rounded, the right margin nearly rectilinear, but somewhat diverging anteriorly, the left margin obliquely expanded and broadly rounded, anterior margin wider and much more curved than the posterior; apex very excentric, depressed and decidedly turned toward the right side, bluntly rounded, smooth except for a few concentric wrinkles; color a very pale corneous, nearly pure white; lines of growth rather strong and irregular; anterior slope with strong radial striæ originating below the septate growth and extending to the anterior margin, similar striæ appear on the left lateral slope, but are scarcely, if at all, visible on the right slope; the septate portion of the shell is small in comparison with the adult expansion, it is narrow and the posterior portion covered by the septum is free from and projects over, but scarcely beyond, the posterior margin of the adult aperture; the first growth of the shell from the septate form is continued on the sides in a nearly direct continuation of the lateral slopes of the septate shell for some little distance, the anterior slope of this stage is also a continuation of the anterior slope of the septate stage but owing to the oblique position assumed by the septate shell is at first somewhat convex, as viewed laterally, later as the side slopes begin to expand, the anterior slope is continued in a nearly straight line to the margin; the left lateral slope of the adult shell below the secondary constriction is concave at first, becoming nearly straight toward the margin; the right lateral slope is less concave above and straighter and more oblique than the left; owing to the small size of the septum and consequent large aperture of the septate shell and the narrow first growth of the adult shell there is no distinct aperture to the septate portion visible in the adult shell from below, the whole interior of the adult shell appears to pass, practically unconstricted, directly into the septate portion; the posterior margin of the adult shell narrow and somewhat abruptly expanded and reflected.

Length 5.5; width 3.75; alt. 1.75 mm.

The septate shell is oblong, the sides being nearly parallel, but slightly expanding anteriorly, the right slightly convex and the left slightly concave; the posterior margin is regularly rounded; the anterior more broadly

rounded; the apex depressed, bluntly rounded, excentric, reaching nearly to the right margin, smooth except for slight concentric wrinkles, lines of growth fine and regular; the anterior slope is slightly convex; the very short posterior slope below the projecting apex to the line of the septum is straight and oblique; the right lateral slope is steep and nearly straight, the left slope very convex; the septum is very short, being less than half of the length of the septate shell, convex on its lower surface, the margin is very short, being less than half of the length of the septate shell, convex on its lower surface, the margin is very concave and on the right side, extends further forward than it does on left, there does not seem to be the distinct thickening of the margin so noticeable in other species; aperture much larger than in any other species yet described.

Length 2; width 1.5; alt. .75 mm.

Type (43455 Coll. Walker) from the Maya Farm, Quirigua, Guatemala, collected by A. A. Hinkley. Cotypes in the collection of Mr. Hinkley.

This fine species is the first from either Central or South America, of which both the septate and adult forms are known.

It differs from all other described species except *crepidulina* Guppy in the small size of the septum and the consequent difference in the position of the aperture of the septate stage in the adult shell. The septum in the specimen figured appears very like the incomplete septum in the North American species, but as the three adult specimens before me are exactly alike in the position of the septate shell, this would seem to be the normal condition in this species. The specimen figured, which is 3.25 mm. in length, has apparently slightly passed the septate stage and begun the growth of the constricted portion of the adult shell and shows the beginnings of the radial striæ.

With the *Gundlachias* was associated a species of *Lævapex*, very like the *excentricus* Morelet. Whether it has any closer relations with the *Gundlachia* remains to be determined as the radula has not yet been examined. While the general aspect of the two species, if such they be, is very similar, the *Lævapex* has a very much more acute apex than the *Gundlachia*.

As shown by the figure, the radula of this species is quite typical of the genus.

A LIST OF SHELLS FROM THE EAST COAST OF FLORIDA.

BY BRYANT WALKER.

The late Dr. Charles A. Davis, the well known peat-expert of the U. S. Bureau of Mines, in addition to his special acquirements in geology and botany, was a good all-round zoologist and had a lively and unaffected interest in the work that any of his friends might be carrying on in that department. It was his kindly habit in his travels about the country to preserve any specimens that he came across that seemed to him likely to be of interest to any of his zoological friends. It will be remembered that the conchologists owe to him the rediscovery of the long lost *Planorbis multivolvus* Case, (NAUT., XXI, p. 16), and also the little *Lymnæa davisii* Walker, (NAUT. XXII, p. 17), which bears his name.

In the spring of 1911 Dr. Davis' professional duties took him to Florida and while there he collected quite a number of samples of "drift," which in due time came into my possession. Several of the localities represented in the collection, such as Miami and St. Augustine, have already been reported upon by previous collectors and there seems to be no occasion to duplicate their work, but quite a number of the places visited by Dr. Davis have not been covered by any of the previous collectors in Florida and a record of the species found by him seems worthy of publication as a contribution to the distribution of the Mollusca along the east coast of the state.

I am indebted to Dr. George H. Clapp for the identification of the *Gastrocoptas* and *Vertigos*.

The list of localities and species represented in the collection is as follows:

MARSHES NEAR CHESTER SHOALS.

Euglandina rosea Fér.
Polygyra cereolus carpenteriana Bld.
Zonitoides minuscula (Binn.).
Zonitoides minuscula alachuana (Dall).
Vitrea dalliana ('Simpson' Pils.).
Pupoides modicus (Pfr.).
Gastrocopta rupicola (Say).
Gastrocopta pellucida hordeacella (Pils.).
Gastrocopta tappaniana (C. B. Ads.)?
Gastrocopta pentodon (Say).
Vertigo milium (Gld.).
Melampus coffeus (L.).
Detracia bulloides (Mont.).
Chondropoma dentatum (Say).
Plecotrema cubense (Pfr.).
Blauneria heteroclita (Mont.).
Microtralia minuscula (Dall).
Truncatella clathrus Lowe.
Truncatella caribæensis pulchella Pfr.
Truncatella bilabiata Pfr.
Littoridina monroensis (Ffld.).

Paludestrina? sp.? A single specimen that I can not approximate to any of the described species.

This is the first record, I believe, for *Plecotrema cubense* from the mainland of Florida. Originally described from Cuba, it was listed from the Bermudas by Dr. Pilsbry in 1900, (Trans. Conn. Acad., X, p. 504, pl. lxii, fig. 11), and there figured by him for the first time. Both he and Mr. John B. Henderson inform me that they have collected it on several of the Keys and I am indebted to both of them for the opportunity of comparing my specimen with theirs.

CHESTER SHOALS REFUGE STATION.

Euglandina rosea (Fer.).
Polygyra auriculata (Pfr.).
Polygyra uvulifera (Shutt.).
Polygyra cereolus (Mühlf.).
Polygyra cereolus carpenteriana (Bld.).
Polygyra cereolus septemvolva Say.
Polygyra cereolus volvoxis (Pfr.).
Praticolella jejuna (Say.).
Melampus coffeus (L.).
Detracia bulloides (Mont.).
Lymnaea humilis Say.
Physa cubensis Pfr.
Planorbis tumidus Pfr.
Planorbis alabamensis Pils.
Chondropoma dentatum (Say).
Truncatella bilabiata Pfr.
Truncatella clathrus Lowe.
Littoridina monroensis (Ffld.).

BETWEEN CHESTER SHOALS AND CAPE CANAVERAL.

Polygyra cereolus (Muhlf.).
Polygyra cereolus carpenteriana (Bld.).
Polygyra cereolus volvoxis (Pfr.).
Polygyra uvulifera (Shutt.).
Zonitoides minuscula (Binn.).
Pupoides modicus (Pfr.).
Gastrocopta pentodon (Say).
Gastrocopta pellucida hordeacella (Pils.).
Gastrocopta rupicola (Say).
Melampus coffeus (L.).
Detracia bulloides (Mont.).
Blauneria heteroclita (Mont.).
Chondropoma dentatum (Say).
Truncatella bilabiata Pfr.

Truncatella clathrus Lowe.

Truncatella caribæensis pulchella Pfr.

Amnicola. sp.? A single immature specimen.

CANAVERAL P. O.

Euglandina rosea (Fer.).

Polygyra cereolus septemvolva Say.

Polygyra cereolus volvoxis (Pfr.).

Polygyra cereolus carpenteriana (Bld.).

Pupoides modicus (Pfr.).

Helicina orbiculata Say.

EAU GALLIE.

Polygyra cereolus septemvolva Say.

Polygyra cereolus volvoxis (Pfr.).

Polygyra uvulifera (Shutt.).

Physa cubensis Pfr.

Helicina orbiculata Say var.

ISLAND OF EAU GALLIE.

Polygyra uvulifera (Shutt.).

Praticolella jejuna (Say).

Lymnaea humilis Say.

Physa cubensis Pfr.

PALM BEACH.

Euglandina rosea (Fer.).

Polygyra auriculata Say.

Polygyra cereolus carpenteriana (Bid.).

Strobilops floridana Pils.

Strobilops hubbari (Brown).
Pupoides modicus (Pfr.).
Gastrocopta contracta (Say).
Gastrocopta rupicola Say.
Gastrocopta pellucida hordeacella (Pils.).
Gastrocopta pentodon (Say).
Vertigo milium (Gld.).
Vitrea dalliana ('Simp.' Pils.).
Vitrea indentata (Say).
Zonitoides arborea (Say).
Zonitoides minuscula (Binn.).
Zonitoides minuscula alachuana (Dall).
Guppya gundlachi (Pfr.).
Thysanophora granum (Streb.).
Physa cubensis Pfr.
Helicina orbiculata Say.

LONG KEY.

Euglandina rosea (Fer.).
Polygyra cereolus (Mühlf.).
Polygyra cereolus carpenteriana (Bid.).
Strophia incanum (Binn.).
Pupoides modicus (Pfr.).
Gastrocopta pentodon (Say).
Gastrocopta rupicola (Say)?
Gastrocopta pellucida hordeacella (Pils.).
Gastrocopta pellucida hordeacella (Pils.) var. Small form.
Thysanophora incrustata (Gld.).
Thysanophora granum (Streb.).
Thysanophora dioscoricola (Guppy).
Guppya gundlachi (Pfr.).
Varicella gracillima floridana Pils.
Succinea campestris Say?
Melampus coffeus (L.).
Detracia bulloides (Mont.).
Microtralia minuscula (Ball).

COLLECTING IN DIGBY, NOVA SCOTIA.

BY LILLIAN DYER THOMPSON.

While traveling through Nova Scotia and New Brunswick last summer, we stayed for about six weeks at Digby, N. S. Digby is about 200 miles northeast of Boston, and is situated near the Bay of Fundy, opposite St. John, N. B. The town is located on the southeast shore of the Annapolis Basin,—a sheet of water about twenty miles long and ten miles wide. This basin is connected with the Bay of Fundy by a channel about three-fourths of a mile wide at its greatest width. This channel, known as Digby Gap, is noted for its rapid tides,—the rate of flow through the Gap being about eight miles an hour. The tide fall at Digby is thirty feet. The shores of the Basin are sandy, with the exception of the two rocky promontories on each side of the Gap; the one which is nearest to Digby being Point Prim. The town is on a small peninsula on either side of which are two inlets of the Annapolis Basin, known as the Racquet, on the west, and the Jacquet, on the east of Digby proper. On the ebb tide these are almost dry, exposing long mud flats.

There is one island in the Basin, about opposite the Gap and at the mouth of Bear River, called Bear Island. From this a long bar extends, called Bear Island Bar, which is covered to a depth of about six feet at low water, and is covered with eelgrass.

Near the Yacht Club pier were found many *Polinices heros*, and their red-brown “sand-collars.” In the Jacquet were many *Litorina littorea* and *Litorina rudis*. On the exposed beach, nearer the town, we found *Mytilus edulis*. On the rocks, in the Racquet, we found *Thais lapillus* and a host of *Acmaea testudinalis* ranging in size from one-eighth of an inch to about an inch in diameter. In the mud, at the base of the rocks, were a multitude of *Buccinum undatum*, *Neptunea decemcostata*, ranging in size from one-

eighth of an inch to about an inch in diameter. In the mud, at the base of the rocks, were a multitude of *Buccinum undatum*, *Neptunea decemcostata*, and *Colus stimpsoni*, all alive and half-buried. Some dead specimens of *Aporrhais occidentalis* were also found, five of them being full-grown.

On the suggestion of Capt. Danforth, we constructed a dredge, and endeavored to dredge Bear Island Bar from his motor-boat. Here we found quantities of *Lacuna vincta*, *Alectrion obsoleta*, *Cylichna alba*, and two *Polinices triseriata*.

There were some soldiers encamped at Digby, and they used to gather *Litorina littorea* and steam and eat them, without any flavoring. They sometimes ate *Thais lapillus* also. One day, after a rain, we found two *Helix hortensis* crawling along the road.

A NEW TYPE OF THE NAYAD-GENUS FUSCONAIA. GROUP OF F. BARNESIANA LEA.

BY A. E. ORTMANN.

During the study of the nayad-fauna of the upper Tennessee, the present writer found that there exists, in this region, a peculiar type of shells, belonging to the genus *Fusconaia*, the various forms of which have been described previously under a great number of specific names, which, however, seem to belong all to one species. In addition, among material received from L. S. Frierson from the Ozark Mountains, a form was discovered which presented the same structure.

The oldest name for the upper Tennessee form is *Unio barnesianus* Lea. A more detailed account of its various phases is to be given elsewhere, and it suffices here to mention only those forms which belong here. According to obesity, I distinguish three local, or ecological races:

1. FUSCONAIA BARNESIANA (Lea) 1838.

U. barnesianus Lea, '38. *U. meredithi* Lea, '58. *U. pudicus* Lea, '60. *U. Lyoni* Lea, '65. *U. tellicoensis* Lea, '72. *U. lenticularis* Lea, '72.

As the normal (most abundant) forms we may regard *U. meredithi*, *pudicus* and *lenticularis*, which differ from each other only in the development of the rays (topotypes examined). *U. barnesianus* is a slightly more elongated individual, with poorly developed rays. *U. tellicoensis* (topotypes examined) is a *lenticularis* slightly more swollen; and *U. Lyoni* forms the transition toward var. *tumescens*, having a little more elevated beaks, greater obesity, and rather distinct rays.

2. FUSCONAIA BARNESIANA BIGBYENSIS (Lea) 1841.

U. bigbyensis Lea, '41. *U. estabrookianus* Lea, '45. *U. fassinans* Lea, '68. *Pleurobema fassinans rhomboidea* Simpson, '00.

The most frequent form is *fassinans rhomboidea* (topotypes examined), with rays poorly developed. *U. bigbyensis* has more distinct rays; *U. estabrookianus* (topotypes examined) is an old, overgrown form, without rays; *U. fassinans* is founded upon an individual (type examined, also topotypes), which is exceptionally elongated, without rays.

3. FUSCONAIA BARNESIANA TUMESCENS (Lea) 1845.

U. tumescens Lea, '45. *U. crudus* Lea, '71. *U. radiosus* Lea, '71.

U. tumescens is the most typical form, greatly swollen, with more or less developed rays; *U. radiosus* (type and topotypes examined) is less swollen, but for the rest like *tumescens*; *U. crudus* (topotypes examined) lacks rays, and has much eroded beaks, but stands close to *radiosus*.

The mutual relations of these forms may be understood by the help of the following key. Only the three largest divisions are to be regarded as varieties, in the other forms the characters are merely individual, although specimens representing only one (or a few) of these "forms" often prevail at a given locality.

a₁. Flat, compressed, dia. of shell less than 40 per cent of the length (var. *bigbyensis*).

b₁. No rays, or rays obscure, color of epidermis brown, dark.

c₁. Rhomboid in shape.

d₁. Large. *Estabrookianus*.

d₂. Smaller. *Fassinans rhomboidea*.

c₂. More ovate, tapering behind. *Fassinans*.

b₂. Rays distinct, well developed over most of the disk. Ground color of epidermis lighter. *Bigbyensis*.

a₂. Moderately convex, dia. 40–49 per cent of length. *Barnesiana typica*.

b₁. Beaks not elevated, shape trapezoidal, rhomboid, or subovate.

c₁. Dia. about 41 or 42 per cent; size small.

d₁. Shape somewhat elongate (trapezoidal); rays obscure.
Barnesianus.

d₂. Shape shorter (rhomboidal).

e₁. Rays obscure. *Lenticularis*.

e₂. Rays present, color of epidermis lighter.

f₁. Rays few. *Meredithi*.

f₂. Rays numerous. *Pudicus*.

c₂. Dia. about 45 per cent; larger. Shape subovate. Rays obscure.
Tellicoensis.

Lyoni.

b₂. Beaks more elevated, shape subtriangular. Dia. 46 per cent, with rather distinct rays. *Lyoni*.

a₃. Much swollen, dia. over 50 per cent. Beaks elevated. (var. *Tumescens*).

b₁. Without rays. Dia. 51 per cent. Beaks much eroded. *Crudus*.

b₂. With rays. Dia. about 56 per cent or more.

c₁. Dia. about 56 per cent. *Radiosus*.

c₂. Dia. about 64 per cent. *Tumescens*.

As to the geographical distribution, it should be briefly stated that the swollen forms (a₃) inhabit the largest rivers; the compressed forms (a₁) are found in the headwaters, and the intermediate forms (a₂) belong to the streams of moderate size. Intergrades are frequent.

ANATOMY.^[7]

All these shells have the same, and an extremely characteristic and unique structure of the soft parts, so that there is not the slightest question that they belong together. I have examined the soft parts of some 200 specimens in the field, and over three dozens have been preserved in alcohol, and have been examined at leisure in the laboratory. They include representatives of the three main varieties, and of practically all of the individual variations.

Gravid females have been found on the following dates: May 11, '13; May 15, '13; May 16, '13; May 20, '13; May 20, '14; May 22, '14; May 25, '14; July 5, '13; July 9, '13; July 10, '13; July 13, '13; July 14, '13. *Glochidia* have been observed on May 20, '14 (immature), and July 14, '13. Thus this species evidently is a summer breeder (tachytictic).

The soft parts are those of the genus *Fusconaia*: the *supraanal* is separated from the *anal* by a very short mantle-connection, which is absent (or torn?) in rare cases. Inner lamina of inner gills free from abdominal sac. All four gills are marsupial. Placentae well developed and subcylindrical.

Branchial opening with well developed papillae, *anal* with distinct, but small papillae. *Palpi* subfalciform, posterior margins connected at base only.

While thus the *Fusconaia* structure is typically developed, this species is quite unique in its color. This concerns chiefly the color of the gonads, eggs, and placentae.

The soft parts are often uniformly pale, whitish, but may shade to orange, and the orange is most prominent on foot, adductors, and mantle-margin; but the paler tints prevail, and often the orange is replaced by yellowish or brown. The gills are pale, but are generally suffused with blackish. The gonads are brown to red, mostly of a peculiar dull lavender color in the female, and the latter color, or purplish brown, is the prevailing color of the eggs and placentae. The charged gills become thus rather dark purple, or purple-brown, shading sometimes to dull red or blackish, in other cases to brownish, brownish pink, brick-red, or even pale brown. These are very peculiar tints, by which this species is easily recognized in the field: four

marsupial gills of this blackish-purple color are not known in any other *Nayad*.

Glochidia have been found only in specimens belonging to the headwaters variety (*barnesiana bigbyensis*). They are subelliptical, slightly higher than long, L. 0.15, H. 0.16 mm.

Although a true *Fusconaia*, this species (or group of forms) stands isolated within the genus, in characters of the shell as well as in the soft parts. It differs from the species of the *subrotunda*-group (ind. *ebena*, *pilaris* etc.) very markedly by its smaller size and by the very shallow beak cavities. The forms of the *undulata*-group (incl. *flava*, and the *cuneolus*- and *corforms*) have generally also somewhat deeper beak cavities, and the shell has a more or less distinct posterior ridge, with a flattening or a shallow groove in front of it, characters which are missing in the *barnesiana*-group. As has been pointed out, in the latter group, the color of eggs and placentae is remarkable: in all other forms of *Fusconaia*, this varies from white to bright red.

I introduce here another species, in order to show that the *barnesiana*-type is also represented outside of the Cumberland-Tennessee drainage, namely in the Ozarks.

FUSCONAIA OZARKENSIS (Call) 1887.

F. ozarkensis Call, Pr. U. S. Mus. 10, '87, p. 499, pl. 27. Tr. St. Louis Ac. 7, '95, p. 33, pl. 18. *Lampsilis ozarkensis* Meek & Clark, Bur. Fisher. Doc. no. 759, '12, p. 18. *Pleurobema utterbacki* Frierson, in: Utterback, Naiad. Missouri (Amer. Midland Natural 4, 1916, p. 86, pl. 5, pl. 20, f. 63).

I have specimens from James River, Galena Stone Co., Mo., and White River, Cotter and Norfolk, Baxter Co., Ark., donated by L. S. Frierson and collected by A. A. Hinkley on July 30 and Aug. 2 and 5, '14, A number of specimens (8) were preserved in alcohol, coll. July 30 and Aug 2, which all were gravid females, and one of each date had glochidia. This marks probably the end of the breeding season, and the species is tachytictic.

There is some confusion with regard to this species. After the first description by Call, it has not again been recorded, except by Meek and Clark, and I believe, the identification of these authors (supported by B. Walker) is correct. But I think that other authors have seen this form, but have not recognized it, and, for instance, Simpson's *pannosus* and *subellipticus* (regarded as varieties of *Pleurobema argenteum* and *breve* respectively) are also this. Frierson's *utterbacki* is surely this, since my specimens were thus labeled by Frierson.

Walker, Frierson, and Simpson (in part) believe this to be a *Pleurobema*, and not a *Lampsilis* (see also Simpson, '00, p. 557, and '14, p. 131), and this comes nearest to the truth, in fact, it is the most plausible assumption to be made from the study of the shell alone. The shell "resembles a very elongated *Quadrula coccinea*," according to Meek and Clark, and the comparison with *Pleurobema argenteum* and *breve* (which, by the way, are synonyms), made by Simpson, is significant. We must keep in mind that Call's fig. 4 represents the normal shape of the shell, while his fig. 1 is rather abnormal, and possibly does not belong here at all. These two figures by no means represent the female and male, as Call believes.

The investigation of the soft parts has shown that this actually is a *Fusconaia*. Corresponding, both in soft parts and shell, to the *barnesiana*-type of the upper Tennessee region. *F. ozarkensis* differs from *barnesiana* by the more elongated (subtrapezoidal) outline of the shell, more anterior

beaks, and the weak development of the rays, which are faint at the best, and often entirely absent. A swollen form of it is not known to me, but specimens from White River are slightly more convex than those from James River (farther up). Also Utterback's quotation of Frierson (p. 87, footnote) make it probable that there are differences in obesity.

ANATOMY.

Supraanal opening probably separated from the *anal* by a short mantle-connection, but in all my specimens this is torn by rough handling. Inner lamina of inner gills free from abdominal sac. All four gills marsupial in the female. *Placentae* well developed and *subcylindrical*.

Anal opening with small papillae, branchial opening with well developed papillae. *Palpi* as usual, their posterior margins connected for about one third of their length or less.

As to the color of the soft parts, which is so characteristic in *barnesiana*, not much can be said, since my material has been too long in alcohol. But in most of my specimens the gills are yet distinctly suffused with black. The *placentae* have been rendered whitish, but here and there traces of a dark stain are preserved (which is disappearing gradually). It is quite possible that the color of the *placentae* originally was similar to that of *barnesiana*.

The *glochidia* are subelliptical, slightly higher than long; L. O. 15, B. O. 18, thus agreeing with those of *F. barnesiana*.

NOTE ON THE RELATION OF SNAIL FAUNA TO FLOODS.

BY A. RICHARDS.

During the years 1911 to 1916, while the writer was a member of the faculty of the University of Texas, a series of incidental observations on the snail fauna of Waller Creek was made. These observations have now come to an end due to the change of residence of the observer. It seems not unwise, therefore, to publish a short note on the subject in the hope that the facts recorded, although fragmentary, may have a bearing on the work of some other follower of snail life.

Waller Creek is a small stream near the University of Texas at Austin. It is some four miles in length and empties into the Colorado River at a distance of perhaps two miles below the University. That portion of the stream close to the University between Fifteenth and Twenty-seventh Streets, was most closely observed, but data was also collected from the region below. During the hot months, from about July 1st to October 1st usually, the stream is dry, or water is to be found only in an occasional pool; during the rest of the year the water flows to a depth of a few inches. The bed is scoured out of limestone (Austin Chalk) and has for much of its length a solid flat rock bottom. The banks of the creek have in general a gradual slope. In time of flood and during heavy rains, this stream rises very rapidly and quickly becomes bank-full, so that the water rushes down in a torrent, the roar of which may at times be heard for a distance of some blocks. The fall of the creek is considerable, being about 75 feet in two miles from Twenty-seventh Street to the Colorado, and this fall in connection with the shape of the bed gives to the current in times of flood a tremendous force.

During the first two years of this observation, 1912–1913, the snail population of the creek in its middle stretches was dense. There were in particular two species very thickly represented, *Planorbis lentus* and *Physa halei*. So numerous were they that one could in a few moments within a very few feet gather a pint of either kind. Wherever a little ripple or a tiny waterfall occurred were many snails oriented in relation to the current, their heads pointing into it. Elsewhere in the more quiet water they were also to be found, but in less numbers. These conditions obtained especially in the early spring; as the breeding season, which in that latitude extends over half the year, passes by, the snails of course become much less numerous.

It is to be noted that previous to the time when the snails had become so abundant, there had been no heavy rains of sufficient importance to be recorded since 1908. Excessive rains occurred in May 1908, November and May 1907, June and March 1905, May and April 1904, July and February 1903, July and November 1902. There was a very severe flood in the creek in April 1900. It will be seen that between the time when my observations began and the last excessive rain considerable time had elapsed and the snails had had the opportunity to reinstate themselves in the creek, assuming that they had suffered in those floods as they have done in the later ones.

In the fall of 1913 there were two floods of unusual proportions in the creek. In October it rose very rapidly, but shortly subsided, and in November, at the time when the entire state was visited by the most severe flood since 1869, it was again subjected to a very thorough scouring. Excessive rains fell on several consecutive days, and streams in the entire Colorado watershed were out of their banks.

After the heavy rains of the earlier part of the month there were several days upon which the rainfall, while comparatively light, was sufficient to keep the creek much higher than its normal level. When the water finally subsided to its normal amount the bed was covered completely with a layer of detritus and soft green humus and algae from a half an inch to an inch in thickness. This deposit and the acids formed from it have been the cause of a much more rapid disintegration of the limestone than had been the case in the immediately preceding years. Loose pieces of limestone which were exposed to the action of the water had in many cases fallen apart by the end

of January. Further rises occurred on April 27 and on May 20th, 1914, but were not sufficient to remove all of the accumulated layer of detritus.

In January 1914, a search for snails where they had before been numerous failed to reveal a single specimen of *Planorbis* and less than half a dozen *Physa*. Even in the deeper pools they could scarcely be found. Later in the spring in the lowest part of the stream a number of very small *Physa*, as well as some clusters of eggs were found. The force of the current had been so great as to wash the snails down to the river, and it is possible that the condition of the water due to the decomposing humus may have affected those which were able to escape the flood danger. That some of the *Physa* were left after the flood may be attributed to their pointed shape which decreases the amount of force that the water was able to exert on them as compared with that on the flat *Planorbis* shells. Except as noted above in April and May 1914, the conditions in the creek remained as normal.

In January 1915, *Physa* has again made its appearance in the middle parts of the stream, although in small numbers only. Diligent search, however, failed to reveal a single *Planorbis*. The snails which were present were found only under fair-sized rocks where they were well protected; they did not occur out in the open stream as had been the case when the creek was more densely populated with snails.

In April 1915, on the 22d and 24th, Waller Creek was swept by three scouring floods which devastated the entire bottoms. The water rose higher during the night than it had at any time since 1886; bridges were washed away and much damage done. The bottom of the creek was again washed clean except under the larger stones and in deep holes in the bed. Snails were not observed in any numbers following this flood during the rest of that year. Early in January 1916, however, *Physa* had again appeared fairly abundantly where they had formerly been very thick, in the region just above the University; later in the spring they became quite numerous here. In January of this year the first *Planorbis* were found that had been seen in the creek since the autumn floods of 1913. Between Fifteenth and Sixteenth Streets in a pocket containing good-sized stones over which the water flows rather swiftly a number of specimens were taken, although none were found above or below this locality. It is noted that below this region the creek is frequently covered with oily scum and that it receives the refuse from the adjoining properties. Except after high water which would clean it out, the

creek in this region would hardly be expected to support much snail life. It must be supposed either that the Planorbis had made their way up to Sixteenth Street during the short time following the floods before the water became badly contaminated, or else that in this particular place a few specimens from the previous years had withstood the floods and reproduced themselves in sufficient numbers to be noticeable in January. Of these two suppositions the latter seems much the more rational.

After the flood of 1915 the water subsided very quickly so that a new layer of humus and algae was not deposited, but that the bed was again restored to its former condition of a clean smooth rock bottom. Upon the return to this condition the snail population increased very rapidly, and at the time when the last observation was made in the early summer of 1916, seemed in a fair way to return to the condition of 1912.

It seems to the writer that the slow return of the snails during the year 1914 was due not only to the repeated rains of the fall and winter of 1913–14, but also to the condition of the creek bed at this time. Although the heavy rains of 1915 were more severe, they were confined to one month, and the creek bed was left in a very much cleaner condition than during the preceding year.

The conditions of life which obtain now in Waller Creek are those of a new life region. This must of necessity be so in an intermittent stream to a certain extent, but owing to the flood conditions here they are doubly so.

The chief facts of interest in regard to the habits of snails as shown by these observations, are these: 1, The snails of both species are to be found commonly in uncontaminated water which is running at a fairly rapid rate, and the most common orientation is with the heads pointing up stream; 2, The snail population in any stream is subject to wide fluctuations depending upon flood conditions; 3, Physa because of its shape is less affected by floods than Planorbis; 4, The return of the snail population to a given stream is determined not only by the frequency and severity of the floods, but also by the condition of the bottom of the stream after the subsidence of the high water; 5, The snails thrive best where there is a constantly renewed supply of clean water which contains little decaying vegetation.

Wabash College, Crawfordsville, Ind.

WILLIAM BULLOCK CLARK.

Dr. William Bullock Clark, professor of geology in the Johns Hopkins University, eminent for his contributions to geology, died suddenly from apoplexy on July 27, at his summer home at North Haven, Maine.

Wm. Bullock Clark was born at Brattleboro, Vermont, December 15, 1860. Since 1894 he was professor of geology in Johns Hopkins University. In 1896 Professor Clark organized the Maryland Geological Society, and has been State Geologist since that time. The admirable volumes on paleontology of Maryland, issued under his direction, are widely used by conchologists interested in fossil mollusks. This series of reports will be his enduring memorial. Professor Clark's chief paleontological interest was in the *Echinoidea*, upon which he published several monographs.

PUBLICATIONS RECEIVED.

A MONOGRAPH OF WEST AMERICAN MELANELLID MOLLUSKS. By Paul Bartsch (Proc. U. S. Nat. Museum, Vol. 53, pp. 295–356, pls. 34–39, Aug. 1917). This completes the discussion of the West American mollusks of the super-family Pyramidelloideae, comprising the family Pyramidellidae, which has been previously treated, and the Melanellidae here considered. The former are readily distinguished by having the “nepionic whorls sinistral and tilted; the axis of the early whorls usually being at right angles to that of the succeeding turns, in the first of which the nuclear whorls are frequently quite strongly imbedded.” In the latter the early whorls are dextral and never tilted or immersed. A review of the work done in this group is followed by the descriptions of the species, including forty-nine new species and one new genus *Eulimostraca*. The illustrations are excellent.

NEW AND LITTLE KNOWN SPECIES OF SOUTH AMERICAN MUSSELS OF THE GENUS DIPLODON. By William B. Marshall (Proc. U. S. Nat. Museum, Vol. 53, pp. 381–388, Pls. 50–55, August, 1917). Two new species *Diplodon felipponei* and *D. fortis* are described and figured, together with six species described by Mr. C. T. Simpson in his Descriptive Catalogue of the Naiades and not previously figured.

NOTES ON THE SHELLS OF THE GENUS EPITONIUM AND ITS ALLIES OF THE PACIFIC COAST OF AMERICA. By William H. Dall (Proc. U. S. Nat. Museum, Vol. 53, pp. 471–488, August, 1917). An interesting account of the various groups and subgenera is followed by descriptions of forty-two new species.

The name *Pictoscala* is proposed for a section, type *Scalaria lineata* Say.

STUDIES ON AUSTRALIAN MOLLUSCA. PT. XIII. By C. Hedley (Proc. Linn. Soc. New South Wales, 1916, vol. 41, pt. 4, pp. 680–719, pls. 46–52, issued April 4, 1917). The author's notes under *Tridacna gigantea* Perry are of such general interest that we quote them in part. "Under the name of *Chama gigas* the father of Natural History seems to have embraced the whole of the modern genus *Tridacna*. For the name *gigas*, as restricted to a single species, the candidates are the shell subsequently named *squamosa* by Lamarck and a huge species whose valves in the Ulrica Museum, together weighed 498 pounds.

"After careful examination, Hanley decided that the furbelowed clam, such as Reeve has figured for *T. squamosa*, ought rightly to bear the name *gigas*. He based his verdict on the ground that the actual shell owned by Linné as representing *gigas*, is the Lamarckian *squamosa*, and that to this apply most of the literary references. Linnean contemporaries such as Born, Regenfuss and Chemnitz, while making casual references to the giant, all agree in figuring and describing *squamosa* as the Linnean *gigas*.

"Discriminating in 1819 between the species his predecessors had confused, Lamarck unlawfully used the name *gigas* for the largest form, while for the Linnean *gigas* he proposed *squamosa*. Attentive to the remarks of Hanley, Hidalgo in 1903, renamed the biggest species *T. lamarcki*. But in 1811, Perry had already used the name *Chama gigantea* for 'the largest shell at present known.' As the young of the giant has not yet been traced to the adult, it is still possible that *squamosa* is a juvenile deeper-water form of the large intertidal and abraded *gigantea*.

"The heaviest known are a pair weighing 550 lbs., which Cuvier and Lamarck relate were presented by the Venetian Republic to Francis I. These still exist, their edges bound with brass, as holy-water basins in the cathedral of St. Sulpice, in Paris.

"The photographs of Saville Kent show the giant clams in their natural position on the Great Barrier Reef, where they occur free and exposed at low tide, standing on their umbones, and showing their brightly colored mantle and so-called eyes as they gape."

There are many other interesting notes bearing on nomenclature, and the animals of Australian species. Six new species are described and twenty-nine species figured.—C. W. J.

AN ANNOTATED LIST OF SHELLS FROM NORTHERN MICHIGAN. By Mina L. Winslow (Occasional papers, Mus. Zool., Univ. Mich., No. 42, July 1, 1917) a list of sixty-five species from Schoolcraft, Alger and Chippewa counties, also a list from Isle Royale.

NOTES.

THE OLDROYD COLLECTION.—Mr. and Mrs. J. S. Oldroyd have given their collection of shells to the Leland Stanford Jr. University, and are now permanently employed in the Museum, Mrs. Oldroyd being the curator. The collection has been placed in the Department of Geology and Mining. The Stanford alumni purchased the collection and library of the late Henry Hemphill, which, with the Law collection and several others, forms an unusually fine working series. Mr. and Mrs. Oldroyd have spent about eight weeks at Friday Harbor, Puget Sound and British Columbia making large collections for the Stanford University, California Academy of Science and University of California.

NORTH CAROLINA LAND SHELLS.—The following species of land shells were picked from leaf-mold collected at Spruce Pine, Mitchell Co., North Carolina, by Samuel G. Gordon while on a mineralogical excursion. The specimens are in the collection of the Acad. Nat. Sci., Phila., *Gastrocopta contracta* Say, *G. pentodon* Say, *Circinaria concava* Say, *Polita indentata* Say, *Taxeodonta lamellidens* Pils., *Gastrodonta elliotti* Redf., *G. guldaris* Say, *Euconulus sterkii* Dall., *Punctum pygmæum* Drap., *Carychium exile* Lea.—E. G. VANATTA.

A CORRECTION.—In my little paper, "Descriptions of New West American Marine Mollusks and Notes on Previously Described Forms," Proc. U. S. Nat. Mus., Vol. 52, pp. 670–671, plate 46, figure 2, 1917, I published *Cerithiopsis (Cerithiopsis) helena* from Panama, type Cat. No. 204128, U. S. N. M. Mr. Vignal, of Paris, has been kind enough to call my attention to the fact that the same combination was used by O. Boettger in 1901 for a fossil in his contribution "Zur Kenntniss der Fauna der mittelmioocänen

Schichten von Kosteĵ im Krassò-Szörényer Komitat,” in “Verhandlungen und Mitteilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt,” p. 128, 1901. It is therefore necessary to bestow a new designation on my shell, and it may be known as *Cerithiopsis (Cerithiopsis) anaitis*.—PAUL BARTSCH.

ALBINISTIC EPIPHRAGMOPHORA FIDELIS.—A few days ago, I had two hours in Gladstone Park. The Park is wooded, and there are large moss-covered rocks. *E. fidelis* was out freely. I found one light one. The one sent you some years ago was uniform in color. This one shows two distinct dark narrow bands around the lower whorl, but not showing anywhere else. This is the fourth one I have found in the Park in twenty years or more, though I have been there often.

—J. G. MALONE, *Portland, Ore.*

The many friends of DR. HERMANN VON IHERING will hear with deep regret that he has been removed by political intrigue from his position of Director of the MUSEU PAULISTA at Sao Paulo, Brazil. This museum was founded by Dr. von Ihering. His eminence as a zoologist and unceasing activity as an investigator of the South American fauna, had won for it an honorable place among scientific institutions. We understand that his successors are men without knowledge of the biological sciences. As the only scientific assistant, Mr. Rudolph von Ihering has resigned, it appears that the scientific activity of the State Museum of Sao Paulo has come to an end—a real calamity to American zoology and paleontology.

Dr. von Ihering is located at present at Hansa de Joinville, State of Santa Catharina. He is in good health, and is engaged in the preparation of his work: “Die biogeographischen Grundgesetze,” several chapters of which will deal with mollusks.—H. A. P. & C. W. J.




PLATE VII

- 1, 2. PSORONAIAS KUXENSIS FRIERSON.
3. EPIPHRAGMOPHORA CALLISTODERMA PILS. & FERR.
4. UNIO TETRALASMUS SAY.
- 5, 6. ZACHRYSIA RAMSDENI PILS.
7. Z. EMARGINATA PFR.

[1.](#) NAUTILUS, vol. xx, pp. 58-60.

[2.](#) NAUTILUS, xxvi, pp. 51-55, 1912.

[3.](#) In a later work Ortmann classifies both *ventricosa* and *cohongoronta* as varieties of *ovata* Say.

[4.](#) The relations of mollusks to fish in Oneida Lake. By Frank Collins Baker. Technical Publication No. 4, New York State College of Forestry at Syracuse University. Pp. 366. Syracuse, N. Y., 1916. We are informed that it may be obtained free by those interested in the study of Mollusca by applying to the dean of the college, Dr. Hugh P. Baker.

5. *Pupa griqualandica* M. and P., 1893; the specimens used being from Pretoria.

6. Ann. Mag. N. H. (8), i, p. 76, pl. i, 1908.

7. In NAUTILUS, 28, 1914, p. 31, I have described the anatomy of “*Pleurobema fassinans*.” This is a mistake: the shells examined belong to *Pleurobema* all right, but are the form known as *U. argenteus* Lea, which belongs to the *oviforme-group*, and should be called: *Pleurobema oviforme argenteum* (Lea). These will be treated more fully elsewhere.



TRANSCRIBER'S NOTES

1. Silently corrected typographical errors.
2. Retained anachronistic, non-standard, and uncertain spellings as printed.
3. Footnotes have been re-indexed using numbers and collected together at the end of the last chapter.

*** END OF THE PROJECT GUTENBERG EBOOK THE NAUTILUS.
VOL. XXXI, NO. 2, OCTOBER 1917 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the

collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with

this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official

version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, “Information about donations to the Project Gutenberg Literary Archive Foundation.”
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.

- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES

EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a)

distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact

links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility: www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.