Individual Case Safety Report (ICSR)

Final Project — Pharmacovigilance Assessment

SIMULATED EDUCATIONAL CASE — NOT A REAL PATIENT REPORT

1. Case Summary

Field	Assessment
Case ID	ICSR-TRAIN-2026-001
Patient	Adult female, 54 years
Suspected medicine	Amoxicillin/Clavulanic acid 625 mg tablet
Indication	Acute bacterial sinusitis
Dose	625 mg orally three times daily
Reaction	Generalised urticaria with facial/lip swelling
Onset	05-Sep-2026
Outcome	Recovered

2. ADR Identification

A suspected hypersensitivity-type reaction consisting of generalised urticaria with facial/lip swelling occurred during treatment with amoxicillin/clavulanic acid. The temporal relationship and positive dechallenge support consideration as an ADR in this simulated case.

3. Seriousness and Severity

Seriousness: Serious — medically important, requiring urgent medical intervention/observation.
Severity: Moderate. The reaction required active treatment but the simulated case does not describe shock, respiratory arrest, intensive-care admission or permanent disability.

4. Expectedness

Expected / labelled for this training scenario. In real pharmacovigilance, expectedness must be checked against the current approved product information/reference safety information.

5. Dechallenge / Rechallenge

Dechallenge: Positive — symptoms improved after discontinuation and treatment.
Rechallenge: Not performed because intentional re-exposure was not clinically justified.

6. WHO-UMC Causality Assessment

Criterion	Finding
Temporal relationship	Reasonable
Alternative causes	No strong alternative cause identified
Dechallenge	Positive and clinically reasonable
Rechallenge	Not performed; not required for Probable/Likely
Overall category	Probable/Likely

7. Clinical Narrative

A 54-year-old female was treated with amoxicillin/clavulanic acid 625 mg orally three times daily for acute bacterial sinusitis beginning 01-Sep-2026. On 05-Sep-2026, during treatment, she developed generalised urticaria accompanied by facial/lip swelling. The suspected medicine was discontinued. Antihistamine and corticosteroid therapy were administered and the patient was observed. Symptoms improved and the patient recovered. Rechallenge was not performed. No strong alternative cause was identified in this simulated case. The overall relationship was assessed as Probable/Likely according

to the WHO-UMC approach.

8. MedDRA-Style Coding

Training coding concepts: Urticaria; Facial swelling/Lip swelling; hypersensitivity reaction. Exact MedDRA LLT/PT selection should be performed using the current licensed MedDRA browser and the verbatim source wording.

9. Drug Safety Associate Conclusion

The case is sufficiently described for this simulated educational ICSR. It is classified as serious due to medical importance, moderate in severity, expected for the training scenario, and Probable/Likely in causality. The patient recovered following withdrawal and treatment. A single case does not establish causality or a safety signal; aggregate evidence and further case assessment would be required in real pharmacovigilance.

10. References

WHO-Uppsala Monitoring Centre (UMC), Standardised Case Causality Assessment; ICH E2A, Clinical Safety Data Management: Definitions and Standards for Expedited Reporting; Indian Pharmacopoeia Commission / PvPI pharmacovigilance resources.