

TUTORIAL-I

NAME : PRALABH AGARWAL

SEC: D (CSE)

SEM: IV

C. Roll no: 29

Univ. roll: 2016908


Date: 10/03/2022

Ques: Asymptotic Notations.

↳ Tending to infinity

They help you find the complexity an algorithm when input is very large.

i) Big O (O)



$$f(n) = O(g(n))$$

$$f(n) \leq c \cdot g(n)$$

$$\forall n \geq n_0$$

for some constant $c > 0$
 $\Rightarrow g(n)$ is 'tight' upper bound of $f(n)$

ii) Big Omega (Ω)


$$f(n) = \Omega(g(n))$$

$g(n)$ is 'tight' lower bound of $f(n)$

$$f(n) = \Omega(g(n))$$

$$f(n) \geq c \cdot g(n)$$

$\forall n \geq n_0$ for some constant $c > 0$



iii) Theta (Θ)

$$f(n) = \Theta(g(n))$$

$g(n)$ is both 'tight' upper & lower bound of $f(n)$


$$f(n) = \Theta(g(n))$$

iff

$$c_1 \cdot g(n) \leq f(n) \leq c_2 \cdot g(n)$$

$$\forall n \geq \max(n_1, n_2)$$

for some constant $c_1 > 0$ & $c_2 > 0$



4) small $O()$

$$f(n) = O(g(n))$$


$g(n)$ is upper bound of $f(n)$

$$f(n) = O(g(n))$$

when $f(n) < c \cdot g(n)$

$$\forall n > n_0$$

$$\Delta \forall c > 0$$



5) small omega (ω)

$$f(n) = \omega(g(n))$$


$g(n)$ is lower bound of $f(n)$

$$f(n) = \omega(g(n))$$

when $f(n) > c \cdot g(n)$

$$\forall n > n_0$$

$$\Delta \forall c > 0$$



Ques 2 What should be time complexity of
 $\text{for}(i=1 \text{ to } n) \{ i = i+2 \}$

$\text{for}(i=1 \text{ to } n) \quad // \quad i = 1, 2, 4, 8, \dots, n$
 $\{ i = i+2 \} \quad // \quad O(1)$

$$\Rightarrow \sum_{i=1}^n 1 + 2 + 4 + 8 + \dots + n$$

GP kth value $\Rightarrow T_k = ar^{k-1}$
 $\Rightarrow 1 \times 2^{k-1}$

$$\Rightarrow n = 2^{k-1}$$

$$\Rightarrow 2n = 2^k$$

$$\Rightarrow \log_2 n = k \log_2 2$$

$$\Rightarrow \log_2 + \log n = k \log 2$$

$$\Rightarrow \log n + 1 = k$$

$$\Rightarrow O(k) = O(1 + \log n)$$

$$= \underline{O(\log n)}$$

Q-3 $T(n) = \{ 3T(n-1) \text{ if } n > 0, \text{ otherwise } 1 \}$

$$T(n) = 3T(n-1) \quad \text{--- (1)}$$

~~Recursion~~

put $n = n-1$

$$T(n-1) = 3T(n-2) \quad \text{--- (2)}$$

from 1 & 2

$$\Rightarrow T(n) = 3(3T(n-2))$$

$$\dots = 9T(n-2) \quad \text{--- (3)}$$

~~Defera~~

putting $n = n-2$ in ①

$$T(n) = 3(T(n-2)) \quad - ④$$

$$\Rightarrow T(n) = 27(T(n-3))$$

$$\Rightarrow T(n) = 3^k(T(n-k))$$

putting $n-k=0$

$$\Rightarrow n=k$$

$$\Rightarrow T(n) = 3^n [T(0-2)]$$

$$\Rightarrow T(n) = 3^n T(0)$$

$$\Rightarrow T(n) = 3^n \times 1 \quad [T(0)=1]$$

$$\Rightarrow \underline{\underline{T(n) = O(3^n)}}$$

4) $T(n) = \{ 2T(n-1) - 1 \text{ if } n > 0, \text{ otherwise } 1 \}$

$$T(n) = 2T(n-1) - 1 \quad - ①$$

Let $n = n-1$

$$\Rightarrow T(n-1) = 2T(n-2) - 1 \quad - ②$$

\Rightarrow from ① & ②

$$\Rightarrow T(n) = 2[2T(n-2) - 1] - 1$$

$$\Rightarrow T(n) = 4T(n-2) - 2 - 1 \quad - ③$$

Let $n = n-2$

$$\Rightarrow T(n-2) = 2T(n-3) - 1 \quad - ④$$

from ③ & ④

$$\Rightarrow T(n) = 4[2T(n-3) - 1] - 2 - 1$$

$$\Rightarrow T(n) = 8T(n-3) - 4 - 2 - 1$$

Alafana

$$\Rightarrow T(n) = 2^k T(n-k) - 2^{k-1} - 2^{k-2} - \dots - 1$$

$$\Rightarrow AP = 2^{k-1} + 2^{k-2} + 2^{k-3} + \dots + 1$$

$$a = 2^{k-1}$$

$$r = 1/2$$

$$\Rightarrow S_k = \frac{a(1-r^n)}{1-r}$$

$$= \frac{2^{k-1}(1-(1/2)^n)}{1-1/2}$$

$$= 2^k (1 - (1/2)^n)$$

$$= 2^k - 1$$

$$\text{Let } n-k = 0$$

$$\Rightarrow n = k$$

$$\Rightarrow T(n) = 2^n T(n-n) - (2^n - 1)$$

$$\Rightarrow T(n) = 2^n \cdot 1 - (2^n - 1)$$

$$\Rightarrow T(n) = 2^n - (2^n - 1)$$

$$\Rightarrow T(n) = 1$$

Done

Q5-5

what should be time complexity of

```
int i=1, s=1;
while (s <= n)
{
    i++; s = s*i;
    printf("#");
}
```

i = 1 2 3 4 5 6 ... n

s = 1 * 3 * 6 * 10 * 15 * 21 ... n

~~also does~~

Sum of s = 1 + 3 + 6 + 10 + ... + T_n - (1)

also s = 1 + 3 + 6 + 10 + ... + T_m + T_n - (2)

from (1) - (2)

$$0 = 1 + 2 + 3 + 4 + \dots + n - T_n$$

$$\Rightarrow T_k = 1 + 2 + 3 + 4 + \dots + k$$

$$\Rightarrow T_k = \frac{1}{2} k(k+1)$$

$$\Rightarrow T(k)$$

\Rightarrow for k iterations.

$$1 + 2 + 3 + \dots + k \leq n$$

$$\Rightarrow \frac{k(k+1)}{2} \leq n$$

$$\Rightarrow \frac{k^2 + k}{2} \leq n$$

$$\Rightarrow O(k^2) \leq n$$

$$\Rightarrow k = O(\sqrt{n})$$

$$\Rightarrow T(n) = O(\sqrt{n})$$

~~Answer~~

Ques 6 Time complexity of -

```
void fn(int n)
```

```
{ int i, count = 0;
```

```
  for (i = 1; i * i <= n; ++i)
```

```
    count++
```

// O(n)

```
}
```

$$\text{as } i^2 \leq n$$

$$\Rightarrow i \leq \sqrt{n}$$

$$i = 1, 2, 3, 4, \dots, \sqrt{n}$$

$$\sum_{i=1}^{\sqrt{n}} 1 + 2 + 3 + 4 + \dots + \sqrt{n}$$

$$\Rightarrow T(n) = \frac{\sqrt{n} \times (\sqrt{n} + 1)}{2}$$

$$\Rightarrow T(n) = \frac{n \times \sqrt{n}}{2}$$

$$\Rightarrow T(n) = \underline{\underline{O(n)}}$$

Ques 7 Time complexity of :-

```
void fn(int n)
```

```
{ int i, j, k, count = 0;
```

```
  for (i = n/2; i <= n; ++i)
```

```
    for (j = 1; j <= n; j = j * 2)
```

```
      for (k = 1; k <= n; k = k * 2)
```

```
        count++;
```

```
}
```

Answer

for $k = k \times 2$

$k = 1, 2, 4, 8, \dots, n$

\Rightarrow G.P. $a = 1, r = 2$

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

$$= \frac{1(2^k - 1)}{2 - 1}$$

$$n \Rightarrow 2^k$$

$$\Rightarrow \underline{\underline{\log n = k}}$$

$$\Rightarrow$$

1	1	k
1	$\log n$	$\log n * \log n$
2	$\log n$	$\log n * \log n$
\vdots	\vdots	\vdots
n	$\log n$	$\log n * \log n$

$$\Rightarrow O(n * \log n * \log n)$$

$$\Rightarrow \underline{\underline{O(n \log^2 n)}}$$

Shubhendra

8) Time Complexity of

```
function int(n)
```

```
{ int(n-1)
```

```
  return;
```

```
  for (i=1 to n)
```

```
    for (j=1 to n)
```

```
      print('*');
```

```
    }
```

```
  }
```

```
function(n-3);
```

```
}
```

// $O(1)$

// $i = 1, 2, 3, 4, \dots, n \Rightarrow O(n)$

// $j = 1, 2, 3, 4, \dots, n \Rightarrow O(n^2)$

$T(n/3)$

$$\Rightarrow T(n) = T(n/3) + n^2$$

$$\Rightarrow a=1, \quad b=3, \quad f(n) = n^2$$

$$c = \log_3 1 = 0$$

$$\Rightarrow n^0 = 1 > (f(n) = n^2)$$

$$\Rightarrow \underline{T(n) = \Theta(n^2)}$$

Q. 10

Ques 9

Time complexity of -
void function (int n)


{ for (i=1 to n)

{ for (j=1; j<=n; j=j+i)

print ("*")

11 06)

11 06)



for i=1 $\Rightarrow j = 1, 2, 3, 4, \dots, n = n$
 for i=2 $\Rightarrow j = 1, 3, 5, \dots, n = n/2$
 for i=3 $\Rightarrow j = 1, 4, 7, \dots, n = n/3$
 \vdots
 for i=n $\Rightarrow j = 1, \dots, 1$

$$\Rightarrow \sum_{j=n}^1 n + \frac{n}{2} + \frac{n}{3} + \frac{n}{4} + \dots + 1$$

$$\Rightarrow \sum_{j=n}^1 n \left[1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} \right]$$

$$\sum_{j=n}^1 n [\log n]$$

$$\Rightarrow T(n) = [n \log n]$$

$$T(n) = O(n \log n)$$

Solution

Q-10 for functions, n^k & c^n , what is the asymptotic relation between these functions?

assume that $k \geq 1$, & $c \geq 1$ are constant.

Find out the value of c & n_0 for which relation holds

as given n^k & c^n

at

relation b/w n^k & c^n is

$$n^k = O(c^n)$$

are ~~of order~~ $n^k \gg n^k$

as $n^k \leq ac^n$

$\forall n \geq n_0$ & some constant ~~near~~ $a > 0$

for $n_0 = 1$

$c = 2$

$$\Rightarrow 1^k \leq a \cdot 2^1$$

$$\Rightarrow \underline{n_0 = 1 \text{ \& } c = 2}$$

~~Defence~~