
a
Heart rate (bpm)
ln EDA tonic (SCL)
EDA phasic (SCR)
Skin temperature (°C)
ln HRV SDNN (ms)
ln HRV RMSSD (ms)
ln HRV LF power (ms²)
ln HRV HF power (ms²)
ln HRV LF/HF ratio
HRV LF (nu)
HRV HF (nu)
ln HRV VLF power (ms²)
ln HRV total power (ms²)
Pupil diameter (mm)
Pupil change rate
Blink duration (s)
Fixation duration (s)
Fixation rate (/s)
Saccade duration (s)
Saccade amplitude (°)
Saccade peak velocity (°/s)
Gaze velocity (°/s)
Gaze dispersion
Gaze distance from centre (px)
Gaze centrality
Eyelid aperture (mm)
General comfort (1–7)
Pleasantness (1–5)
State anxiety STAI-6 (1–5)
Restorativeness PRS-11 (1–5)
rmcorr r
Black box / bold = BH-FDR q < 0.05 (one family of 104 tests). Within-person rmcorr across the 4 phases on 44 participant × phase means (ln applied per 10 s epoch for log outcomes, then averaged). Vertical line separates Empatica (left) and eye-tracker (right) outcomes.
b
r_rm = -0.77 [-0.88, -0.57], q = 7.2e-05, n = 39 cells / 10 participants
Bike-Urban (BikeU)
Walk-Urban (WalkU)
Bike-Green (BikeG)
Walk-Green (WalkG)
Restorativeness PRS-11 (1–5) (within-person deviation)
Blink duration (s) (within-person deviation)
c
r_rm = 0.67 [0.40, 0.83], q = 2.0e-03, n = 39 cells / 10 participants
Restorativeness PRS-11 (1–5) (within-person deviation)
Pupil diameter (mm) (within-person deviation)