

Introduction to Quantum Information and Communication

Theory Assignment-2

Moida Praneeth Jain, 2022101093

Exercise 4.1.3

Given:

- A is a square operator acting on Hilbert space \mathcal{H}_S
- I_R is the identity operator acting on a Hilbert space \mathcal{H}_R isomorphic to \mathcal{H}_S
- $|\Gamma\rangle_{RS}$ is the unnormalized maximally entangled vector.

To Prove:

$$\text{Tr}\{A\} = \langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS}$$

Proof:

In the computational basis

$$|\Gamma\rangle_{RS} = \sum_{i=0}^{d-1} |i\rangle_R |i\rangle_S$$

$$\langle \Gamma |_{RS} = \sum_{i=0}^{d-1} \langle i |_R \langle i |_S$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \left(\sum_{i=0}^{d-1} \langle i |_R \langle i |_S \right) (I_R \otimes A_S) \left(\sum_{j=0}^{d-1} |j\rangle_R |j\rangle_S \right)$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \left(\sum_{i=0}^{d-1} \langle i |_R \langle i |_S \right) \left(\sum_{j=0}^{d-1} (I_R \otimes A_S) (|j\rangle_R \otimes |j\rangle_S) \right)$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \left(\sum_{i=0}^{d-1} \langle i |_R \langle i |_S \right) \left(\sum_{j=0}^{d-1} (I_R |j\rangle_R) \otimes (A_S |j\rangle_S) \right)$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \left(\sum_{i=0}^{d-1} \langle i |_R \langle i |_S \right) \left(\sum_{j=0}^{d-1} |j\rangle_R \otimes (A_S |j\rangle_S) \right)$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \sum_{i,j=0}^{d-1} (\langle i |_R \otimes \langle i |_S) (|j\rangle_R \otimes (A_S |j\rangle_S))$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \sum_{i,j=0}^{d-1} (\langle i | j \rangle_R \otimes \langle i |_S A_S | j \rangle_S)$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \sum_{i,j=0}^{d-1} (\delta_{i,j} \otimes \langle i |_S A_S | j \rangle_S)$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \sum_{i=0}^{d-1} \langle i |_S A_S | i \rangle_S$$

$$\langle \Gamma |_{RS} I_R \otimes A_S | \Gamma \rangle_{RS} = \text{Tr}\{A\}$$

Hence, proven.

Exercise 4.1.16

Given:

- Commuting projectors Π_1 and Π_2
- $0 \leq \Pi_1, \Pi_2 \leq I$

To Prove:

For arbitrary density operator ρ

$$\text{Tr}\{(I - \Pi_1 \Pi_2)\rho\} \leq \text{Tr}\{(I - \Pi_1)\rho\} + \text{Tr}\{(I - \Pi_2)\rho\}$$

Proof:

TO DO

Exercise 4.2.2

Given:

- Ensemble $\{p_X(x), \rho_x\}$ of density operators
- POVM with elements $\{\Lambda_x\}$
- Operator τ such that $\tau \geq p_X(x)\rho_x$

To Prove:

$$\text{Tr}\{\tau\} \geq \sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\}$$

Proof:

$$\sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\} = \sum_x \text{Tr}\{\Lambda_x p_X(x) \rho_x\}$$

$$\sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\} \leq \sum_x \text{Tr}\{\Lambda_x \tau\}$$

$$\sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\} \leq \text{Tr}\left\{\sum_x \Lambda_x \tau\right\}$$

$$\sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\} \leq \text{Tr}\left\{\tau \sum_x \Lambda_x\right\}$$

$$\sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\} \leq \text{Tr}\{\tau I\}$$

$$\sum_x p_X(x) \text{Tr}\{\Lambda_x \rho_x\} \leq \text{Tr}\{\tau\}$$

Hence, proven.

Now for the case of encoding n bits into a d -dimensional subspace.

$$\{2^{-n}, \rho_i\}_{i \in \{0,1\}^n}$$

Consider

$$p_X(x)\rho_x = 2^{-n}\rho_i$$

$$p_X(x)\rho_x = 2^{-n} \sum_j \lambda_j |j\rangle\langle j|$$

$$2^{-n}I - p_X(x)\rho_x = 2^{-n}I - 2^{-n} \sum_j \lambda_j |j\rangle\langle j|$$

$$2^{-n}I - p_X(x)\rho_x = 2^{-n} \sum_j |j\rangle\langle j| - 2^{-n} \sum_j \lambda_j |j\rangle\langle j|$$

$$2^{-n}I - p_X(x)\rho_x = 2^{-n} \sum_j (1 - \lambda_j) |j\rangle\langle j|$$

Since $0 \leq \lambda_j \leq 1 \forall j$, $1 - \lambda_j \geq 0 \forall j$. All the eigenvalues of the matrix in LHS are non-negative.

$$2^{-n}I - p_X(x)\rho_x \geq 0$$

$$2^{-n}I \geq p_X(x)\rho_x$$

\therefore We consider $\tau = 2^{-n}I$

Now, we know that the probability of success is upper bounded by $\text{Tr}\{\tau\}$

$$\text{Tr}\{\tau\} = \text{Tr}\{2^{-n}I\}$$

$$\text{Tr}\{\tau\} = 2^{-n} \text{Tr}\{I\}$$

Since I is d -dimensional,

$$\text{Tr}\{\tau\} = d2^{-n}$$

Thus, the expected success probability is bounded above by $d2^{-n}$

Exercise 4.3.1

Given:

- A' has a Hilbert space structure isomorphic to that of system A
- $\forall x, y \ F_{AA'} |x\rangle_A |y\rangle_{A'} = |y\rangle_A |x\rangle_{A'}$

To Prove:

$$P(\rho_A) = \text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\}$$

Proof:

$$\rho_A = \sum_i \lambda_i |i\rangle_A \langle i|_A$$

$$\rho_{A'} = \sum_j \lambda_j |j\rangle_{A'} \langle j|_{A'}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\{F_{AA'}(\rho_A \otimes \rho_{A'})\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\left\{F_{AA'}\left(\left(\sum_i \lambda_i |i\rangle_A \langle i|_A\right) \otimes \left(\sum_j \lambda_j |j\rangle_{A'} \langle j|_{A'}\right)\right)\right\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\left\{F_{AA'}\left(\sum_{i,j} \lambda_i \lambda_j |i\rangle_A \langle i|_A \otimes |j\rangle_{A'} \langle j|_{A'}\right)\right\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\left\{F_{AA'}\left(\sum_{i,j} \lambda_i \lambda_j |i\rangle_A \langle i|_A \otimes |j\rangle_{A'} \langle j|_{A'}\right)\right\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\left\{F_{AA'}\left(\sum_{i,j} \lambda_i \lambda_j (|i\rangle_A \langle j|_{A'}) (\langle i|_A \langle j|_{A'})\right)\right\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\left\{\left(\sum_{i,j} \lambda_i \lambda_j (F_{AA'} |i\rangle_A \langle j|_{A'}) (\langle i|_A \langle j|_{A'})\right)\right\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\left\{\sum_{i,j} \lambda_i \lambda_j (|j\rangle_A \langle i|_{A'}) (\langle i|_A \langle j|_{A'})\right\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \sum_{i,j} \lambda_i \lambda_j \text{Tr}\{(|j\rangle_A \langle i|_{A'}) (\langle i|_A \langle j|_{A'})\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \sum_{i,j} \lambda_i \lambda_j \text{Tr}\{(\langle i|_A \langle j|_{A'}) (|j\rangle_A \langle i|_{A'})\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \sum_{i,j} \lambda_i \lambda_j \text{Tr}\{ \langle i|j\rangle_A \otimes \langle j|i\rangle_{A'} \}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \sum_{i,j} \lambda_i \lambda_j \langle i|j\rangle_A \langle j|i\rangle_{A'}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \sum_{i,j} \lambda_i \lambda_j \delta_{i,j}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \sum_i \lambda_i^2$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = \text{Tr}\{\rho_A^2\}$$

$$\text{Tr}\{(\rho_A \otimes \rho_{A'})F_{AA'}\} = P(\rho_A)$$

Hence, proven.