BTVNANOCOMMISSIONING · ZB_ZC_RATIO WORKFLOW

Z(→ℓℓ) + b/c AK4 jets in Run 3

Summer22 DY→ℓℓ MC with AK4 jets, 20 files per sample: samples, cutflow, kinematics, flavour and pileup checks

19 October 2026

Campaign and setup

Goal: measure σ(Z+b) / σ(Z+c) with an AK4 jet recoiling against Z→ee/μμ (switched from the boosted AK8 jet after the group meeting). First step: tune the selection on DY→ℓℓ MC.

MC CAMPAIGN

Summer22

Run3Summer22 = 2022 eras C–D (pre-EE)

FORMAT

NanoAODv12

130X MC conditions

ENERGY (√s)

13.6 TeV

Run 3 pp collisions

NORMALIZATION

7.98 fb^{-1}

2022 C–D golden JSON; brilcalc check pending

MC only, no data yet. Processed: 20 files per sample (the first 20 files of each sample in the fileset, 100 of 1887 DY files; run_zbzc_ak4_20files.sh). Code: BTVNanoCommissioning, branch coffea_machine, workflow Zb_Zc_ratio (AK4, commit 31ba6bc, with the GenJet-match counters).

MC samples: DY→ℓℓ, stitched in pT(ℓℓ)

```
/DYto2L-2Jets_MLL-50[_PTLL-X]_TuneCP5_13p6TeV_amcatnloFXFX-pythia8/
  Run3Summer22NanoAODv12-130X_mcRun3_2022_realistic_v5-vN/NANOAODSIM
```

Sample	LHE pT(ℓℓ) used	σ [pb]	DAS files	DAS events	20 files: events
Inclusive (MLL-50)	< 100 GeV	6346	525	93.8 M	3,266,113
PTLL-100	100–200 GeV	105.0	418	56.4 M	2,933,762
PTLL-200	200–400 GeV	11.24	335	44.7 M	2,735,083
PTLL-400	400–600 GeV	0.643	424	41.6 M	1,560,230
PTLL-600	> 600 GeV	0.0994	185	9.50 M	1,236,732

PTLL-X samples are inclusive above X, so each is kept to its own LHE pT(ℓℓ) slice. σ(PTLL-X) = 6346 pb × fraction of inclusive events with pT(ℓℓ) > X (measured on 27.7 M inclusive events). "20 files: events" = events of 20 files inside the slice. With no boosted-jet requirement the inclusive sample (pT(ℓℓ) < 100 GeV) gives 87% (Z→ee) / 89% (Z→μμ) of the scaled yield from only 5–6% of the raw selected events, so its large per-event weights limit the MC statistics.

Event selection, per channel

Z→ee / Z→μμ (unchanged)

- Trigger: HLT_Ele30_WPTight_Gsf / HLT_IsoMu24
- Leptons pT > 25 GeV, |η| < 2.4; leading pT > 35 GeV
- e: tight cut-based ID, IP cuts, EB–EE gap veto
- μ: tight ID, PF isolation < 0.15
- Opposite sign (OS)
- 71 < m$\ell\ell$ < 111 GeV
- PuppiMET < 50 GeV

Leading AK4 jet ("jet" step)

- AK4 PUPPI jets, ΔR > 0.4 from selected leptons
- Tight-lepton-veto jet ID (with NanoAODv12 jetId fix)
- pT > 30 GeV, |η| < 2.5; ≥ 1 such jet, leading one used
- No Z pT requirement

Weights and flavour

- genWeight × pileup × lepton ID/iso/reco SFs
- No trigger SF yet
- Jet flavour from hadronFlavour: b (5), c (4), light; no bb/cc split for AK4
- Also labelled by |partonFlavour|: b, c, uds, g, none (pages 21+, CMS recommendation on p. 21)
- Tagger: PNet AK4 B, CvL, CvB

Cutflow, Z→ee: raw MC events per step

Sample	all	trigger	leptons	OS	Zmass	MET	jet	jet / MET
Inclusive	3,266,113	541,950	165,143	164,108	158,226	150,078	39,242	26.1%
PTLL-100	2,933,762	840,163	238,941	236,499	218,939	193,805	186,102	96.0%
PTLL-200	2,735,083	929,177	344,674	339,940	309,699	256,244	252,727	98.6%
PTLL-400	1,560,230	576,282	264,078	259,407	233,151	171,946	171,561	99.8%
PTLL-600	1,236,732	306,140	138,357	135,652	112,766	73,911	73,866	99.9%

723,498 raw MC events pass the full Z→ee selection; the most (35%) come from PTLL-200. The AK4 jet step keeps 26% of inclusive (low-pT) Z events and ≥ 96% at pT($\ell\ell$) > 100 GeV.

Summer22 DY→$\ell\ell$ MC · 20 files per sample · AK4 jets · "all" = events in the sample's LHE pT($\ell\ell$) slice · 5

Cutflow, Z→μμ: raw MC events per step

Sample	all	trigger	leptons	OS	Zmass	MET	jet	jet / MET
Inclusive	3,266,113	690,771	290,730	290,720	280,562	266,331	68,684	25.8%
PTLL-100	2,933,762	886,704	372,657	372,607	344,587	305,050	292,532	95.9%
PTLL-200	2,735,083	949,986	504,655	504,584	457,929	375,168	369,611	98.5%
PTLL-400	1,560,230	591,000	365,818	365,780	324,198	226,470	225,901	99.7%
PTLL-600	1,236,732	435,749	279,760	279,731	241,960	138,241	138,146	99.9%

1,094,874 raw MC events pass the full Z→μμ selection; the most (34%) come from PTLL-200. The AK4 jet step keeps 26% of inclusive (low-pT) Z events and ≥ 96% at pT(ℓℓ) > 100 GeV.

Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · "all" = events in the sample's LHE pT(ℓℓ) slice · 6

Yields by AK4 jet flavour (7.98 fb^{-1})

Channel	light	c	b	total
Z→ee	373,416.5	34,579.9	16,853.2	424,849.6
fraction	0.879	0.081	0.040	
Z→μμ	674,884.4	62,972.2	32,587.5	770,444.2
fraction	0.876	0.082	0.042	

20 files per sample (100 of 1887 DY files)

723,498 (Z→ee) and 1,094,874 (Z→μμ) raw events pass. The inclusive sample (pT($\ell\ell$) < 100 GeV) still carries large per-event weights, so the far tails (e.g. N_jet ≥ 6) rest on a few events. No b-tag requirement applied yet.


AK4: 1 file vs 20 files per sample

	1 file Z→ee	20 files Z→ee	1 file Z→μμ	20 files Z→μμ
Raw events selected	21,367	723,498	33,106	1,094,874
Scaled yield	403,007.9	424,849.6	784,702.2	770,444.2
b fraction	0.042	0.040	0.045	0.042
c fraction	0.106	0.081	0.083	0.082
light fraction	0.852	0.879	0.872	0.876

Same selection and code; 20× more files per sample, about 33× more raw events. The 1-file test had only N_eff ≈ 466 (Z→ee) / 915 (Z→μμ) effective events. The largest change is the Z→ee c fraction, 0.106 → 0.081, 1.7σ of the 1-file statistical error, so the 1-file numbers were a fair estimate. The AK8 vs AK4 comparison (1-file test only) is in the test deck.

Dilepton mass


m(ℓℓ) after the full selection (71–111 GeV window): peaks at the Z mass in both channels. Linear y-axis.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 9

Dilepton pT


pT(ℓℓ) of the Z candidate after the full selection. No Z pT cut: dominated by low pT, unlike the boosted AK8 selection.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 10

Leading-lepton pT


Leading lepton of the Z candidate, after the full selection (pT > 35 GeV required).



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 11

Subleading-lepton pT


Subleading lepton of the Z candidate, after the full selection (pT > 25 GeV required).



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 12

Leading AK4 jet pT


Leading lepton-cleaned AK4 jet with tight-lepton-veto ID (pT > 30 GeV required).



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 13

Leading AK4 jet η


Pseudorapidity of the selected AK4 jet (|η| < 2.5 required).



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 14

AK4 jet multiplicity


Number of selected AK4 jets (pT > 30 GeV, |η| < 2.5, ID, lepton-cleaned) in events passing the full selection.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 15

LHE pT($\ell\ell$)


Generator-level pT($\ell\ell$) after the full selection: checks the sample stitching (slices at 100/200/400/600 GeV).



Summer22 DY→$\ell\ell$ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 16

Leading AK4 jet: PNet B


ParticleNet b-tag score of the selected jet; b jets pile up near 1, light jets near 0.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 17

Leading AK4 jet: PNet CvL


ParticleNet charm-vs-light score of the selected jet.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 18

Leading AK4 jet: PNet CvB


ParticleNet charm-vs-b score of the selected jet.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · stacked by leading-jet flavour; lower: fraction per bin · Z→ee left, Z→μμ right · 19

PNet charm-tagging plane: CvL vs CvB

Selected AK4 jets per flavour, both channels summed, normalized to unit area. b jets sit at high CvL and low CvB, light jets at low CvL and high CvB. c jets are at high CvB, but a large part of them overlaps the light-jet corner (low CvL), with the rest at high CvL.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · Z→ee + Z→μμ · 20

Jet flavour truth: CMS vs our choice

CMS / BTV recommendation

- Hadron-based flavour by ghost association: generated B and C hadrons get their momenta scaled by 10^{-18} and are re-clustered with the jet constituents (same algorithm, same R), so the jet is unchanged but now contains the hadron "ghosts"
- hadronFlavour = 5 if ≥ 1 B hadron, 4 if ≥ 1 C hadron and no B, else 0 (udsg)
- PAT default since CMSSW_7_0_5_patch1; in NanoAOD as Jet_hadronFlavour
- BTV b/c-tag SFs are defined for it: correctionlib input "flavor: hadron flavor definition: 5=b, 4=c, 0=udsg" (2022_Summer22 btagging / ctagging)
- partonFlavour (ghost partons, hadron flavour has priority) only to split light jets into uds / g; the ΔR < 0.3 parton matching is the Run 1 legacy definition

What Zb_Zc_ratio (AK4) does

- Takes Jet_hadronFlavour of the leading selected AK4 jet straight from NanoAODv12: 5 → b, 4 → c, else → light (Zb_Zc_ratio.jet_flavour)
- No re-clustering or ΔR matching of our own: same ghost-association labels as CMSSW
- No b/bb or c/cc split: NanoAOD AK4 jets carry no B/C hadron counts (the AK8 version uses FatJet_nBHadrons / nCHadrons)
- |Jet_partonFlavour| kept as a second axis (b, c, uds, g, none) as a cross-check only: 99% of hadron-b and 95% of hadron-c jets agree (next pages)
- Same convention as the BTV SFs, so b/c-tag SFs can be applied directly once added

Official reference: CMS TWiki "Jet Flavour Identification (MC Truth)"

https://twiki.cern.ch/twiki/bin/view/CMSPublic/SWGuideBTagMCTools#Hadron_parton_based_jet_flavour

See also CMS, JINST 13 (2018) P05011 (arXiv:1712.07158), jet flavour definition

hadronFlavour vs partonFlavour

hadronFlavour (default)

- Ghost-matched B / C hadrons clustered into the jet
- b (5) if any B hadron, else c (4) if any C hadron, else 0
- BTV standard; b/c-tag SFs are measured for it

partonFlavour


- Ghost-matched partons (last before hadronization) in the jet
- Hardest heavy parton wins: b (5), c (4), else hardest light parton
- Splits light jets into uds (1–3) and gluon (21); 0 = no parton matched

Fraction of scaled yield	hadron Z→ee	parton Z→ee	hadron Z→μμ	parton Z→μμ
b	0.040	0.041	0.042	0.044
c	0.081	0.087	0.082	0.087
light (hadron) / uds + g + none (parton)	0.879	0.872	0.876	0.870
of which g / none (parton)		0.273 / 0.004		0.278 / 0.004

Summer22 DY→$\ell\ell$ MC · 20 files per sample · AK4 jets · leading AK4 jet, scaled to 7.98 fb^{-1} · 22

Hadron vs parton flavour migration


Each row shows where the jets of one hadronFlavour land in partonFlavour. 99% of hadron-b and 95% of hadron-c jets also have partonFlavour b / c (both channels). Hadron-light jets split into uds (67%) and gluon (31%); 1.2% carry a c or b parton without a matching hadron.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · rows: hadronFlavour, each sums to 100% (scaled) · small negative entries = NLO negative wei


Leading AK4 jet pT, by partonFlavour

Same as page 13, stacked by |partonFlavour| of the leading jet.



Leading AK4 jet η, by partonFlavour


Same as page 14, stacked by |partonFlavour| of the leading jet.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · by leading-jet partonFlavour; lower: fraction per bin · Z→ee left, Z→μμ right · 25

AK4 jet multiplicity, by partonFlavour


Same as page 15, stacked by |partonFlavour| of the leading jet.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · by leading-jet partonFlavour; lower: fraction per bin · Z→ee left, Z→μμ right · 26

PNet B, by partonFlavour


Same as page 17, stacked by |partonFlavour| of the leading jet.



Summer22 DY→ℓℓ MC · 20 files per sample · AK4 jets · by leading-jet partonFlavour; lower: fraction per bin · Z→ee left, Z→μμ right · 27


PNet CvL, by partonFlavour

Same as page 18, stacked by |partonFlavour| of the leading jet.



PNet CvB, by partonFlavour

Same as page 19, stacked by |partonFlavour| of the leading jet.



Leading jets with partonFlavour 0 are pileup

A jet with no GenJet (pT > 10 GeV) within ΔR < 0.4 (Jet_genJetIdx = −1) is a pileup jet. 395 of 399 (99%) leading jets with partonFlavour 0 have no GenJet match (131 Z→ee, 264 Z→μμ).

Leading-jet partonFlavour	Z→ee no GenJet / all	Z→ee %	Z→μμ no GenJet / all	Z→μμ %
b	5 / 33,829	0.01	6 / 50,225	0.01
c	14 / 63,633	0.02	26 / 95,070	0.03
uds	97 / 451,608	0.02	187 / 684,307	0.03
g	363 / 174,296	0.21	621 / 265,005	0.23
none	131 / 132	99.24	264 / 267	98.88
all	610 / 723,498	0.08	1,104 / 1,094,874	0.10

- Not every pileup jet has partonFlavour 0: most unmatched leading jets carry a gluon label (a soft ISR/UE gluon is ghost-matched, but there is no GenJet).
- Pileup barely reaches the heavy-flavour yields: 7 of 82,303 hadron-b and 36 of 151,037 hadron-c leading jets are unmatched. Scaled, 1.8% (Z→ee) and 1.7% (Z→μμ) of leading jets are unmatched. This is above the raw fraction because the unmatched jets sit in the high-weight inclusive sample, but it is close to the 1-file value, so it is not a statistical fluke.

N_jet bumps go away with more MC

20 files per sample instead of 1. Z→ee N_jet = 6: 45.5 (N_eff 0.7) → 45.5 (N_eff 0.7); Z→μμ N_jet = 6: 108.4 (N_eff 2.9) → 108.4 (N_eff 2.9), N_jet = 8: 0.0 (N_eff 0.0) → 0.0 (N_eff 0.0). Both spectra now fall smoothly up to N_jet = 7. The new Z→ee N_jet = 8 point, 33.7 (N_eff 1.1), is again a single inclusive-sample event: N_jet ≥ 6 still needs the full run.

