BTVNANOCOMMISSIONING · ZB_ZC_RATIO WORKFLOW

Z(→ℓℓ) + b/c AK4 jets in Run 3

Summer22 DY→ℓℓ MC with AK4 jets: samples, cutflow, kinematics, and a comparison with the boosted AK8 selection

11 October 2026

Campaign and setup

Goal: measure σ(Z+b) / σ(Z+c) with an AK4 jet recoiling against Z→ee/μμ (switched from the boosted AK8 jet after the group meeting). First step: tune the selection on DY→ℓℓ MC.

MC CAMPAIGN

Summer22

Run3Summer22 = 2022 eras C–D (pre-EE)

FORMAT

NanoAODv12

130X MC conditions

ENERGY (√s)

13.6 TeV

Run 3 pp collisions

NORMALIZATION

7.98 fb^{-1}

2022 C–D golden JSON; brilcalc check pending

MC only, no data yet. Processed: 1 file per sample (test run), the same files as the AK8 test. Code: BTVNanoCommissioning, branch coffea_machine, workflow Zb_Zc_ratio (AK4, commit f8748a6); the AK8 selection runs as Zb_Zc_ratio_AK8.

MC samples: DY→ℓℓ, stitched in pT(ℓℓ)

```
/DYto2L-2Jets_MLL-50[_PTLL-X]_TuneCP5_13p6TeV_amcatnloFXFX-pythia8/
  Run3Summer22NanoAODv12-130X_mcRun3_2022_realistic_v5-vN/NANOAODSIM
```

Sample	LHE pT(ℓℓ) used	σ [pb]	DAS files	DAS events	Test: events used
Inclusive (MLL-50)	< 100 GeV	6346	525	93.8 M	197,672
PTLL-100	100–200 GeV	105.0	418	56.4 M	161,921
PTLL-200	200–400 GeV	11.24	335	44.7 M	91,186
PTLL-400	400–600 GeV	0.643	424	41.6 M	2,727
PTLL-600	> 600 GeV	0.0994	185	9.50 M	939

PTLL-X samples are inclusive above X, so each is kept to its own LHE pT(ℓℓ) slice. σ(PTLL-X) = 6346 pb × fraction of inclusive events with pT(ℓℓ) > X (measured on 27.7 M inclusive events). "Test: events used" = events of 1 file inside the slice. With no boosted-jet requirement the inclusive sample (pT(ℓℓ) < 100 GeV) gives 87% (Z→ee) / 89% (Z→μμ) of the scaled yield from only ~12% of the raw selected events, so its large per-event weights limit the MC statistics.

Event selection, per channel

Z→ee / Z→μμ (unchanged)

- Trigger: HLT_Ele30_WPTight_Gsf / HLT_IsoMu24
- Leptons pT > 25 GeV, |η| < 2.4; leading pT > 35 GeV
- e: tight cut-based ID, IP cuts, EB–EE gap veto
- μ: tight ID, PF isolation < 0.15
- Opposite sign (OS)
- 71 < m$\ell\ell$ < 111 GeV
- PuppiMET < 50 GeV

Leading AK4 jet ("jet" step)

- AK4 PUPPI jets, ΔR > 0.4 from selected leptons
- Tight-lepton-veto jet ID (with NanoAODv12 jetId fix)
- pT > 30 GeV, |η| < 2.5; ≥ 1 such jet, leading one used
- No Z pT requirement

Weights and flavour

- genWeight × pileup × lepton ID/iso/reco SFs
- No trigger SF yet
- Jet flavour from hadronFlavour: b (5), c (4), light; no bb/cc split for AK4
- Tagger: PNet AK4 B, CvL, CvB

Cutflow, Z→ee: raw MC events per step

Sample	all	trigger	leptons	OS	Zmass	MET	jet	jet / MET
Inclusive	197,672	32,660	9,802	9,745	9,396	8,872	2,347	26.5%
PTLL-100	161,921	46,448	13,156	13,037	12,032	10,642	10,214	96.0%
PTLL-200	91,186	31,202	11,594	11,431	10,398	8,584	8,466	98.6%
PTLL-400	2,727	1,013	458	447	396	287	286	99.7%
PTLL-600	939	229	101	100	82	54	54	100.0%

21,367 raw MC events pass the full Z→ee selection; the most (48%) come from PTLL-100. The AK4 jet step keeps 26% of inclusive (low-pT) Z events and ≥ 96% at pT($\ell\ell$) > 100 GeV.

Summer22 DY→$\ell\ell$ MC · 1 file per sample · AK4 jets · "all" = events in the sample's LHE pT($\ell\ell$) slice · 5

Cutflow, Z→μμ: raw MC events per step

Sample	all	trigger	leptons	OS	Zmass	MET	jet	jet / MET
Inclusive	197,672	42,062	17,685	17,685	17,054	16,139	4,159	25.8%
PTLL-100	161,921	48,592	20,606	20,604	19,043	16,788	16,142	96.2%
PTLL-200	91,186	31,717	16,868	16,865	15,258	12,453	12,298	98.8%
PTLL-400	2,727	1,048	659	659	592	407	405	99.5%
PTLL-600	939	319	199	199	178	102	102	100.0%

33,106 raw MC events pass the full Z→μμ selection; the most (49%) come from PTLL-100. The AK4 jet step keeps 26% of inclusive (low-pT) Z events and ≥ 96% at pT($\ell\ell$) > 100 GeV.

Summer22 DY→$\ell\ell$ MC · 1 file per sample · AK4 jets · "all" = events in the sample's LHE pT($\ell\ell$) slice · 6

Yields by AK4 jet flavour (7.98 fb^{-1})

Channel	light	c	b	total
Z→ee	343,532.5	42,631.0	16,844.4	403,007.9
fraction	0.852	0.106	0.042	
Z→μμ	683,956.5	65,108.4	35,637.3	784,702.2
fraction	0.872	0.083	0.045	

Still 1 file per sample

No negative yields now: 21,367 (Z→ee) and 33,106 (Z→μμ) raw events pass, vs ~1.4k / 2.1k with AK8. Single negative-weight amcatnlo events still leave jumps and empty bins in the tails of the plots. No b-tag requirement applied yet.


AK8 vs AK4: same files, same Z selection

	AK8 Z→ee	AK4 Z→ee	AK8 Z→μμ	AK4 Z→μμ
Raw events selected	1,440	21,367	2,121	33,106
Scaled yield	2,259.7	403,007.9	2,784.7	784,702.2
b fraction (b + bb)	0.043	0.042	−0.045	0.045
c fraction (c + cc)	0.118	0.106	0.244	0.083
light fraction	0.840	0.852	0.800	0.872
Jet requirement	pT > 200, mSD > 40	pT > 30	pT > 200, mSD > 40	pT > 30

AK8 flavour is from B/C hadron counts inside the jet (bb/b/cc/c); AK4 flavour is hadronFlavour. The AK4 selection keeps ~15× more events because it does not require a boosted Z. AK8 heavy-flavour fractions are unreliable at this statistics (negative Z→μμ b yield).

Dilepton mass


m(ℓℓ) after the full selection (71–111 GeV window): peaks at the Z mass in both channels.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 9

Dilepton pT


pT(ℓℓ) of the Z candidate after the full selection. No Z pT cut: dominated by low pT, unlike the boosted AK8 selection.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 10

Leading-lepton pT


Leading lepton of the Z candidate, after the full selection (pT > 35 GeV required).



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 11

Subleading-lepton pT


Subleading lepton of the Z candidate, after the full selection (pT > 25 GeV required).



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 12

Leading AK4 jet pT


Leading lepton-cleaned AK4 jet with tight-lepton-veto ID (pT > 30 GeV required).



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 13

Leading AK4 jet η


Pseudorapidity of the selected AK4 jet (|η| < 2.5 required).



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 14

AK4 jet multiplicity


Number of selected AK4 jets (pT > 30 GeV, |η| < 2.5, ID, lepton-cleaned) in events passing the full selection.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 15

LHE pT(ℓℓ)


Generator-level pT(ℓℓ) after the full selection: checks the sample stitching (slices at 100/200/400/600 GeV).



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 16

Leading AK4 jet: PNet B


ParticleNet b-tag score of the selected jet; b jets pile up near 1, light jets near 0.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 17

Leading AK4 jet: PNet CvL


ParticleNet charm-vs-light score of the selected jet.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 18

Leading AK4 jet: PNet CvB


ParticleNet charm-vs-b score of the selected jet.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · stacked by leading AK4 jet flavour · Z→ee left, Z→μμ right · 19

PNet charm-tagging plane: CvL vs CvB

Selected AK4 jets per flavour, both channels summed, normalized to unit area. b jets sit at high CvL and low CvB, light jets at low CvL and high CvB. c jets are at high CvB, but a large part of them overlaps the light-jet corner (low CvL), with the rest at high CvL.



Summer22 DY→ℓℓ MC · 1 file per sample · AK4 jets · Z→ee + Z→μμ · 20