

③ **MODE: (M)** most frequently occurring value.

UNGROUPED

Observation having maximum frequency

GROUPED

$$M = l + \left(\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

- ①: lower limit of the modal class
- ②: class size
- ③: Frequency of the modal class
- ④: frequency of the class succeeding modal class
- ⑤: class preceding modal class
- ⑥: class preceding modal class

EMPIRICAL RELATION between Mean, Median and Mode.

$$\text{MODE} = 3 \text{ MEDIAN} - 2 \text{ MEAN}$$

NOTE: Measures of Central tendency give only representative value so, is INSUFFICIENT in many scenarios. To overcome, this we learn MEASURES OF DISPERSION.

Prepared by Neha Agrawal Mathematically Inclined

STATISTICS (RECALL)

Neha Agrawal Mathematically Inclined!

KDS xpo gygi!

(I) **QUICK RECALL:**

MEASURES OF CENTRAL TENDENCY:

① **MEAN / ARITHMETIC MEAN / AVERAGE (\bar{X})**

UNGROUPED

For observations x_1, x_2, \dots, x_n

$$\bar{X} = \frac{\sum_{i=1}^n x_i}{n}$$

GROUPED

For observations x_1, x_2, \dots, x_n with frequencies f_1, f_2, \dots, f_n

$$\bar{X} = \frac{\sum_{i=1}^n f_i x_i}{\sum_{i=1}^n f_i}$$

ASSUMED MEAN (SHORT CUT) METHOD

$$\bar{X} = A + \frac{\sum_{i=1}^n f_i d_i}{\sum_{i=1}^n f_i}$$

$$d_i = x_i - A$$

Assumed Mean

(M)

② **MEDIAN:** Middle most or Central value when data is arranged in **ASCENDING** order.

UNGROUPED

n is odd $M = \left(\frac{n+1}{2} \right)^{\text{th}} \text{ term}$

n is even $\left(\frac{\frac{n}{2} + \frac{n+2}{2}}{2} \right)^{\text{th}} \text{ term}$

GROUPED

$$M = l + \frac{\frac{n}{2} - C.f.}{f} \times h$$

- ①: lower lim of median class
- ②: Frequency
- ③: Cumulative freq. of preceding class

(II) MEASURES OF DISPERSION

1. RANGE = Max value - Min value of the Distribution

2. MEAN DEVIATION:

DATA ABOUT \bar{X} ABOUT MEDIAN (M)

UNGROUPED	DATA ABOUT \bar{X}	ABOUT MEDIAN (M)
	$\frac{\sum x_i - \bar{x} }{n}$	$\frac{\sum x_i - M }{n}$
GROUPED	$\frac{\sum f_i x_i - \bar{x} }{N}$ $N = \sum_{i=1}^n f_i$	$\frac{\sum f_i x_i - M }{N}$ $N = \sum_{i=1}^n f_i$

* Shortcut method for calculating \bar{X} or M is available on Page 1

3. VARIANCE and STANDARD DEVIATION

Var σ^2 S.D. σ

$$S.D. = +\sqrt{\text{Var}}$$

mean of the squares of the deviations from mean is called variance.

Prepared by

Neha Agrawal

Mathematically

Inclined

STATISTICS

Neha Agrawal

Mathematically

Inclined

KDS too gya! \heartsuit

DATA TYPE	VARIANCE	STANDARD DEVIATION
UNGROUPED	$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$	$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}$
GROUPED	$\sigma^2 = \frac{1}{N} \sum_{i=1}^n f_i (x_i - \bar{x})^2$ $N = \sum_{i=1}^n f_i$	$\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^n f_i (x_i - \bar{x})^2}$

ALTERNATE FORMULAE:

UNGROUPED $\sigma^2 = \frac{1}{n} \left(\sum x_i^2 \right) - \bar{x}^2$

GROUPED $\sigma^2 = \frac{1}{n} \sum f_i x_i^2 - \left(\frac{\sum f_i x_i}{N} \right)^2$

(III) MAGIC TABLE

Quantity	Change of Origin	Change of Scale
Mean	$+/- a$	$a(\text{old})$ or old/a
Median	$+/- a$	$a(\text{old})$ or old/a
Mode	$+/- a$	$a(\text{old})$ or old/a
Var	same	$a^2(\text{old})$
S.D.	same	$ a (\text{old})$

(IV) COEFFICIENT OF VARIATION

$CV = \frac{\sigma}{\bar{x}} \times 100$ used for comparing DISPERION of 2 series