

Workout Injury Risk Report

2026-09-29 10:09

Health Summary

BMI 19.8 (Normal) - Goal: Bulk

Great work finishing your session today! At a BMI of 19.8, you are in a healthy weight range that provides a fantastic, lean baseline for your bulking goal. To effectively build muscle on a vegetarian diet, focus on staying in a modest calorie surplus while getting plenty of protein from complete sources like eggs, as well as dairy, legumes, and tofu. Combining this nutrition strategy with progressive resistance training and solid rest will give your body everything it needs to pack on quality size.

Workout Summary


Duration: 02:48

Exercises Completed

- Lateral Raises: 9 reps
- Bicep Curls: 7 reps
- Shoulder Press: 7 reps
- Hammer Curls: 6 reps

Mishaps & Injury Risk

Goal-Post Press (Elbows Flaring Early)



Evidence: Based on an official training guideline (US Army FM 7-22)

During your overhead press set, your elbows swung out to the sides before your arms had straightened. This meant your shoulders held the weight out wide instead of pushing it straight up, which partly turns the press into a lateral raise under a heavy load. Standard guidance in FM 7-22 specifies that the overhead press starts with the shoulders directly below the hands and pushes straight up.

How to prevent it


Fix it now: Keep your forearms vertical, with your wrists stacked over your elbows, and push the weight in a straight line up over your shoulders.

Corrective drill: Shoulder Stability Drill (FM 7-22, p. 6-25): I, T, Y, L and W raises lying face down, 5 slow reps each, bodyweight only. FM 7-22 uses it to build shoulder strength and stability, and in rehab after shoulder injuries.

Progress safely: Lighten the load until the bar or dumbbells travel straight up on every rep. Add weight in small steps and only after a full session of clean reps. Injury risk rises with the size of the jump from what your body is used to (HHS, p. 90).

Sources: US Army FM 7-22 (2012), p. 6-47, overhead press starting position; Kolber et al., J Strength Cond Res (2010), technique as a shoulder-injury risk factor

Elbow Drift (Curl)



Evidence: Technique standard - no specific injury study; general shoulder-injury evidence applies

During your curl, your elbow drifted forward and away from your side, so the front of your shoulder started helping to lift the weight. Consequently, your biceps does less of the work while your shoulder picks up load it does not need to carry. No study has linked this specific fault to injury, so you can treat this mainly as technique coaching. Nevertheless, staying mindful of this is helpful because the shoulder is the most commonly injured area in weight training.

How to prevent it


Fix it now: Pin your elbows to your sides and move only your forearms. Use a weight you can lift without swinging.

Corrective drill: Curl at a slow, controlled pace, as in FM 7-22's machine biceps curl (p. 6-62), with no arching back to move the weight.

Progress safely: If you have to swing your body or let your elbows drift to finish a rep, the weight is too heavy. Drop it until every rep is clean.

Sources: US Army FM 7-22 (2012), p. 6-62, biceps curl; Kolber et al., J Strength Cond Res (2010), general shoulder-injury evidence

Elbow Drift (Curl)



Evidence: Technique standard - no specific injury study; general shoulder-injury evidence applies

During your set, your elbow drifted forward and away from your side as you curled, causing the front of your shoulder to assist in lifting the weight. This results in your biceps doing less work while your shoulder carries unnecessary load. Because no study has linked this specific fault to injury, this is primarily a technique fault to treat as technique coaching. It remains good to track, however, since the shoulder is the most commonly injured area in weight training.

How to prevent it


Fix it now: Pin your elbows to your sides and move only your forearms. Use a weight you can lift without swinging.

Corrective drill: Curl at a slow, controlled pace, as in FM 7-22's machine biceps curl (p. 6-62), with no arching back to move the weight.

Progress safely: If you have to swing your body or let your elbows drift to finish a rep, the weight is too heavy. Drop it until every rep is clean.

Sources: US Army FM 7-22 (2012), p. 6-62, biceps curl; Kolber et al., *J Strength Cond Res* (2010), general shoulder-injury evidence

Arms Raised Above Shoulder Height



Evidence: Based on an official training guideline (US Army FM 7-22)

During your set, you raised your arms above shoulder height during the lateral raise. Raising your arms sideways above shoulder level with weight is a position commonly linked to shoulder impingement. Guidance in FM 7-22 is explicit on this exercise, stating that you must not raise your arms above parallel to the ground. Keeping within this range is worth your focus, as the shoulder accounts for up to 36% of weight-training injuries.

How to prevent it

Fix it now: Stop when your arms are parallel to the floor, keep your elbows slightly bent, and lead with your elbows, not your hands.

Corrective drill: Shoulder Stability Drill (FM 7-22, p. 6-25): I, T, Y, L and W raises lying face down, 5 slow reps each, bodyweight only. FM 7-22 uses it to build shoulder strength and stability, and in rehab after shoulder injuries.

Progress safely: Use light dumbbells. If you need to swing or shrug to reach shoulder height, the weight is too heavy.

Sources: US Army FM 7-22 (2012), p. 6-54, lateral raise precautions; Kolber et al., *J Strength Cond Res* (2010), prevalence of shoulder injuries in resistance training

General Injury Prevention

- Increase load gradually. Injury risk to bones, muscles and joints is tied to the gap between what you usually do and what you do next. Make small increases and give your body time to adapt. (HHS Physical Activity Guidelines (2018), p. 90)
- Warm up before strength work by doing your exercises with lighter weight first. (HHS Physical Activity Guidelines (2018), p. 91)
- More is not better. Soreness and fatigue that never fully clear, together with falling performance, are early signs of overtraining. Rest, recovery and good nutrition bring more improvement than training harder. (US Army FM 7-22 (2012), p. 5-3)
- If you have been injured before, that body part is at higher risk of injury again, so progress it more slowly than the rest. (HHS Physical Activity Guidelines (2018), p. 89 and 91)

References

- [1] Headquarters, Department of the Army. FM 7-22: Army Physical Readiness Training. Washington, DC; October 2012 (incl. Change 1, May 2013). (Government guideline)
- [2] Kolber MJ, Beekhuizen KS, Cheng MS, Hellman MA. Shoulder injuries attributed to resistance training: a brief review. J Strength Cond Res. 2010;24(6):1696-1704. doi:10.1519/JSC.0b013e3181dc4330 (Peer-reviewed review)
- [3] U.S. Department of Health and Human Services. Physical Activity Guidelines for Americans, 2nd edition. Washington, DC: HHS; 2018. Chapter 7, Active and Safe. (Government guideline)

This report gives general exercise-technique guidance based on the sources above. It is not a medical diagnosis. If you have pain, especially sharp or lasting pain, stop the exercise and see a doctor or physiotherapist.