

Your Workout Form Report

2026-09-29 11:30

Health Summary

BMI 19.8 (Normal) - Goal: Bulk

Great job finishing your workout today! At a healthy BMI of 19.8, you have an excellent baseline to safely begin bulking and building lean muscle mass. To support your strength goals on a vegetarian diet, focus on maintaining a consistent caloric surplus and leveraging high-quality protein sources like eggs, dairy, and legumes to fuel your recovery and muscle growth.

Workout Summary

Duration: 03:21

Exercises Completed

- Bicep Curls: 12 reps
- Pullups: 1 reps
- Shoulder Press: 8 reps
- Lateral Raises: 7 reps
- Tricep Dips: 4 reps
- Leg Raises: 2 reps

At a Glance

We spotted 3 form issues in this workout. Here is the quick version; the details are further down.

- **Elbows Flaring Out Too Early (Shoulder Press):** Keep your forearms upright, with your wrists directly above your elbows, and push the weight in a straight line up over your shoulders.
- **Snapping Knees Straight (Leg Extension):** Straighten your legs slowly until they are straight but not locked, pause for a moment, then lower them slowly.
- **Elbow Moving Forward (Curl):** Keep your elbows pinned to your sides and move only your forearms.

Form Issues in Detail

AI summary unavailable - showing the standard explanations. The advice and sources below are unchanged.

Elbows Flaring Out Too Early (Shoulder Press)

Exercise: Shoulder Press | How sure we are: Based on an official training manual (US Army)

What happened

Your elbows swung out to the sides before your arms had straightened, so your shoulders held the weight out wide instead of pushing it straight up. Part of the press turns into a side raise with a heavy weight, which is harder on the shoulders. The standard way is to start with your hands above your shoulders and push straight up.

How to fix it

Next time: Keep your forearms upright, with your wrists directly above your elbows, and push the weight in a straight line up over your shoulders.

Exercise that helps: Shoulder raises lying face down, no weights. Lie on your stomach and lift both arms 8-15 cm (3-6 inches) off the floor, first straight above your head (like the letter I), then straight out to the sides (like a T). Do 5 slow reps of each. This strengthens the small muscles that keep your shoulder joint steady.

Going forward: Use less weight until it travels straight up on every rep. Add weight in small steps, and only after a whole workout of good reps. Injuries are most likely when you jump far beyond what your body is used to.

Sources: US Army FM 7-22 (2012), p. 6-47, shoulder press; Kolber et al., J Strength Cond Res (2010), technique and shoulder injuries; US Army FM 7-22 (2012), p. 6-25 to 6-27, shoulder stability drill; HHS Physical Activity Guidelines (2018), p. 90, increase activity gradually

Snapping Knees Straight (Leg Extension)

Exercise: Leg Extension | How sure we are: Strong - backed by research studies and an official training manual

What happened

You kicked your legs up hard and snapped your knees straight at the top, so momentum finished the rep instead of your thigh muscles, and your knee joint was jolted into its end position under weight. Official guidance straightens the legs fully but without locking them, and lowers them slowly. Research also shows the pressure behind the kneecap in this exercise is highest in the middle of the movement, so moving slowly through the whole range matters, not just at the top.

How to fix it

Next time: Straighten your legs slowly until they are straight but not locked, pause for a moment, then lower them slowly.

Exercise that helps: Practise with a light weight: lift slowly, stop just before your knees lock, hold for a moment, and lower slowly, until that pace feels natural.

Going forward: Choose a weight you can lower more slowly than you lift. Add weight in small steps, and only after a whole workout of good reps. Injuries are most likely when you jump far beyond what your body is used to.

Sources: US Army FM 7-22 (2012), p. 6-32, straighten legs without locking; Escamilla et al., Med Sci Sports Exerc (1998), knee pressure during leg extensions; HHS Physical Activity Guidelines (2018), p. 90, increase activity gradually

Elbow Moving Forward (Curl)

Exercise: Bicep Curls | How sure we are: Technique tip - good form, but no study links this exact mistake to injury

What happened

Your elbow moved forward and away from your side while you curled, so your shoulder started helping to lift the weight. This mainly makes the exercise less effective for your arm, and puts some load on your shoulder that it doesn't need to carry. No study links this exact mistake to injury, so treat it as a technique tip.

How to fix it

Next time: Keep your elbows pinned to your sides and move only your forearms. Use a weight you can lift without swinging.

Exercise that helps: Curl slowly and under control, without leaning back to help lift the weight.

Going forward: If you have to swing your body or move your elbows to finish a rep, the weight is too heavy. Lower it until every rep is clean.

Sources: US Army FM 7-22 (2012), p. 6-62, biceps curl; Kolber et al., J Strength Cond Res (2010), shoulder injuries in weight training

Tips to Stay Injury-Free

- Increase weight gradually. Injuries are most likely when you jump far beyond what your body is used to, so make small increases and give your body time to adapt.
- Warm up before lifting by doing each exercise with a lighter weight first.
- More is not always better. If soreness and tiredness never fully go away and your performance keeps dropping, you may be overtraining. Rest, sleep and good food will help you improve more than training harder.
- If you have hurt a body part before, it is more likely to get hurt again, so build that area up more slowly.

Sources: HHS Physical Activity Guidelines (2018), p. 90; HHS Physical Activity Guidelines (2018), p. 91; US Army FM 7-22 (2012), p. 5-3; HHS Physical Activity Guidelines (2018), p. 89 and 91

References

- [1] Headquarters, Department of the Army. FM 7-22: Army Physical Readiness Training. Washington, DC; October 2012 (incl. Change 1, May 2013). (Government guideline)
- [2] Kolber MJ, Beekhuizen KS, Cheng MS, Hellman MA. Shoulder injuries attributed to resistance training: a brief review. J Strength Cond Res. 2010;24(6):1696-1704. doi:10.1519/JSC.0b013e3181dc4330 (Peer-reviewed review)
- [3] U.S. Department of Health and Human Services. Physical Activity Guidelines for Americans, 2nd edition. Washington, DC: HHS; 2018. Chapter 7, Active and Safe. (Government guideline)
- [4] Escamilla RF, Fleisig GS, Zheng N, Barrentine SW, Wilk KE, Andrews JR. Biomechanics of the knee during closed kinetic chain and open kinetic chain exercises. Med Sci Sports Exerc. 1998;30(4):556-569. (Peer-reviewed lab study)

This report gives general exercise-technique guidance based on the sources above. It is not a medical diagnosis. If you have pain, especially sharp or lasting pain, stop the exercise and see a doctor or physiotherapist.