

FUNCTION

1. CARTESIAN PRODUCT OF TWO SETS :

Given two non-empty sets A and B. The cartesian product $A \times B$ is the set of all ordered pairs of the form (a, b) where the first entry comes from set A & second comes from set B.

$$A \times B = \{(a, b) \mid a \in A, b \in B\}$$

e.g. $A = \{1, 2, 3\}$ $B = \{p, q\}$

$$A \times B = \{(1, p), (1, q), (2, p), (2, q), (3, p), (3, q)\}$$

Note :

- If either A or B is the null set, then $A \times B$ will also be empty set, i.e. $A \times B = \phi$
- If $n(A) = p, n(B) = q$, then $n(A \times B) = pq$, where $n(X)$ denotes the number of elements in set X.
- A **Relation** R from set A to B is any subset of $A \times B$. If $A R B$ & $(a, b) \in R$ then b is image of a under R and a is preimage of b under R.

Note : If $n(A) = m, n(B) = n$, then number of relations defined from set A to B are 2^{mn} .

2. FUNCTION :

A relation R from set A to set B is called a function if each element of A is uniquely associated with some element of B. It is denoted by the symbol :

$$f : A \rightarrow B \text{ or } A \xrightarrow{f} B$$

which reads 'f' is a function from A to B 'or' f maps A to B,

If an element $a \in A$ is associated with an element $b \in B$, then b is called 'the f image of a' or 'image of a under f' or 'the value of the function f at a'. Also a is called the 'pre-image of b' or 'argument of b under the function f'. We write it as

$$b = f(a) \text{ or } f : a \rightarrow b \text{ or } f : (a, b)$$

Thus a function 'f' from set A to set B is subset of $A \times B$ in which each a belonging to A appears in one and only one ordered pair belonging to f.

Representation of Function :

(a) **Ordered pair :** Every function from $A \rightarrow B$ satisfies the following conditions :

- $f \subset A \times B$
- $\forall a \in A$ there exist $b \in B$ and
- $(a, b) \in f$ & $(a, c) \in f \Rightarrow b = c$

(b) **Formula based (uniformly/nonuniformly) :**


(i) $f : \mathbb{R} \rightarrow \mathbb{R}, y = f(x) = 4x, f(x) = x^2$ (uniformly defined)

(ii) $f(x) = \begin{cases} x+1 & -1 \leq x < 4 \\ -x & 4 \leq x < 7 \end{cases}$ (non-uniformly defined)


(iii) $f(x) = \begin{cases} x^2 & x \geq 0 \\ -x-1 & x < 0 \end{cases}$ (non-uniformly defined)

(c) **Graphical representation :**

If a vertical line cuts a given graph at more than one point then it can not be the graph of a function.



Graph (1)



Graph (2)

Graph(1) represent a function but graph(2) does not represent a function.

Every function is a relation but every relation is not necessarily a function.

3. DOMAIN, CO-DOMAIN & RANGE OF A FUNCTION :

Let $f: A \rightarrow B$, then the set A is known as the domain of f & the set B is known as co-domain of f . The set of f images of all the elements of A is known as the range of f .

Domain of $f = \{a \mid a \in A, (a, f(a)) \in f\}$

Range of $f = \{f(a) \mid a \in A, f(a) \in B\}$

- If only the rule of function is given then the domain of the function is the set of those real numbers, where function is defined.
- For a continuous function, the interval from minimum to maximum value of a function gives the range
- It should be noted that range is a subset of co-domain.

Note :

- The complete set of all positive real numbers is denoted by \mathbb{R}^+ or $\mathbb{R}_{>0}$
- The complete set of all negative real numbers is denoted by \mathbb{R}^- or $\mathbb{R}_{<0}$
- The complete set of all real numbers other than zero is denoted by \mathbb{R}^* or $\mathbb{R}_{\neq 0}$
- The complete set of all integers is denoted by \mathbb{Z} .

Illustration 1 : Find the domain of following functions :

$$(i) y = \sqrt{5-2x} \quad (ii) y = \frac{1}{\sqrt{x-|x|}}$$

Solution : (i) $5-2x \geq 0 \Rightarrow x \leq \frac{5}{2} \therefore$ Domain is $(-\infty, 5/2]$

(ii) $x-|x| > 0 \Rightarrow |x| < x \Rightarrow x$ cannot take any real values \therefore Domain is ϕ

Illustration 2 : Find the range of following functions :

$$(i) f(x) = \log_{\sqrt{2}}((x-1)^2 + 4) \quad (ii) f(x) = 3 - \cos x$$

Solution : (i) $f(x) = \log_{\sqrt{2}}((x-1)^2 + 4)$
 $4 \leq (x-1)^2 + 4 < \infty$
 $\Rightarrow \log_{\sqrt{2}} 4 \leq \log_{\sqrt{2}}((x-1)^2 + 4) < \infty$
 $\Rightarrow 4 \leq \log_{\sqrt{2}}((x-1)^2 + 4) < \infty$
 \therefore Range of $f(x) = [4, \infty)$
 (ii) $f(x) = 3 - \cos x$
 $-1 \leq \cos x \leq 1$
 $2 \leq 3 - \cos x \leq 4$
 \therefore Range of $f(x) = [2, 4]$

Do yourself - 1 :

(i) Find the domain of following functions :

$$(a) y = 1 - \log_{10} x \quad (b) y = \frac{1}{\sqrt{x^2 - 4x}}$$

(ii) Find the range of the following function :

$$(a) \log_4 \left| x + \frac{1}{x} \right| \quad (b) f(x) = \sin(3x^2 + 1)$$

$$(c) f(x) = 2 \sin \left(2x + \frac{\pi}{4} \right) \quad (d) f(x) = \cos \left(2x + \frac{\pi}{4} \right)$$

4. IMPORTANT TYPES OF FUNCTIONS :

(a) Polynomial Function :

If a function f is defined by $f(x) = a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_{n-1} x + a_n$ where n is a non negative integer and $a_0, a_1, a_2, \dots, a_n$ are real numbers and $a_0 \neq 0$, then f is called a polynomial function of degree n . If n is odd, then polynomial is of odd degree, if n is even, then polynomial is of even degree.

Note :

- Range of odd degree polynomial is always \mathbb{R} .
- Range of even degree polynomial is never \mathbb{R} .
- A polynomial of degree one with no constant term is called an odd linear function.
i.e. $f(x) = ax$, $a \neq 0$
- $f(x) = ax + b$, $a \neq 0$ is a linear polynomial
- $f(x) = c$ is non linear polynomial (its degree is zero)
- $f(x) = 0$ is a polynomial but its degree is **not** defined
- There are four polynomial functions, satisfying the relation ;
 $f(x) \cdot f(1/x) = f(x) + f(1/x)$. They are : (a) $f(x) = 0$ (b) $f(x) = 2$
(c) $f(x) = x^n + 1$ & (d) $f(x) = 1 - x^n$, where n is a positive integer.

(b) Algebraic Function :

y is an algebraic function of x , if it is a function that satisfies an algebraic equation of the form $P_0(x) y^n + P_1(x) y^{n-1} + \dots + P_{n-1}(x) y + P_n(x) = 0$. Where n is a positive integer and $P_0(x), P_1(x), \dots$ are Polynomials in x . e.g. $x^3 + y^3 - 3xy = 0$.

In other words a function ' f ' is called an algebraic function if it can be constructed using finite number of algebraic operations (such as addition, subtraction, multiplication, division, and taking radicals) within polynomials.


Example : $f(x) = \sqrt{x^2 + 1}$; $g(x) = \frac{x^4 - 16x^2}{x + \sqrt{x}} + (x - 2) \sqrt[3]{x + 1}$

Note :

- All polynomial functions are Algebraic but not the converse.
- A function that is not algebraic is called **Transcendental Function**.

Basic algebraic function :


(i) $y = x^2$



Domain : \mathbb{R}

Range : $\mathbb{R}^+ \cup \{0\}$ or $[0, \infty)$


(ii) $y = \frac{1}{x}$



Domain : $\mathbb{R} - \{0\}$ or \mathbb{R}_0

Range : $\mathbb{R} - \{0\}$


(iii) $y = \frac{1}{x^2}$



Domain : \mathbb{R}_0

Range : \mathbb{R}^+ or $(0, \infty)$

(iv) $y = x^3$



Domain : \mathbb{R}

Range : \mathbb{R}

(c) Rational function :

A rational function is a function of the form $y = f(x) = \frac{g(x)}{h(x)}$, where $g(x)$ & $h(x)$ are polynomials

&

$h(x) \neq 0$, **Domain** : $\mathbb{R} - \{x \mid h(x)=0\}$

Any rational function is automatically an algebraic function.


(d) Trigonometric functions :

(i) Sine function

$$f(x) = \sin x$$

Domain : \mathbb{R}

Range : $[-1, 1]$, period 2π




(ii) Cosine function

$$f(x) = \cos x$$

Domain : \mathbb{R}

Range : $[-1, 1]$, period 2π




(iii) Tangent function

$$f(x) = \tan x$$

Domain : $\mathbb{R} - \left\{x \mid x = \frac{(2n+1)\pi}{2}, n \in \mathbb{I}\right\}$

Range : \mathbb{R} , period π




(iv) Cosecant function

$$f(x) = \operatorname{cosec} x$$

Domain : $\mathbb{R} - \{x \mid x = n\pi, n \in \mathbb{I}\}$

Range : $\mathbb{R} - (-1, 1)$, period 2π




(v) Secant function

$$f(x) = \sec x$$

$$\text{Domain : } \mathbb{R} - \{x | x = (2n + 1) \pi/2 : n \in \mathbb{I}\}$$

$$\text{Range : } \mathbb{R} - (-1, 1), \text{ period } 2\pi$$




(vi) Cotangent function

$$f(x) = \cot x$$

$$\text{Domain : } \mathbb{R} - \{x | x = n\pi, n \in \mathbb{I}\}$$

$$\text{Range : } \mathbb{R}, \text{ period } \pi$$



(e) Exponential and Logarithmic Function :

A function $f(x) = a^x$ ($a > 0$), $a \neq 1$, $x \in \mathbb{R}$ is called an exponential function. The inverse of the exponential function is called the logarithmic function, i.e. $g(x) = \log_a x$.




Note that $f(x)$ & $g(x)$ are inverse of each other & their graphs are as shown. (If functions are mirror image of each other about the line $y = x$)




Domain of a^x is \mathbb{R}

Range \mathbb{R}^+

Domain of $\log_a x$ is \mathbb{R}^+

Range \mathbb{R}





Note-1 : $f(x) = a^{1/x}$, $a > 0$ **Domain :** $\mathbb{R} - \{0\}$ **Range :** $\mathbb{R}^+ - \{1\}$

Note-2 : $f(x) = \log_x a = \frac{1}{\log_a x}$ **Domain :** $\mathbb{R}^+ - \{1\}$ **Range :** $\mathbb{R} - \{0\}$
 $(a > 0) (a \neq 1)$


(f) Absolute Value Function :

A function $y = f(x) = |x|$ is called the absolute value function or Modulus function. It is defined as :

$$y = |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

For $f(x) = |x|$, domain is \mathbb{R} and range is $[0, \infty)$

For $f(x) = \frac{1}{|x|}$, domain is $\mathbb{R} - \{0\}$ and range is \mathbb{R}^+ .



(g) Signum Function :

A function $y = f(x) = \text{Sgn}(x)$ is defined as follows :


$$y = f(x) = \begin{cases} 1 & \text{for } x > 0 \\ 0 & \text{for } x = 0 \\ -1 & \text{for } x < 0 \end{cases}$$

It is also written as $\text{Sgn } x = |x|/x$; $x \neq 0$;
 $= 0$; $x = 0$

Note : $f(x) = (\text{sgn}(x))x \Rightarrow f(x) = |x|$

Domain : \mathbb{R}


Range : $\{-1, 0, 1\}$



(h) **Greatest integer or step up function :**

The function $y = f(x) = [x]$ is called the greatest integer function where $[x]$ denotes the greatest integer less than or equal to x . Note that for :

x	$[x]$
$[-2, -1)$	-2
$[-1, 0)$	-1
$[0, 1)$	0
$[1, 2)$	1



Domain : \mathbb{R}

Range : \mathbb{I}

Properties of greatest integer function :

- (i) $[x] \leq x < [x] + 1$ and $x - 1 < [x] \leq x$, $0 \leq x - [x] < 1$
- (ii) $[x + m] = [x] + m$, if m is an integer.

$$(iii) [x] + [-x] = \begin{cases} 0, & x \in \mathbb{I} \\ -1, & x \notin \mathbb{I} \end{cases}$$

Illustration 3 : If $y = 2[x] + 3$ & $y = 3[x - 2] + 5$, then find $[x + y]$ where $[.]$ denotes greatest integer function.

Solution : $y = 3[x - 2] + 5 = 3[x] - 1$
 so $3[x] - 1 = 2[x] + 3$
 $[x] = 4 \Rightarrow 4 \leq x < 5$
 then $y = 11$
 so $x + y$ will lie in the interval $[15, 16)$
 so $[x + y] = 15$

Illustration 4 : Find the value of $\left[\frac{1}{2}\right] + \left[\frac{1}{2} + \frac{1}{1000}\right] + \dots + \left[\frac{1}{2} + \frac{2946}{1000}\right]$ where $[.]$ denotes greatest integer function ?


Solution : $\left[\frac{1}{2}\right] + \left[\frac{1}{2} + \frac{1}{1000}\right] + \dots + \left[\frac{1}{2} + \frac{499}{1000}\right] + \left[\frac{1}{2} + \frac{500}{1000}\right] + \dots + \left[\frac{1}{2} + \frac{1499}{1000}\right] + \left[\frac{1}{2} + \frac{1500}{1000}\right] + \dots$
 $+ \left[\frac{1}{2} + \frac{2499}{1000}\right] + \left[\frac{1}{2} + \frac{2500}{1000}\right] + \dots + \left[\frac{1}{2} + \frac{2946}{1000}\right]$
 $= 0 + 1 \times 1000 + 2 \times 1000 + 3 \times 447 = 3000 + 1341 = 4341$

Ans.

(i) Fractional part function :

It is defined as : $g(x) = \{x\} = x - [x]$

e.g. the fractional part of the number 2.1 is $2.1 - 2 = 0.1$ and the fractional part of -3.7 is 0.3 . The period of this function is 1 and graph of this function is as shown.



x	{x}
$[-2, -1)$	$x+2$
$[-1, 0)$	$x+1$
$[0, 1)$	x
$[1, 2)$	$x-1$

Domain : \mathbb{R}

Range : $[0, 1)$

Properties of fractional part function :

$$(i) \quad 0 \leq \{x\} < 1 \quad (ii) \quad \{[x]\} = [\{x\}] = 0 \quad (iii) \quad \{\{x\}\} = \{x\}$$

$$(iv) \quad \{x+m\} = \{x\}, \quad m \in \mathbb{I} \quad (v) \quad \{x\} + \{-x\} = \begin{cases} 1, & x \notin \mathbb{I} \\ 0, & x \in \mathbb{I} \end{cases}$$

Illustration 5 : Solve the equation $|2x - 1| = 3[x] + 2\{x\}$ where $[.]$ denotes greatest integer and $\{.\}$ denotes fractional part function.

Solution : We are given that, $|2x - 1| = 3[x] + 2\{x\}$

Let, $2x - 1 \leq 0$ i.e. $x \leq \frac{1}{2}$. The given equation yields.

$$1 - 2x = 3[x] + 2\{x\}$$

$$\Rightarrow 1 - 2[x] - 2\{x\} = 3[x] + 2\{x\} \Rightarrow 1 - 5[x] = 4\{x\} \Rightarrow \{x\} = \frac{1 - 5[x]}{4}$$

$$\Rightarrow 0 \leq \frac{1 - 5[x]}{4} < 1 \Rightarrow 0 \leq 1 - 5[x] < 4 \Rightarrow -\frac{3}{5} < [x] \leq \frac{1}{5}$$

Now, $[x] = 0$ as zero is the only integer lying between $-\frac{3}{5}$ and $\frac{1}{5}$

$$\Rightarrow \{x\} = \frac{1}{4} \Rightarrow x = \frac{1}{4} \text{ which is less than } \frac{1}{2}, \text{ Hence } \frac{1}{4} \text{ is one solution.}$$

Now, let $2x - 1 > 0$ i.e. $x > \frac{1}{2}$


$$\Rightarrow 2x - 1 = 3[x] + 2\{x\} \Rightarrow 2[x] + 2\{x\} - 1 = 3[x] + 2\{x\}$$

$$\Rightarrow [x] = -1 \Rightarrow -1 \leq x < 0 \text{ which is not a solution as } x > \frac{1}{2}$$

$$\Rightarrow x = \frac{1}{4} \text{ is the only solution.}$$



(j) **Identity function :**

The function $f: A \rightarrow A$ defined by $f(x) = x \quad \forall x \in A$ is called the identity of A and is denoted by I_A . It is easy to observe that identity function defined on R is a bijection.



(k) **Constant function :**

A function $f: A \rightarrow B$ is said to be a constant function if every element of A has the same f image in B . Thus $f: A \rightarrow B; f(x) = c, \forall x \in A, c \in B$ is a constant function. Note that the range of a constant function is a singleton.



Do yourself - 2 :

- (i) Let $\{x\}$ & $[x]$ denotes the fraction and integral part of a real number x respectively, then match the column.

Column-I

- (A) $[x^2] \geq 4$
 (B) $[x]^2 - 5[x] + 6 = 0$
 (C) $x = \{x\}$
 (D) $[x] < -5$

Column-II

- (p) $x \in [2, 4)$
 (q) $x \in (-\infty, -2] \cup [2, \infty)$
 (r) $x \in (-\infty, -5)$
 (s) $x \in \{-2\}$
 (t) $x \in [0, 1)$

5. ALGEBRAIC OPERATIONS ON FUNCTIONS :

If f & g are real valued functions of x with domain set A, B respectively, $f + g, f - g, (f \cdot g)$ & (f/g) as follows :

(a) $(f \pm g)(x) = f(x) \pm g(x)$ domain in each case is $A \cap B$

(b) $(f \cdot g)(x) = f(x) \cdot g(x)$ domain is $A \cap B$

(c) $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$ domain $A \cap B - \{x | g(x) = 0\}$

Illustration 6 : Find the domain of the following function :

(i) $y = \log_{(x-4)}(x^2 - 11x + 24)$ (ii) $f(x) = \log_2 \left(-\log_{1/2} \left(1 + \frac{1}{\sqrt[4]{x}} \right) - 1 \right)$

Solution :

(i) $y = \log_{(x-4)}(x^2 - 11x + 24)$

Here 'y' would assume real value if,

$x - 4 > 0$ and $\neq 1, x^2 - 11x + 24 > 0 \Rightarrow x > 4$ and $\neq 5, (x - 3)(x - 8) > 0$

$\Rightarrow x > 4$ and $\neq 5, x < 3$ or $x > 8 \Rightarrow x > 8 \Rightarrow \text{Domain } (y) = (8, \infty)$

(ii) We have $f(x) = \log_2 \left(-\log_{1/2} \left(1 + \frac{1}{\sqrt[4]{x}} \right) - 1 \right)$

$f(x)$ is defined if $-\log_{1/2} \left(1 + \frac{1}{\sqrt[4]{x}} \right) - 1 > 0$

or if $\log_{1/2} \left(1 + \frac{1}{\sqrt[4]{x}} \right) < -1$ or if $\left(1 + \frac{1}{\sqrt[4]{x}} \right) > (1/2)^{-1}$

or if $1 + \frac{1}{\sqrt[4]{x}} > 2$ or if $\frac{1}{\sqrt[4]{x}} > 1$ or if $x^{1/4} < 1$ or if $0 < x < 1$

$\therefore D(f) = (0, 1)$

Illustration 7 : Find the domain $f(x) = \frac{1}{\sqrt{[|x| - 5] - 11}}$ where $[.]$ denotes greatest integer function.

Solution : $[|x| - 5] > 11$

so $[|x| - 5] > 11$ or $[|x| - 5] < -11$

$[|x|] > 16$ $[|x|] < -6$

$|x| \geq 17$ or $|x| < -6$ (Not Possible)

$\Rightarrow x \leq -17$ or $x \geq 17$

so $x \in (-\infty, -17] \cup [17, \infty)$

Illustration 8 : Find the range of following functions :

(i) $f(x) = \frac{1}{8 - 3 \sin x}$ (ii) $f(x) = \log_{\sqrt{2}} (2 - \log_2 (16 \sin^2 x + 1))$

Solution : (i) $f(x) = \frac{1}{8 - 3 \sin x}$

$-1 \leq \sin x \leq 1$

\therefore Range of $f = \left[\frac{1}{11}, \frac{1}{5} \right]$

(ii) $f(x) = \log_{\sqrt{2}} (2 - \log_2 (16 \sin^2 x + 1))$

$1 \leq 16 \sin^2 x + 1 \leq 17$

$\therefore 0 \leq \log_2 (16 \sin^2 x + 1) \leq \log_2 17$

$\therefore 2 - \log_2 17 \leq 2 - \log_2 (16 \sin^2 x + 1) \leq 2$

Now consider $0 < 2 - \log_2 (16 \sin^2 x + 1) \leq 2$

$\therefore -\infty < \log_{\sqrt{2}} [2 - \log_2 (16 \sin^2 x + 1)] \leq \log_{\sqrt{2}} 2 = 2$

\therefore the range is $(-\infty, 2]$

Illustration 9 : Find the range of $f(x) = \frac{x - [x]}{1 + x - [x]}$, where $[.]$ denotes greatest integer function.

Solution : $y = \frac{x - [x]}{1 + x - [x]} = \frac{\{x\}}{1 + \{x\}}$
 $\therefore \frac{1}{y} = \frac{1}{\{x\}} + 1 \Rightarrow \frac{1}{\{x\}} = \frac{1-y}{y} \Rightarrow \{x\} = \frac{y}{1-y}$
 $0 \leq \{x\} < 1 \Rightarrow 0 \leq \frac{y}{1-y} < 1$
 Range = $[0, 1/2)$

Do yourself - 3 :

(i) Find domain of following functions :

(a) $f(x) = \sin(\sqrt{1-x^2}) + \sqrt{x+2} + \frac{1}{\log_{10}^{(x+1)}}$

(b) $f(x) = \sqrt{\frac{(2x+1)}{x^3 - 3x^2 + 2x}}$

(ii) Find range of following functions :

(a) $f(x) = \log_2(\log_{1/2}(x^2 + 4x + 4))$

(b) $f(x) = \frac{1}{2 - \cos 3x}$

6. EQUAL OR IDENTICAL FUNCTION :

Two function f & g are said to be equal if :

- (a) The domain of f = the domain of g
- (b) The co-domain of f = co-domain of g and
- (c) $f(x) = g(x)$, for every x belonging to their common domain (i.e. should have the same graph)

Illustration 10 : The functions $f(x) = \log(x-1) - \log(x-2)$ and $g(x) = \log\left(\frac{x-1}{x-2}\right)$ are identical when x lies in the interval

- (A) $[1, 2]$ (B) $[2, \infty)$ (C) $(2, \infty)$ (D) $(-\infty, \infty)$

Solution : Since $f(x) = \log(x-1) - \log(x-2)$.
 Domain of $f(x)$ is $x > 2$ or $x \in (2, \infty)$ (i)
 $g(x) = \log\left(\frac{x-1}{x-2}\right)$ is defined if $\frac{x-1}{x-2} > 0 \Rightarrow x \in (-\infty, 1) \cup (2, \infty)$ (ii)
 From (i) and (ii), $x \in (2, \infty)$. **Ans. (C)**

Do yourself - 4 :

(i) Are the following functions identical ?

$$(a) f(x) = \frac{x}{x^2} \text{ \& } \phi(x) = \frac{x^2}{x} \quad (b) f(x) = x \text{ \& } \phi(x) = \sqrt{x^2} \quad (c) f(x) = \log_{10} x^2 \text{ \& } \phi(x) = 2\log_{10}|x|$$

7. HOMOGENEOUS FUNCTIONS :

A function is said to be homogeneous with respect to any set of variables when each of its terms is of the same degree with respect to those variables.

For examples $5x^2 + 3y^2 - xy$ is homogenous in x & y . Symbolically if, $f(tx, ty) = t^n f(x, y)$ then $f(x, y)$ is homogeneous function of degree n .

Illustration 11 : Which of the following function is not homogeneous ?

$$(A) x^3 + 8x^2y + 7y^3 \quad (B) y^2 + x^2 + 5xy \quad (C) \frac{xy}{x^2 + y^2} \quad (D) \frac{2x - y + 1}{2y - x + 1}$$

Solution : It is clear that (D) does not have the same degree in each term. **Ans. (D)**

8. BOUNDED FUNCTION :

A function is said to be bounded if there exists a finite M such that $|f(x)| \leq M, \forall x \in D_f$.

9. IMPLICIT & EXPLICIT FUNCTION :

A function defined by an equation not solved for the dependent variable is called an **implicit function**. e.g. the equations $x^3 + y^3 = 1$ & $x^y = y^x$, defines y as an implicit function. If y has been expressed in terms of x alone then it is called an **Explicit function**.

Illustration 12 : Which of the following function is implicit function ?

$$(A) y = \frac{x^2 + e^x + 5}{\sqrt{1 - \cos^{-1} x}} \quad (B) y = x^2 \quad (C) xy - \sin(x + y) = 0 \quad (D) y = \frac{x^2 \log x}{\sin x}$$

Solution : It is clear that in (C) y is not clearly expressed in x . **Ans. (C)**

Do yourself - 5 :

(i) Find the boundness of the function $f(x) = \frac{x^2}{x^4 + 1}$

(ii) Which of the following function is implicit function ?

$$(A) xy - \cos(x + y) = 0$$

$$(B) y = x^3$$

$$(C) y = \log(x^2 + x + 1)$$

$$(D) y = |x|$$

(iii) Convert the implicit form into the explicit function :

$$(a) xy = 1 \quad (b) x^2y = 1.$$

10. APPLICATIONS OF FUNCTIONAL RULE :

Illustration 13 : Determine all functions f satisfying the functional relation.

$$f(x) + f\left(\frac{1}{1-x}\right) = \frac{2(1-2x)}{x(1-x)} \text{ where } x \in \mathbb{R} - \{0, 1\}$$

Solution : Given $f(x) + f\left(\frac{1}{1-x}\right) = \frac{2(1-2x)}{x(1-x)} = \frac{2}{x} - \frac{2}{1-x}$... (i)

Replacing x by $\frac{1}{1-x}$ we obtain

$$f\left(\frac{1}{1-x}\right) + f\left(\frac{1}{1-\frac{1}{1-x}}\right) = 2(1-x) - \frac{2}{1-\frac{1}{1-x}}$$

$$\Rightarrow f\left(\frac{1}{1-x}\right) + f\left(1 - \frac{1}{x}\right) = -2x + \frac{2}{x}$$
 ... (ii)

Again replacing x by $\left(1 - \frac{1}{x}\right)$ in (i) we obtain

$$\Rightarrow f\left(1 - \frac{1}{x}\right) + f\left(\frac{1}{1-\left(1-\frac{1}{x}\right)}\right) = \frac{2}{1-\frac{1}{x}} - \frac{2}{1-\left(1-\frac{1}{x}\right)}$$

$$\Rightarrow f\left(1 - \frac{1}{x}\right) + f(x) = \frac{2x}{x-1} - 2x$$
 ... (iii)


subtracting (ii) from (i) then

$$f(x) - f\left(1 - \frac{1}{x}\right) = 2x - \frac{2}{1-x}$$

Now adding (iii) and (iv) we get

$$2f(x) = \frac{2x}{x-1} - \frac{2}{1-x}$$


$$\Rightarrow f(x) = \frac{x+1}{x-1}$$



(v) Drawing the graph of $y = f(|x|)$ from the known graph of $y = f(x)$


It is clear that, $f(|x|) = \begin{cases} f(x), & x \geq 0 \\ f(-x), & x < 0 \end{cases}$. Thus $f(|x|)$ would be an even function, graph of $f(|x|)$ and

$f(x)$ would be identical in the first and the fourth quadrants (as $x \geq 0$) and as such the graph of $f(|x|)$ would be symmetric about the y-axis (as $|x|$ is even).




(vi) Drawing the graph of $|y| = f(x)$ from the known graph of $y = f(x)$

Clearly $|y| \geq 0$. If $f(x) < 0$, graph of $|y| = f(x)$ would not exist. And if $f(x) \geq 0$, $|y| = f(x)$ would give $y = \pm f(x)$. Hence graph of $|y| = f(x)$ would exist only in the regions where $f(x)$ is non-negative and will be reflected about the x-axis only in those regions.




(vii) Drawing the graph of $y = f(x + a)$, $a \in \mathbb{R}$ from the known graph of $y = f(x)$



(i) If $a > 0$, shift the graph of $f(x)$ through 'a' units towards left of $f(x)$.


(ii) If $a < 0$, shift the graph of $f(x)$ through 'a' units towards right of $f(x)$.

(viii) Drawing the graph of $y = af(x)$ from the known graph of $y = f(x)$



It is clear that the corresponding points (points with same x co-ordinates) would have their ordinates in the ratio of 1 : a.

(ix) Drawing the graph of $y = f(ax)$ from the known graph of $y = f(x)$.



Let us take any point $x_0 \in \text{domain of } f(x)$. Let $ax = x_0$ or $x = \frac{x_0}{a}$.

Clearly if $0 < a < 1$, then $x > x_0$ and $f(x)$ will stretch by $\frac{1}{a}$ units along the x-axis and if $a > 1$, $x < x_0$, then $f(x)$ will compress by 'a' units along the x-axis.

Note :

- (i) A function $h(x)$ is defined as $h(x) = \max. \{f(x), g(x)\}$ then

$$h(x) = \begin{cases} f(x) & f(x) \geq g(x) \\ g(x) & g(x) > f(x) \end{cases}$$

- (ii) A function $h(x)$ is defined as $h(x) = \min. \{f(x), g(x)\}$ then

$$h(x) = \begin{cases} f(x) & f(x) \leq g(x) \\ g(x) & g(x) < f(x) \end{cases}$$

Illustration 14: Find $f(x) = \max \{1 + x, 1 - x, 2\}$.

Solution : From the graph it is clear that

$$f(x) = \begin{cases} 1 - x & x < -1 \\ 2 & -1 \leq x \leq 1 \\ 1 + x & x > 1 \end{cases}$$





Illustration 15: Draw the graph of $y = |2 - |x - 1||$.

Solution :






Illustration 16: Draw the graph of $y = 2 - \frac{4}{|x-1|}$

Solution :




Illustration 17: Draw the graph of $y = |e^{|x|} - 2|$

Solution :





Illustration 18: Draw the graph of $f(x) = \cos x \cos(x+2) - \cos^2(x+1)$.

Solution :

$$\begin{aligned}
 f(x) &= \cos x \cos(x+2) - \cos^2(x+1) \\
 &= \frac{1}{2} [\cos(2x+2) + \cos 2] - \frac{1}{2} [\cos(2x+2) + 1] \\
 &= \frac{1}{2} \cos 2 - \frac{1}{2} < 0.
 \end{aligned}$$



Do yourself - 6 :

(i) Draw graph of following functions :

(a) $y = \lfloor \ln |x| \rfloor + 1$


(b) $y = \min. \{x^2 + 1, 3 - x\}$

12. CLASSIFICATION OF FUNCTIONS :**One-One Function (Injective mapping) :**

A function $f : A \rightarrow B$ is said to be a one-one function or injective mapping if different elements of A have different f images in B .

Thus there exist $x_1, x_2 \in A$ & $f(x_1), f(x_2) \in B$, $f(x_1) = f(x_2) \Leftrightarrow x_1 = x_2$ or $x_1 \neq x_2 \Leftrightarrow f(x_1) \neq f(x_2)$.


Diagrammatically an injective mapping can be shown as

**Many-one function (not injective) :**

A function $f : A \rightarrow B$ is said to be a many one function if two or more elements of A have the same f image in B .

Thus $f : A \rightarrow B$ is many one there exist $x_1, x_2 \in A$, $f(x_1) = f(x_2)$ but $x_1 \neq x_2$.

Diagrammatically a many one mapping can be shown as


**Note :**

- (i) If a line parallel to x-axis cuts the graph of the function at most at one point, then the function is one-one.
- (ii) If any line parallel to x-axis cuts the graph of the function at least at two points, then f is many-one.
- (iii) If continuous function $f(x)$ is always increasing or decreasing in whole domain, then $f(x)$ is one-one.
- (iv) All linear functions are one-one.
- (v) All trigonometric functions in their domain are many one
- (vi) All even degree polynomials are many one
- (vii) Linear by Linear is one-one
- (viii) Quadratic by quadratic with no common factor is many one.

Onto function (Surjective mapping) :

If the function $f : A \rightarrow B$ is such that each element in B (co-domain) is the f image of at least one element in A , then we say that f is a function from A 'onto' B . Thus $f : A \rightarrow B$ is surjective iff $\forall b \in B, \exists$ some $a \in A$ such that $f(a) = b$.

Diagrammatically surjective mapping can be shown as




Note that : if range is same as co-domain, then $f(x)$ is onto.

Into function :

If $f: A \rightarrow B$ is such that there exists atleast one element in co-domain which is not the image of any element in domain, then $f(x)$ is into.

Diagrammatically into function can be shown as




Note :

- A polynomial function of degree even define from $\mathbb{R} \rightarrow \mathbb{R}$ will always be into.
- A polynomial function of degree odd defined from $\mathbb{R} \rightarrow \mathbb{R}$ will always be onto.
- Quadratic by quadratic without any common factor define from $\mathbb{R} \rightarrow \mathbb{R}$ is always an into function.


Thus a function can be one of these four types :

- one-one onto (injective & surjective)


(also known as **Bijjective** mapping)



- one-one into (injective but not surjective)



- many-one onto (surjective but not injective)



- many-one into (neither surjective nor injective)




Illustration 19 : Let $A = \{x : -1 \leq x \leq 1\}$ B be a mapping $f: A \rightarrow B$. For each of the following functions from A to B , find whether it is bijective or non-bijective.

(a) $f(x) = x|x|$

(b) $f(x) = x^3$

(c) $f(x) = \sin \frac{\pi x}{2}$

Solution :

$$(a) \quad f(x) = x|x| = \begin{cases} -x^2, & -1 < x < 0 \\ x^2, & 0 \leq x < 1 \end{cases}$$

Graphically,

The graph shows $f(x)$ is one-one, as the straight line parallel to x -axis cuts only at one point.

Here, range

$$f(x) \in [-1, 1]$$

Thus, range = co-domain

Hence, onto.

Therefore, $f(x)$ is one-one onto or (Bijective).

$$(b) \quad f(x) = x^3,$$

Graphically;

Graph shows $f(x)$ is one-one onto

(i.e. Bijective)




[as explained in above example]

$$(c) \quad f(x) = \sin \frac{\pi x}{2}$$

Graphically;

Which shows $f(x)$ is one-one and onto as range

= co-domain.

Therefore, $f(x)$ is bijective.**Illustration 20 :**Let $f : \mathbb{N} \rightarrow \mathbb{Z}$ be a function defined as $f(x) = x - 1000$. Show that $f(x)$ is an into function.**Solution :**

$$\text{Let } f(x) = y = x - 1000 \Rightarrow x = y + 1000 = g(y) \text{ (say)}$$

here $g(y)$ is defined for each $y \in \mathbb{Z}$, but $g(y) \notin \mathbb{N}$ for $y \leq -1000$.Hence $f(x)$ is into.**Illustration 21 :**Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = x + \sqrt{x^2}$, then f is

(A) injective

(B) surjective

(C) bijective

(D) None of these

Solution :

$$\text{We have, } f(x) = x + \sqrt{x^2} = x + |x|$$

Clearly, f is not one-one as $f(-1) = f(-2) = 0$ and $-1 \neq -2$ Also, f is not onto as $f(x) \geq 0 \quad \forall x \in \mathbb{R}$ \therefore range of $f = (0, \infty) \subset \mathbb{R}$ **Ans. (D)****Illustration 22 :**Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined as $f(x) = 2x^3 + 6x^2 + 12x + 3 \cos x - 4 \sin x$; then f is -

(A) Injective

(B) Surjective

(C) Bijective

(D) Not Surjective

Solution :

$$\text{We have } f(x) = 2x^3 + 6x^2 + 12x + 3 \cos x - 4 \sin x$$

$$\Rightarrow f'(x) = 6x^2 - 12x + 12 - 3 \sin x - 4 \cos x$$

$$f'(x) = \underbrace{6(x-1)^2 + 6}_{g(x)} - \underbrace{(3 \sin x + 4 \cos x)}_{h(x)}$$

$$\text{range of } g(x) = [6, \infty)$$

$$\text{range of } h(x) = [-5, 5]$$

hence $f'(x)$ always lies in the interval $[1, \infty)$

$$\Rightarrow f'(x) > 0$$

Hence $f(x)$ is increasing i.e. one-one function

Now $x \rightarrow \infty \Rightarrow f \rightarrow \infty$ & $x \rightarrow -\infty \Rightarrow f \rightarrow -\infty$ & $f(x)$ is continuous

hence its range is $\mathbb{R} \Rightarrow f$ is onto so f is bijective.

Ans. (C)

Illustration 23 : Let $f(x) = \frac{x^2 + 3x + a}{x^2 + x + 1}$, where $f : \mathbb{R} \rightarrow \mathbb{R}$. Find the value of parameter 'a' so that the given function is one-one.

Solution : $f(x) = \frac{x^2 + 3x + a}{x^2 + x + 1}$

$$f'(x) = \frac{(x^2 + x + 1)(2x + 3) - (x^2 + 3x + a)(2x + 1)}{(x^2 + x + 1)^2} = \frac{-2x^2 + 2x(1 - a) + (3 - a)}{(x^2 + x + 1)^2}$$

Let, $g(x) = -2x^2 + 2x(1 - a) + (3 - a)$

$g(x)$ will be negative if $4(1 - a)^2 + 8(3 - a) < 0$

$$\Rightarrow 1 + a^2 - 2a + 6 - 2a < 0 \Rightarrow (a - 2)^2 + 3 < 0$$

which is not possible. Therefore function is not monotonic.

Hence, no value of a is possible.

Do yourself - 7 :

- (i) Is the function $f : \mathbb{N} \rightarrow \mathbb{N}$ (the set of natural numbers) defined by $f(x) = 2x + 3$ surjective ?
- (ii) Let $A = \mathbb{R} - \{3\}$, $\mathbb{R} = \mathbb{R} - \{1\}$ and let $f : A \rightarrow B$ defined by $f(x) = \frac{x-2}{x-3}$. Check whether the function $f(x)$ is bijective or not.
- (iii) A mapping $f : A \rightarrow [-1, 1]$ defined by $f(x) = \sin x$, $\forall x \in \mathbb{R}$, where A is a subset of \mathbb{R} (the set of all real numbers) is one-one and onto if A is the interval, then A belongs to

- (A) $[0, 2\pi]$ (B) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (C) $[-\pi, \pi]$ (D) $[0, \pi]$

13. COMPOSITE OF UNIFORMLY & NON-UNIFORMLY DEFINED FUNCTION:

Let $f : A \rightarrow B$ & $g : B \rightarrow C$ be two functions. Then the function $g \circ f : A \rightarrow C$ defined by $(g \circ f)(x) = g(f(x))$ $\forall x \in A$ is called the composite of the two functions f & g .

Diagrammatically $\xrightarrow{x} \boxed{f} \xrightarrow{f(x)} \boxed{g} \xrightarrow{g(f(x))}$

Thus the image of every $x \in A$ under the function $g \circ f$ is the g -image of f -image of x .

Note that $g \circ f$ is defined only if $\forall x \in A$, $f(x)$ is an element of the domain of 'g' so that we can take its g -image.

Properties of composite functions:

- (a) In general composite of functions is not commutative i.e. $g \circ f \neq f \circ g$.

- (b) The composition of functions is associative i.e. if f, g, h are three functions such that $fo(goh)$ & $(fog)oh$ are defined, then $fo(goh) = (fog)oh$.
- (c) The composition of two bijections is a bijection i.e. if f & g are two bijections such that gof is defined, then gof is also a bijection.

Illustration 24: If $f(x) = x^2 + 1$, $g(x) = \frac{1}{x-1}$, then find $(fog)(x)$ and $(gof)(x)$.

Solution : Given, $f(x) = x^2 + 1$ (1) $g(x) = \frac{1}{x-1}$... (2)

$$\text{Now } (fog)(x) = f(g(x)) = f\left(\frac{1}{x-1}\right) = f(z), \text{ where } z = \frac{1}{x-1}$$

$$= z^2 + 1 \quad [\because f(x) = x^2 + 1]$$

$$= \left(\frac{1}{x-1}\right)^2 + 1 = \frac{1}{(x-1)^2} + 1$$

Note : Domain of $(fog)(x)$ is $x \in \mathbb{R} - \{1\}$

$$(gof)(x) = g(f(x)) = g(x^2 + 1) = g(u), \text{ where } u = x^2 + 1$$

$$= \frac{1}{u-1} = \frac{1}{x^2 + 1 - 1} = \frac{1}{x^2}$$

Note : Domain of $(gof)(x)$ is $x \in \mathbb{R} - \{0\}$

Illustration 25: If f be the greatest integer function and g be the modulus function, then

$$(gof)\left(-\frac{5}{3}\right) - (fog)\left(-\frac{5}{3}\right) =$$

- (A) 1 (B) -1 (C) 2 (D) 4

Solution : Given $(gof)\left(-\frac{5}{3}\right) - (fog)\left(-\frac{5}{3}\right) = g\left\{f\left(-\frac{5}{3}\right)\right\} - f\left\{g\left(-\frac{5}{3}\right)\right\} = g(-2) - f\left(\frac{5}{3}\right) = 2 - 1 = 1$

Ans.(A)

Illustration 26: Let $f(x) = \begin{cases} x+1, & x \leq 1 \\ 2x+1, & 1 < x \leq 2 \end{cases}$ and $g(x) = \begin{cases} x^2, & -1 \leq x < 2 \\ x+2, & 2 \leq x \leq 3 \end{cases}$, find (fog)

Solution : $f(g(x)) = \begin{cases} g(x)+1, & g(x) \leq 1 \\ 2g(x)+1, & 1 < g(x) \leq 2 \end{cases}$

Here, $g(x)$ becomes the variable that means we should draw the graph.

It is clear that $g(x) \leq 1$; $\forall x \in [-1, 1]$

and $1 < g(x) \leq 2$; $\forall x \in (1, \sqrt{2}]$

$$\Rightarrow f(g(x)) = \begin{cases} x^2 + 1, & -1 \leq x \leq 1 \\ 2x^2 + 1, & 1 < x \leq \sqrt{2} \end{cases}$$




Illustration 27: Find the domain and range of $h(x) = g(f(x))$, where

$$f(x) = \begin{cases} [x], & -2 \leq x \leq -1 \\ |x| + 1, & -1 < x \leq 2 \end{cases} \text{ and } g(x) = \begin{cases} [x], & -\pi \leq x \leq 0 \\ \sin x, & 0 \leq x \leq \pi \end{cases}, [\cdot] \text{ denotes the greatest integer function.}$$


Solution : $h(x) = g(f(x)) = \begin{cases} [f(x)], & -\pi \leq f(x) < 0 \\ \sin(f(x)), & 0 \leq f(x) \leq \pi \end{cases}$

From graph of $f(x)$, we get

$$h(x) = \begin{cases} [[x]], & -2 \leq x \leq -1 \\ \sin(|x| + 1), & -1 < x \leq 2 \end{cases}$$

\Rightarrow Domain of $h(x)$ is $[-2, 2]$

and Range of $h(x)$ is $\{-2, -1\} \cup [\sin 3, 1]$



Do yourself - 8 :

(i) $f(x) = x^3 - x$ & $g(x) = \sin 2x$, find

(a) $f(f(1))$

(b) $f(f(-1))$

(c) $f\left(g\left(\frac{\pi}{2}\right)\right)$

(d) $f\left(g\left(\frac{\pi}{4}\right)\right)$

(e) $g(f(1))$

(f) $g\left(g\left(\frac{\pi}{2}\right)\right)$

(ii) If $f(x) = \begin{cases} x+1; & 0 \leq x < 2 \\ |x|; & 2 \leq x < 3 \end{cases}$, then find $f \circ f(x)$.

14. INVERSE OF A FUNCTION :

Let $f : A \rightarrow B$ be a one-one & onto function, then there exists a unique function $g : B \rightarrow A$ such that $f(x) = y \Leftrightarrow g(y) = x, \forall x \in A \text{ & } y \in B$. Then g is said to be inverse of f .

Thus $g = f^{-1} : B \rightarrow A = \{(f(x), x) | (x, f(x)) \in f\}$.

Properties of inverse function :


(a) The inverse of a bijection is unique.

(b) If $f : A \rightarrow B$ is a bijection & $g : B \rightarrow A$ is the inverse of f , then $f \circ g = I_B$ and $g \circ f = I_A$, where I_A & I_B are identity functions on the sets A & B respectively. If $f \circ f = I$, then f is inverse of itself.

(c) The inverse of a bijection is also a bijection.

(d) If f & g are two bijections $f : A \rightarrow B, g : B \rightarrow C$ then the inverse of $g \circ f$ exists and $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$.

- (e) Since $f(a) = b$ if and only if $f^{-1}(b) = a$, the point (a, b) is on the graph of 'f' if and only if the point (b, a) is on the graph of f^{-1} . But we get the point (b, a) from (a, b) by reflecting about the line $y = x$.



The graph of f^{-1} is obtained by reflecting the graph of f about the line $y = x$.

Drawing the graph of $y = f^{-1}(x)$ from the known graph of $y = f(x)$

For drawing the graph of $y = f^{-1}(x)$ take the reflection of $y = f(x)$ about the line $y = x$. The reflected part would give us the graph of $y = f^{-1}(x)$.

e.g. let us draw the graph of $y = \sin^{-1}x$. We know that $y = f(x) = \sin x$ is invertible if $f :$

$$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1]$$

$$\Rightarrow \text{the inverse mapping would be } f^{-1} : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right].$$




Illustration 28: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = (e^x - e^{-x})/2$. Is $f(x)$ invertible? If so, find its inverse.

Solution : Let us check for invertibility of $f(x)$:

(a) One-One :

$$f(x) = \frac{1}{2}(e^x - e^{-x}) \Rightarrow f'(x) = \frac{1}{2}(e^x + e^{-x})$$

$$\Rightarrow f'(x) > 0, f(x) \text{ is increasing function}$$

$$\therefore f(x) \text{ is one-one function.}$$

(b) Onto :

As x tends to larger and larger values so does $f(x)$ and when $x \rightarrow \infty$, $f(x) \rightarrow \infty$.

Similarly as $x \rightarrow -\infty$, $f(x) \rightarrow -\infty$ i.e. $-\infty < f(x) < \infty$ so long as $x \in (-\infty, \infty)$

Hence the range of f is same as the set R . Therefore $f(x)$ is onto.

Since $f(x)$ is both one-one and onto, $f(x)$ is invertible.

(c) To find $f^{-1}(x)$: Interchange x & y

$$\frac{e^y - e^{-y}}{2} = x \Rightarrow e^{2y} - 2xe^y - 1 = 0$$

$$\Rightarrow e^y = \frac{2x \pm \sqrt{4x^2 + 4}}{2} \Rightarrow e^y = x \pm \sqrt{1 + x^2}$$

Since $e^y > 0$, hence negative sign is ruled out and

$$\text{Hence } e^y = x + \sqrt{1 + x^2}$$

Taking logarithm, we have $y = \ln(x + \sqrt{1 + x^2})$ or $f^{-1}(x) = \ln(x + \sqrt{1 + x^2})$

Illustration 29 : Find the inverse of the function $f(x) = \log_a \left(x + \sqrt{x^2 + 1} \right)$; $a > 1$ and assuming it to be an onto function.

Solution : Given $f(x) = \log_a \left(x + \sqrt{x^2 + 1} \right)$

$$\therefore f'(x) = \frac{\log_a e}{\sqrt{1 + x^2}} > 0$$

which is a strictly increasing functions.

Thus, $f(x)$ is injective, given that $f(x)$ is onto. Hence the given function $f(x)$ is invertible.

Interchanging x & y

$$\Rightarrow \log_a \left(y + \sqrt{y^2 + 1} \right) = x$$

$$\Rightarrow y + \sqrt{y^2 + 1} = a^x \quad \dots(i)$$

$$\text{and } \sqrt{y^2 + 1} - y = a^{-x} \quad \dots(ii)$$

$$\text{From (i) and (ii), we get } y = \frac{1}{2}(a^x - a^{-x}) \text{ or } f^{-1}(x) = \frac{1}{2}(a^x - a^{-x})$$

Illustration 30 : Find the inverse of the function $f(x) = \ln(x^2 + 3x + 1)$; $x \in [1, 3]$ and assuming it to be an onto function.

Solution : Given $f(x) = \ln(x^2 + 3x + 1)$

$$\therefore f'(x) = \frac{2x + 3}{x^2 + 3x + 1} > 0 \forall x \in [1, 3]$$

Which is a strictly increasing function. Thus $f(x)$ is injective, given that $f(x)$ is onto. Hence the given function $f(x)$ is invertible.

Interchanging x & y

$$\Rightarrow (y)^2 + 3(y) + 1 - e^x = 0$$

$$\therefore y = \frac{-3 \pm \sqrt{9 - 4 \cdot (1 - e^x)}}{2} = \frac{-3 \pm \sqrt{5 + 4e^x}}{2}$$

$$\Rightarrow y = \frac{-3 + \sqrt{5 + 4e^x}}{2} \quad (\text{as } y \in [1, 3])$$

$$\text{Hence } f^{-1}(x) = \frac{-3 + \sqrt{5 + 4e^x}}{2}$$

Do yourself - 9 :

- (i) Let $f : [-1, 1] \rightarrow [-1, 1]$ defined by $f(x) = x|x|$, find $f^{-1}(x)$.
 (ii) $f(x) = 1 + \ln(x + 2)$, find $f^{-1}(x)$.

15. ODD & EVEN FUNCTIONS :

Consider a function $f(x)$ such that both x and $-x$ are in its domain then

If $\begin{cases} f(-x) = f(x) & \text{then } f \text{ is said to be an even function} \\ f(-x) = -f(x) & \text{then } f \text{ is said to be an odd function} \end{cases}$

Note :

- $f(x) - f(-x) = 0 \Rightarrow f(x)$ is even & $f(x) + f(-x) = 0 \Rightarrow f(x)$ is odd.
- A function may neither be odd nor even.
- The only function which is defined on the entire number line & is even and odd at the same time is $f(x) = 0$.
- Every constant function is even function.
- Inverse of an even function is not defined.
- Every even function is symmetric about the y -axis & every odd function is symmetric about the origin.

Special Note : If a function $f(x)$ is defined as $f(a+x) = f(a-x)$ then this function is symmetric about line $x = a$

- (vii) Every function which has ' $-x$ ' in its domain whenever ' x ' is in its domain, can be expressed as the sum of an even & an odd function .

$$\text{i.e. } f(x) = \underbrace{\frac{f(x) + f(-x)}{2}}_{\text{EVEN}} + \underbrace{\frac{f(x) - f(-x)}{2}}_{\text{ODD}}$$

(viii) If $f(x)$ is odd and defined at $x = 0$, then $f(0) = 0$.

$f(x)$	$g(x)$	$f(x) + g(x)$	$f(x) - g(x)$	$f(x) \cdot g(x)$	$f(x)/g(x)$	$(g \circ f)(x)$	$(f \circ g)(x)$
odd	odd	odd	odd	even	even	odd	odd
even	even	even	even	even	even	even	even
odd	even	neither odd nor even	neither odd nor even	odd	odd	even	even
even	odd	neither odd nor even	neither odd nor even	odd	odd	even	even

Illustration 31 : Which of the following functions is (are) even, odd or neither :

(i) $f(x) = x^2 \sin x$ (ii) $f(x) = \sin x - \cos x$ (iii) $f(x) = \frac{e^x + e^{-x}}{2}$

Solution :

(i) $f(-x) = (-x)^2 \sin(-x) = -x^2 \sin x = -f(x)$. Hence $f(x)$ is odd.

(ii) $f(-x) = \sin(-x) - \cos(-x) = -\sin x - \cos x$.

Hence $f(x)$ is neither even nor odd.

(iii) $f(-x) = \frac{e^{-x} + e^{-(-x)}}{2} = \frac{e^{-x} + e^x}{2} = f(x)$. Hence $f(x)$ is even

Illustration 32 : Identify the given functions as odd, even or neither :

(i) $f(x) = \frac{x}{e^x - 1} + \frac{x}{2} + 1$ (ii) $f(x + y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$

Solution :

(i) $f(x) = \frac{x}{e^x - 1} + \frac{x}{2} + 1$

Clearly domain of $f(x)$ is $\mathbb{R} \sim \{0\}$. We have,

$$\begin{aligned} f(-x) &= \frac{-x}{e^{-x} - 1} - \frac{x}{2} + 1 = \frac{-e^x \cdot x}{1 - e^x} - \frac{x}{2} + 1 = \frac{(e^x - 1 + 1)x}{(e^x - 1)} - \frac{x}{2} + 1 \\ &= x + \frac{x}{e^x - 1} - \frac{x}{2} + 1 = \frac{x}{e^x - 1} + \frac{x}{2} + 1 = f(x) \end{aligned}$$

Hence $f(x)$ is an even function.

(ii) $f(x + y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$

Replacing x, y by zero, we get $f(0) = 2f(0) \Rightarrow f(0) = 0$

Replacing y by $-x$, we get $f(x) + f(-x) = f(0) = 0 \Rightarrow f(x) = -f(-x)$

Hence $f(x)$ is an odd function.

Do yourself - 10 :

(i) Which of the following functions is (are) even, odd or neither :

(a) $f(x) = x^3 \sin 3x$ (b) $f(x) = \frac{e^{x^2} + e^{-x^2}}{2x}$

(c) $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ (d) $f(x) = x^2 + 2^x$

16. PERIODIC FUNCTION :

A function $f(x)$ is called periodic if there exists a positive number T such that $f(x+T) = f(x) = f(x-T)$, for all values of x within the domain of f . Smallest positive T (if exists) is called fundamental period of function $f(x)$.

Note :

- (i) Odd powers of $\sin x$, $\cos x$, $\sec x$, $\csc x$ are periodic with period 2π .
- (ii) None zero integral powers of $\tan x$, $\cot x$ are periodic with period π .
- (iii) Non zero even powers or modulus of $\sin x$, $\cos x$, $\sec x$, $\csc x$ are periodic with period π .
- (iv) $f(T) = f(0) = f(-T)$, where ' T ' is the period.
- (v) if $f(x)$ has a period T then **$f(ax+b)$ has a period $T/|a|$** ($a \neq 0$).

Proof : Let $f(x+T) = f(x)$ and $f[a(x+T') + b] = f(ax+b)$
 $f(ax+b+aT') = f(ax+b)$

$$f(y+aT') = f(y) = f(y+T) \Rightarrow T = aT' \Rightarrow T' = \frac{T}{a}$$

- (vi) If $f(x)$ & $g(x)$ are periodic with period T_1 & T_2 respectively, then a period (need not be fundamental) of $f(x) \pm g(x)$ is L.C.M. of (T_1, T_2) .
- (a) LCM of T_1 & T_2 is defined when T_1/T_2 is rational.

$$(b) \text{ LCM of } \left\{ \frac{a}{b}, \frac{p}{q} \right\} = \frac{\text{LCM of } (a, p)}{\text{HCF of } (b, q)}$$

In case if there exists a positive K such that $K < \text{LCM of } T_1 \text{ and } T_2$ and overall function repeats itself after every K , then fundamental period of the function will be K .

- (vii) Every constant function is always periodic, whose fundamental period is undefined.
- (viii) Periodic functions are non invertible.

Illustration 33 : Find the fundamental periods (if periodic) of the following functions, where $[.]$ denotes the greatest integer function

$$(i) f(x) = e^{\ell n(\sin x)} + \tan^3 x - \csc(3x-5) \quad (ii) f(x) = x - [x-b], b \in \mathbb{R}$$

$$(iii) f(x) = \frac{|\sin x + \cos x|}{|\sin x| + |\cos x|} \quad (iv) f(x) = \tan \frac{\pi}{2} [x]$$

$$(v) f(x) = \cos(\sin x) + \cos(\cos x) \quad (vi) f(x) = \frac{(1+\sin x)(1+\sec x)}{(1+\cos x)(1+\csc x)}$$

$$(vii) f(x) = e^{x-[x]+|\cos \pi x|+|\cos 2\pi x|+\dots+\|\cos n\pi\|}$$

Solution :

$$(i) f(x) = e^{\ell n(\sin x)} + \tan^3 x - \csc(3x-5)$$

Period of $e^{\ell n \sin x} = 2\pi$, $\tan^3 x = \pi$

$$\csc(3x-5) = \frac{2\pi}{3}$$

$$\therefore \text{Period} = 2\pi$$

$$(ii) f(x) = x - [x-b] = b + \{x-b\}$$

$$\therefore \text{Period} = 1$$

$$(iii) f(x) = \frac{|\sin x + \cos x|}{|\sin x| + |\cos x|}$$

Since period of $|\sin x + \cos x| = \pi$ and period of $|\sin x| + |\cos x|$ is $\frac{\pi}{2}$. Hence $f(x)$ is periodic with π as its period

$$(iv) f(x) = \tan \frac{\pi}{2} [x]$$

$$\tan \frac{\pi}{2} [x + T] = \tan \frac{\pi}{2} [x] \Rightarrow \frac{\pi}{2} [x + T] = n\pi + \frac{\pi}{2} [x]$$

$$\therefore T = 2$$

$$\therefore \text{Period} = 2$$

(v) Let $f(x)$ is periodic then $f(x + T) = f(x)$

$$\Rightarrow \cos(\sin(x + T)) + \cos(\cos(x + T)) = \cos(\sin x) + \cos(\cos x)$$

If $x = 0$ then $\cos(\sin T) + \cos(\cos T)$

$$= \cos(0) + \cos(1) = \cos\left(\cos \frac{\pi}{2}\right) + \cos\left(\sin \frac{\pi}{2}\right)$$

$$\text{On comparing } T = \frac{\pi}{2}$$

$$(vi) f(x) = \frac{(1 + \sin x)(1 + \sec x)}{(1 + \cos x)(1 + \csc x)} = \frac{(1 + \sin x)(1 + \cos x) \sin x}{\cos x (1 + \sin x)(1 + \cos x)}$$

$$\Rightarrow f(x) = \tan x$$

Hence $f(x)$ has period π .

$$(vii) f(x) = e^{x - [x] + |\cos \pi x| + |\cos 2\pi x| + \dots + |\cos n\pi x|}$$

$$\text{Period of } x - [x] = 1$$

$$\text{Period of } |\cos \pi x| = 1$$

$$\text{Period of } |\cos 2\pi x| = \frac{1}{2}$$

.....

$$\text{Period of } |\cos n\pi x| = \frac{1}{n}$$

So period of $f(x)$ will be L.C.M. of all period = 1

Illustration 34: Find the fundamental periods (if periodic) of the following functions, where $[.]$ denotes the greatest integer function

$$(i) f(x) = e^{x - [x]} + \sin x \quad (ii) f(x) = \sin \frac{\pi x}{\sqrt{2}} + \cos \frac{\pi x}{\sqrt{3}} \quad (iii) f(x) = \sin \frac{\pi x}{\sqrt{3}} + \cos \frac{\pi x}{2\sqrt{3}}$$

Solution :

$$(i) \text{ Period of } e^{x - [x]} = 1$$

$$\text{period of } \sin x = 2\pi$$

\therefore L.C.M. of rational and an irrational number does not exist.

\therefore not periodic.

$$(ii) \text{ Period of } \sin \frac{\pi x}{\sqrt{2}} = \frac{2\pi}{\pi/\sqrt{2}} = 2\sqrt{2}$$

$$\text{Period of } \cos \frac{\pi x}{\sqrt{3}} = \frac{2\pi}{\pi/\sqrt{3}} = 2\sqrt{3}$$

\therefore L.C.M. of two different kinds of irrational number does not exist.
 \therefore not periodic.

$$(iii) \text{ Period of } \sin \frac{\pi x}{\sqrt{3}} = \frac{2\pi}{\pi/\sqrt{3}} = 2\sqrt{3}$$

$$\text{Period of } \cos \frac{\pi x}{2\sqrt{3}} = \frac{2\pi}{\pi/2\sqrt{3}} = 4\sqrt{3}$$

\therefore L.C.M. of two similar irrational number exist.

\therefore Periodic with period = $4\sqrt{3}$

Ans.

Do yourself - 11 :

(i) Find the fundamental periods (if periodic) of the following functions.

(a) $f(x) = \ell n(\cos x) + \tan^3 x$.

(b) $f(x) = e^{x-[x]}$, [.] denotes greatest integer function

(c) $f(x) = \left| \sin \frac{x}{2} \right| + \left| \cos \frac{x}{2} \right|$

17. GENERAL :

If x, y are independent variables and f is a continuous function, then :

(a) $f(xy) = f(x) + f(y) \Rightarrow f(x) = k \ell n x$

(b) $f(xy) = f(x) \cdot f(y) \Rightarrow f(x) = x^n, n \in \mathbb{R}$ or $f(x) = 0$

(c) $f(x+y) = f(x) \cdot f(y) \Rightarrow f(x) = a^{kx}$ or $f(x) = 0$

(d) $f(x+y) = f(x) + f(y) \Rightarrow f(x) = kx$, where k is a constant.


Miscellaneous Illustration :

Illustration 35: ABCD is a square of side ℓ . A line parallel to the diagonal BD at a distance 'x' from the vertex A cuts two adjacent sides. Express the area of the segment of the square with A at a vertex, as a function of x . Find this area at $x = 1/\sqrt{2}$ and at $x = 2$, when $\ell = 2$.

Solution : There are two different situations

Case-I : when $x = AP \leq OA$, i.e., $x \leq \frac{\ell}{\sqrt{2}}$

$$\text{ar}(\triangle AEF) = \frac{1}{2} x \cdot 2x = x^2 \quad (\because PE = PF = AP = x)$$



Case-II : when $x = AP > OA$, i.e., $x > \frac{\ell}{\sqrt{2}}$ but $x \leq \sqrt{2}\ell$


$$\text{ar(ABEFDA)} = \text{ar(ABCD)} - \text{ar}(\Delta CFE)$$

$$= \ell^2 - \frac{1}{2}(\sqrt{2}\ell - x) \cdot 2(\sqrt{2}\ell - x) \quad [\because CP = \sqrt{2}\ell - x]$$

$$= \ell^2 - (2\ell^2 + x^2 - 2\sqrt{2}\ell x) = 2\sqrt{2}\ell x - x^2 - \ell^2$$

\therefore the required function $s(x)$ is as follows :

$$s(x) = \begin{cases} x^2, & 0 \leq x \leq \frac{\ell}{\sqrt{2}} \\ 2\sqrt{2}\ell x - x^2 - \ell^2, & \frac{\ell}{\sqrt{2}} < x \leq \sqrt{2}\ell \end{cases} ; \text{ area of } s(x) = \begin{cases} \frac{1}{2} & \text{at } x = \frac{1}{\sqrt{2}} \\ 8(\sqrt{2} - 1) & \text{at } x = 2 \end{cases}$$



Ans.

Illustration 36 : If the function $f(x)$ satisfies the functional rule, $f(x + y) = f(x) + f(y) \forall x, y \in \mathbb{R}$ & $f(1) = 5$, then find $\sum_{n=1}^m f(n)$.

Solution :

Here, $f(x + y) = f(x) + f(y)$; put $x = t - 1, y = 1$
 $f(t) = f(t - 1) + f(1)$ (1)

$$\begin{aligned} \therefore f(t) &= f(t - 1) + 5 \\ \Rightarrow f(t) &= \{f(t - 2) + 5\} + 5 \\ \Rightarrow f(t) &= f(t - 2) + 2(5) \\ \Rightarrow f(t) &= f(t - 3) + 3(5) \end{aligned}$$

$$\begin{aligned} &\dots\dots\dots \\ &\dots\dots\dots \\ \Rightarrow f(t) &= f\{t - (t - 1)\} + (t - 1)5 \\ \Rightarrow f(t) &= f(1) + (t - 1)5 \\ \Rightarrow f(t) &= 5 + (t - 1)5 \\ \Rightarrow f(t) &= 5t \end{aligned}$$

$$\therefore \sum_{n=1}^m f(n) = \sum_{n=1}^m (5n) = 5[1 + 2 + 3 + \dots + m] = \frac{5m(m + 1)}{2}$$

$$\text{Hence, } \sum_{n=1}^m f(n) = \frac{5m(m + 1)}{2}.$$

ANSWERS FOR DO YOURSELF

1: (i) (a) $x \in (0, \infty)$ (b) $x \in (-\infty, 0) \cup (4, \infty)$

(ii) (a) $\left[\frac{1}{2}, \infty\right)$ (b) $[-1, 1]$ (c) $[-2, 2]$ (d) $[-1, 1]$



2: (i) (A) $\rightarrow (p, q, r, s)$, (B) $\rightarrow (p)$, (C) $\rightarrow (t)$, (D) $\rightarrow (r)$

3: (i) (a) $(-1, 0) \cup (0, 1]$ (b) $\left(-\infty, -\frac{1}{2}\right] \cup (0, 1) \cup (2, \infty)$

(ii) (a) $(-\infty, \infty)$ (b) $\left[\frac{1}{3}, 1\right]$

4: (i) (a) no (b) no (c) yes

5: (i) $\left[0, \frac{1}{2}\right]$ (ii) A (iii) (a) $y = \frac{1}{x}$ (b) $y = \frac{1}{x^2}$



7: (i) not onto (ii) yes (iii) B

8: (i) (a) 0 (b) 0 (c) 0 (d) 0 (e) 0 (f) 0 (ii) $\begin{cases} x+2, & 0 \leq x < 1 \\ x+1, & 1 \leq x < 2 \\ x, & 2 \leq x < 3 \end{cases}$

9: (i) $f^{-1}(x) = \begin{cases} -\sqrt{-x}, & -1 \leq x \leq 0 \\ \sqrt{x}, & 0 \leq x \leq 1 \end{cases}$ (ii) $y = -2 + e^{x-1}$

10: (i) (a) even (b) odd (c) odd (d) neither even nor odd

11: (i) (a) 2π (b) 1 (c) π

EXERCISE (O-1)

Subjective Type Questions

1. Find the domain of definition of the given functions :

(i) $y = \sqrt{-px} \ (p > 0)$ FN0001

(ii) $y = \frac{1}{x^2 + 1}$ FN0002

(iii) $y = \frac{1}{x^3 - x}$ FN0003

(iv) $y = \frac{1}{\sqrt{x^2 - 4x}}$ FN0004

(v) $y = \sqrt{x^2 - 4x + 3}$ FN0005

(vi) $y = \frac{x}{\sqrt{x^2 - 3x + 2}}$ FN0006

(vii) $y = \sqrt{1 - |x|}$ FN0007

(viii) $y = \log_x 2$. FN0008

(ix) $y = \frac{1}{\log_{10}(1-x)} + \sqrt{x+2}$ FN0009

(x) $y = \sqrt{x} + \sqrt[3]{\frac{1}{x-2}} - \log_{10}(2x-3)$ FN0010

(xi) $y = \frac{3}{4-x^2} + \log_{10}(x^3 - x)$ FN0011

(xii) $y = \frac{1}{\sqrt{\sin x}} + \sqrt[3]{\sin x}$ FN0012

(xiii) $y = \log_{10}(\sqrt{x-4} + \sqrt{6-x})$ FN0013

(xiv) $y = \log_{10}[1 - \log_{10}(x^2 - 5x + 16)]$ FN0014

2. Find the range of the following functions :

(i) $f(x) = \frac{x-1}{x+2}$

FN0015

(ii) $f(x) = \frac{2}{x}$

FN0016

(iii) $f(x) = \frac{1}{x^2 - x + 1}$

FN0017

(iv) $f(x) = \frac{x^2 - x + 1}{x^2 + x + 1}$

FN0018

(v) $f(x) = e^{(x-1)^2}$

FN0019

(vi) $f(x) = x^3 - x^2 + x + 1$

FN0020

(vii) $f(x) = \log(x^8 + x^4 + x^2 + 1)$

FN0021

(viii) $f(x) = \sin^2 x - 2\sin x + 4$

FN0022

(ix) $f(x) = \sin(\log_2 x)$

FN0023

(x) $f(x) = 2^{x^2} + 1$

FN0024

(xi) $f(x) = \frac{e^{2x} - e^x + 1}{e^{2x} + e^x + 1}$

FN0025

(xii) $f(x) = \frac{1}{8 - 3\sin x}$

FN0026

3. The number of integers lying in the domain of the function $f(x) = \sqrt{\log_{0.5} \left(\frac{5-2x}{x} \right)}$ is -

(A) 3

(B) 2

(C) 1

(D) 0

FN0027

4. The range of the function $f: \mathbb{N} \rightarrow \mathbb{Z}; f(x) = (-1)^{x-1}$, is -(A) $[-1, 1]$ (B) $\{-1, 1\}$ (C) $\{0, 1\}$ (D) $\{0, 1, -1\}$

FN0028

5. The range of the function $f(x) = e^{-x} + e^x$, is -(A) $f(x) \geq 1$ (B) $f(x) \leq 1$ (C) $f(x) \geq 2$ (D) $f(x) \leq 2$

FN0029



6. A function f has domain $[-1, 2]$ and range $[0, 1]$. The domain and range respectively of the function g defined by $g(x) = 1 - f(x + 1)$ is
- (A) $[-1, 1]; [-1, 0]$ (B) $[-2, 1]; [0, 1]$ (C) $[0, 2]; [-1, 0]$ (D) $[1, 3]; [-1, 0]$

FN0030

7. For the function $f(x) = \frac{e^x + 1}{e^x - 1}$, if $n(d)$ denotes the number of integers which are not in its domain and $n(r)$ denotes the number of integers which are not in its range, then $n(d) + n(r)$ is equal to -
- (A) 2 (B) 3 (C) 4 (D) Infinite

FN0031

- 8.** Find a formula for each function graphed

**FN0032**

9. If $[a]$ denotes the greatest integer less than or equal to a and $-1 \leq x < 0$, $0 \leq y < 1$, $1 \leq z < 2$, then
- $$\begin{vmatrix} [x]+1 & [y] & [z] \\ [x] & [y]+1 & [z] \\ [x] & [y] & [z]+1 \end{vmatrix}$$
- is equal to -
- (A) $[x]$ (B) $[y]$ (C) $[z]$ (D) none of these

FN0033

10. The range of the function $f(x) = \operatorname{sgn}\left(\frac{\sin^2 x + 2 \sin x + 4}{\sin^2 x + 2 \sin x + 3}\right)$ is (where $\operatorname{sgn}(\cdot)$ denotes signum function)-
- (A) $\{-1, 0, 1\}$ (B) $\{-1, 0\}$ (C) $\{1\}$ (D) $\{0, 1\}$

FN0034

- 11.** The range of the function $f(x) = \sqrt{4 - x^2} + \sqrt{x^2 - 1}$ is
- (A) $[\sqrt{3}, \sqrt{7}]$ (B) $[\sqrt{3}, \sqrt{5}]$ (C) $[\sqrt{2}, \sqrt{3}]$ (D) $[\sqrt{3}, \sqrt{6}]$

FN0035

- 12.** If $[x]$ and $\{x\}$ denotes the greatest integer function less than or equal to x and fractional part function respectively, then the number of real x , satisfying the equation $(x-2)[x] = \{x\} - 1$, is-
- (A) 0 (B) 1 (C) 2 (D) infinite

FN0036

13. If $2f(x) - 3f\left(\frac{1}{x}\right) = x^2$, x is not equal to zero, then $f(2)$ is equal to-

(A) $-\frac{7}{4}$ (B) $\frac{5}{2}$ (C) -1 (D) none of these

FN0037

14. If $f(x) = \frac{4^x}{4^x + 2}$, then $f(x) + f(1-x)$ is equal to-

(A) 0 (B) -1 (C) 1 (D) 4

FN0038

15. If $x^4 f(x) - \sqrt{1 - \sin 2\pi x} = |f(x)| - 2f(x)$, then $f(-2)$ equals

(A) $\frac{1}{17}$ (B) $\frac{1}{11}$ (C) $\frac{1}{19}$ (D) 0

FN0039

16. The graph of a function f is given.

(a) State the value of $f(-1)$.

(b) For what values of x is $f(x) = 2$

(c) State the domain and range of f .


(d) On what interval is f increasing?

(e) Estimated value of $f(2)$ is -

(A) 2.2 (B) 2.8 (C) 2.5 (D) 3

(f) Estimated value of x such that $f(x) = 0$, is -

(A) -2.5 (B) 0.8 (C) -2.9 (D) 0.3



FN0040

17. Graph the function $F(x) = \begin{cases} 3-x, & x \leq 1 \\ 2x, & x > 1 \end{cases}$

FN0041

18. The graphs of f and g are given.

(a) State the value of $f(-4)$ and $g(3)$


(b) For what value of x is $f(x) = g(x)$?

(c) Estimate the solution of the equation $f(x) = -1$.

(d) On what interval is f decreasing?

(e) State the domain and range of f .


(f) State the domain and range of g .



FN0042


19. Solve the following inequalities using graph of $f(x)$:

- (a) $0 \leq f(x) \leq 1$
 (b) $-1 \leq f(x) \leq 2$
 (c) $2 \leq f(x) \leq 3$
 (d) $f(x) > -1$ & $f(x) < 0$




FN0043

20. Which of the following is the graph of $y = |x - 1| + |x - 3|$?



FN0044

21. Which of the following is the graph of $y = \frac{|\sin x|}{\sin x}$?



FN0045

22. Let $f : \mathbb{R} - \left\{ \frac{-15}{2} \right\} \rightarrow \mathbb{R} - \left\{ \frac{1}{2} \right\}$ be defined by $f(x) = \frac{x+10}{2x+15}$ then $f(x)$ is-

- (A) one-one but not onto
 (B) many one but not-onto
 (C) one-one and onto
 (D) many one and onto

FN0046

23. $f : \mathbb{R} \rightarrow \mathbb{R}$ $f(x) = \frac{2x^2 - 5x + 3}{8x^2 + 9x + 11}$, then f is -

- (A) one-one onto
 (B) many-one onto
 (C) one-one into
 (D) many one into

FN0047

24. If $f : \mathbb{R} \rightarrow \mathbb{R}$ & $f(x) = \frac{\sin([x]\pi)}{x^2 + 2x + 3} + 2x - 1 + \sqrt{x(x-1) + \frac{1}{4}}$ (where $[x]$ denotes integral part of x), then $f(x)$ is -

- (A) one-one but not onto
 (B) one-one & onto
 (C) onto but not one-one
 (D) neither one-one nor onto

FN0048

25. Which of the following function is surjective but not injective

(A) $f: \mathbb{R} \rightarrow \mathbb{R} \quad f(x) = x^4 + 2x^3 - x^2 + 1$

(B) $f: \mathbb{R} \rightarrow \mathbb{R} \quad f(x) = x^3 + x + 1$

(C) $f: \mathbb{R} \rightarrow \mathbb{R}^+ \quad f(x) = \sqrt{1+x^2}$

(D) $f: \mathbb{R} \rightarrow \mathbb{R} \quad f(x) = x^3 + 2x^2 - x + 1$

FN0049

26. If $f: (-\infty, 3] \rightarrow [7, \infty)$; $f(x) = x^2 - 6x + 16$, then which of the following is true -

(A) $f^{-1}(x) = 3 + \sqrt{x-7}$

(B) $f^{-1}(x) = 3 - \sqrt{x-7}$

(C) $f^{-1}(x) = \frac{1}{x^2 - 6x + 16}$

(D) f is many-one

FN0050

27. $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = \ln(x + \sqrt{x^2 + 1})$. Another function $g(x)$ is defined such that $g \circ f(x) = x \quad \forall x \in \mathbb{R}$. Then $g(2)$ is -

(A) $\frac{e^2 + e^{-2}}{2}$

(B) e^2

(C) $\frac{e^2 - e^{-2}}{2}$

(D) e^{-2}

FN0051

28. If $f(x) = x|x|$ then $f^{-1}(x)$ equals-

(A) $\sqrt{|x|}$

(B) $(\operatorname{sgn} x) \cdot \sqrt{|x|}$

(C) $-\sqrt{|x|}$

(D) Does not exist

(where $\operatorname{sgn}(x)$ denotes signum function of x)

FN0052

29. Let $P(x) = kx^3 + 2k^2x^2 + k^3$. The sum of all real numbers k for which $(x-2)$ is a factor of $P(x)$, is

(A) 4

(B) 8

(C) -4

(D) -8

FN0053

30. Period of function $f(x) = \min\{\sin x, |x|\} + \frac{x}{\pi} - \left[\frac{x}{\pi} \right]$ (where $[.]$ denotes greatest integer function) is -

(A) $\pi/2$

(B) π

(C) 2π

(D) 4π

FN0054

Multiple Correct Answer

31. Which of the following function(s) have the same domain and range ?

(A) $f(x) = \sqrt{1-x^2}$

(B) $g(x) = \frac{1}{x}$

(C) $h(x) = \sqrt{x}$

(D) $l(x) = \sqrt{4-x}$

FN0055

32. Which of the following pair(s) of function have same graphs?

(A) $f(x) = \frac{\sec x}{\cos x} - \frac{\tan x}{\cot x}$, $g(x) = \frac{\cos x}{\sec x} + \frac{\sin x}{\operatorname{cosec} x}$

(B) $f(x) = \operatorname{sgn}(x^2 - 4x + 5)$, $g(x) = \operatorname{sgn}\left(\cos^2 x + \sin^2\left(x + \frac{\pi}{3}\right)\right)$ where sgn denotes signum function.

(C) $f(x) = e^{\ln(x^2 + 3x + 3)}$, $g(x) = x^2 + 3x + 3$

(D) $f(x) = \frac{\sin x}{\sec x} + \frac{\cos x}{\operatorname{cosec} x}$, $g(x) = \frac{2\cos^2 x}{\cot x}$

FN0056

33. Let $f(x) = \sin^6 x + \cos^6 x$, then -

(A) $f(x) \in [0, 1] \forall x \in \mathbb{R}$

(B) $f(x) = 0$ has no solution

(C) $f(x) \in \left[\frac{1}{4}, 1\right] \forall x \in \mathbb{R}$

(D) $f(x)$ is an injective function

FN0057

34. Let $f(x) = \begin{cases} x^2 - 3x + 4 & ; x < 3 \\ x + 7 & ; x \geq 3 \end{cases}$ and $g(x) = \begin{cases} x + 6 & ; x < 4 \\ x^2 + x + 2 & ; x \geq 4 \end{cases}$, then which of the following is/are true -

(A) $(f + g)(1) = 9$ (B) $(f - g)(3.5) = 1$ (C) $(f \cdot g)(0) = 24$ (D) $\left(\frac{f}{g}\right)(5) = \frac{8}{3}$

FN0058

35. If a function is defined by an implicit equation $2^{|x|+|y|} + 2^{|x|-|y|} = 2$, then -

(A) Domain of function is singleton

(B) Range of function is singleton

(C) graph of the function intersects the line $y = x$

(D) maximum value of function is 2

FN0059

36. Let $f(x) = x^2 + 3x + 2$, then number of solutions to -

(A) $f(|x|) = 2$ is 1

(B) $f(|x|) = 2$ is 3

(C) $|f(x)| = 0.125$ is 4

(D) $|f(|x|)| = 0.125$ is 8

FN0060

37. For each real x , let $f(x) = \max\{x, x^2, x^3, x^4\}$, then $f(x)$ is -

(A) x^4 for $x \leq -1$

(B) x^2 for $-1 < x \leq 0$

(C) $f\left(\frac{1}{2}\right) = \frac{1}{2}$

(D) $f\left(\frac{1}{2}\right) = \frac{1}{4}$

FN0061

Matrix Match Type

38. Match the functions given in column-I correctly with mappings given in column-II.

Column-I

(A) $f: \left[-\frac{1}{2}, \frac{1}{2}\right] \rightarrow \left[\frac{4}{7}, \frac{4}{3}\right]$

$$f(x) = \frac{1}{x^2 + x + 1}$$

(B) $f: [-2, 2] \rightarrow [-1, 1]$

$$f(x) = \sin x$$

(C) $f: \mathbb{R} - \mathbb{I} \rightarrow \mathbb{R}$

$$f(x) = \ell n\{x\}, \text{ (where } \{.\} \text{ represents fractional part function)}$$

(D) $f: (-\infty, 0] \rightarrow [1, \infty), f(x) = (1 + \sqrt{-x}) + (\sqrt{-x} - x)$

Column-II

(P) Injective mapping

(Q) Non-injective mapping

(R) Surjective mapping

(S) Non-surjective mapping

(T) Bijective mapping

FN0062

Linked Comprehension Type**Paragraph for Question 39 & 40**

$$\text{Let } f(x) = \begin{cases} x & ; x < 0 \\ 1-x & ; x \geq 0 \end{cases} \quad \& \quad g(x) = \begin{cases} x^2 & ; x < -1 \\ 2x+3 & ; -1 \leq x \leq 1 \\ x & ; x > 1 \end{cases}$$

On the basis of above information, answer the following questions :

39. Range of $f(x)$ is -

(A) $(-\infty, 1]$

(B) $(-\infty, \infty)$

(C) $(-\infty, 0]$

(D) $(-\infty, 2]$

FN0063

40. Range of $g(f(x))$ is -

(A) $(-\infty, \infty)$

(B) $[1, 3) \cup (3, \infty)$

(C) $[1, \infty)$

(D) $[0, \infty)$

FN0063

EXERCISE (O-2)**Straight Objective Type**

1. Range of function $f(x) = \log_2 \left(\frac{4}{\sqrt{x+2} + \sqrt{2-x}} \right)$ is given by

(A) $(0, \infty)$

(B) $\left[\frac{1}{2}, 1\right]$

(C) $[1, 2]$

(D) $\left[\frac{1}{4}, 1\right]$

FN0064

2. Range of $f(x) = \frac{\sec x + \tan x - 1}{\tan x - \sec x + 1}$; $x \in \left(0, \frac{\pi}{2}\right)$ is-

(A) $(0, 1)$

(B) $(1, \infty)$

(C) $(-1, 0)$

(D) $(-\infty, -1)$

FN0065

3. If $f(x, y) = \max(x, y) + \min(x, y)$ and $g(x, y) = \max(x, y) - \min(x, y)$, then the value of

$$f\left(g\left(-\frac{2}{3}, -\frac{3}{2}\right), g(-3, -4)\right) \text{ is greater than -}$$

- (A) 1 (B) 2 (C) 3 (D) 4

FN0066

4. The number of integral values of x satisfying the inequality $[x-5][x-3] + 2 < [x-5] + 2[x-3]$ (where $[.]$ represents greatest integer function) is -

- (A) 0 (B) 1 (C) 2 (D) 3

FN0067

5. Suppose, $f(x, n) = \sum_{k=1}^n \log_x \left(\frac{k}{x}\right)$, then the value of x satisfying the equation $f(x, 10) = f(x, 11)$, is

- (A) 9 (B) 10 (C) 11 (D) none

FN0068

6. A lion moves in the region given by the graph $y - |y| - x + |x| = 0$. Then on which of the following curve a person can move so that he does not encounter lion -

- (A) $y = e^{-|x|}$ (B) $y = \frac{1}{x}$ (C) $y = \text{signum}(x)$ (D) $y = -|4 + |x||$

FN0069

7. Let $f(x) = \frac{x}{1-x}$ and let α be a real number. If $x_0 = \alpha$, $x_1 = f(x_0)$, $x_2 = f(x_1)$, & $x_{2011} = -\frac{1}{2012}$ then the value of α is -

- (A) $\frac{2011}{2012}$ (B) 1 (C) 2011 (D) -1

FN0070

8. If $f_1(x) = 2^{f_2(x)}$, where $f_2(x) = 2012^{f_3(x)}$, where $f_3(x) = \left(\frac{1}{2013}\right)^{f_4(x)}$, where $f_4(x) = \log_{2013} \log_x 2012$, then the range of $f_1(x)$ is -

- (A) $(2, \infty)$ (B) $(2012, \infty)$ (C) $(0, \infty)$ (D) $(-\infty, \infty)$

FN0071

9. If functions $f(x)$ and $g(x)$ are defined on $\mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = \begin{cases} x+3 & , x \in \text{rational} \\ 4x & , x \in \text{irrational} \end{cases}$,

$$g(x) = \begin{cases} x + \sqrt{5} & , x \in \text{irrational} \\ -x & , x \in \text{rational} \end{cases} \text{ then } (f-g)(x) \text{ is-}$$

- (A) one-one & onto (B) neither one-one nor onto
(C) one-one but not onto (D) onto but not one-one

FN0072

10. Let $f: A \rightarrow B$ be an onto function such that $f(x) = \sqrt{x-2-2\sqrt{x-3}} - \sqrt{x-2+2\sqrt{x-3}}$, then set 'B' is-

(A) $[-2,0]$ (B) $[0,2]$ (C) $[-3,0]$ (D) $[-1,0]$

FN0073

11. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $f(x) = x^3 + ax^2 + bx - 8$. If $f(x) = 0$ has three real roots & $f(x)$ is a bijective function, then $(a+b)$ is equal to

(A) 0 (B) 6 (C) -6 (D) 12

FN0074

12. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \ln(x + \sqrt{x^2 + 1})$, then number of solutions of $|f^{-1}(x)| = e^{-|x|}$ is :-

(A) 1 (B) 2 (C) 3 (D) Infinite

FN0075

13. $f(x) = [x-1] + \{x\}^{[x]}$, $x \in (1,3)$, then $f^{-1}(x)$ is -

(where $[.]$ denotes greatest integer function and $\{.\}$ denotes fractional part function)

(A) $\begin{cases} x+1 & x \in (1,2) \\ 2+\sqrt{x-1} & x \in [2,3) \end{cases}$ (B) $\begin{cases} x-1 & x \in (1,2) \\ 2-\sqrt{x-1} & x \in [2,3) \end{cases}$

(C) $\begin{cases} x-1 & x \in (0,1) \\ 2-\sqrt{x-1} & x \in [1,2) \end{cases}$ (D) $\begin{cases} x+1 & x \in (0,1) \\ 2+\sqrt{x-1} & x \in [1,2) \end{cases}$

FN0076

14. Which of the following functions is an odd function ?

(A) $|x-2| + (x+2) \operatorname{sgn}(x+2)$ (B) $\frac{1}{x(e^x-1)} + \frac{1}{2x}$

(C) $\log(\sin x + \sqrt{1+\sin^2 x})$ (D) $e^{-4x}(e^{2x}-1)^4$

(where $\operatorname{sgn}(x)$ denotes signum function of x)

FN0077

15. Period of $f(x) = \{x\} + \left\{x + \frac{1}{3}\right\} + \left\{x + \frac{2}{3}\right\}$ is equal to (where $\{.\}$ denotes fractional part function)

(A) 1 (B) $\frac{2}{3}$ (C) $\frac{1}{2}$ (D) $\frac{1}{3}$

FN0078

16. Let $f(x) = 2x - \left\{\frac{x}{\pi}\right\}$ and $g(x) = \cos x$, where $\{.\}$ denotes fractional part function, then period of $g \circ f(x)$ is -

(A) $\frac{\pi}{2}$ (B) π (C) $\frac{3\pi}{2}$ (D) $\frac{\pi}{4}$

FN0079

17. The period of the function $\frac{\sin x + \sin 5x}{\cos x + \cos 5x}$ is -

- (A) $\frac{\pi}{3}$ (B) $\frac{\pi}{2}$ (C) π (D) 2π

FN0080

18. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a real valued function such that $f(10+x) = f(10-x) \forall x \in \mathbb{R}$ and $f(20+x) = -f(20-x) \forall x \in \mathbb{R}$. Then which of the following statements is true -

- (A) $f(x)$ is odd and periodic (B) $f(x)$ is odd and aperiodic
(C) $f(x)$ is even and periodic (D) $f(x)$ is even and aperiodic

FN0081

Multiple Correct Answer Type

19. The range of the function $f(\theta) = \sqrt{8\sin^2 \theta + 4\cos^2 \theta - 8\sin \theta \cos \theta}$ is -

- (A) $[\sqrt{5}-1, \sqrt{5}+1]$ (B) $[0, \sqrt{5}+1]$
(C) $[\sqrt{6-\sqrt{20}}, \sqrt{6+\sqrt{20}}]$ (D) none of these

FN0082

20. For the function $f(x) = |x+3| - |x+1| - |x-1| + |x-3|$, identify correct option(s)

- (A) Range of $f(x)$ is $(-\infty, 4]$ (B) maximum value of $f(x)$ is 4
(C) $f(x) = 4$ has infinite solutions (D) $f(x) = 0$ has infinite solutions

FN0083

21. The values of x in $[-2\pi, 2\pi]$, for which the graph of the function $y = \sqrt{\frac{1+\sin x}{1-\sin x}} - \sec x$ and

$y = -\sqrt{\frac{1-\sin x}{1+\sin x}} + \sec x$, coincide are

- (A) $\left[-2\pi, -\frac{3\pi}{2}\right) \cup \left(\frac{3\pi}{2}, 2\pi\right]$ (B) $\left(-\frac{3\pi}{2}, -\frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$
(C) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (D) $[-2\pi, 2\pi] - \left\{\pm\frac{\pi}{2}, \pm\frac{3\pi}{2}\right\}$

FN0084

22. Which of the following statement(s) is(are) correct ?

- (A) If f is a one-one mapping from set A to A , then f is onto.
(B) If f is an onto mapping from set A to A , then f is one-one
(C) Let f and g be two functions defined from $\mathbb{R} \rightarrow \mathbb{R}$ such that $g \circ f$ is injective, then f must be injective.
(D) If set A contains 3 elements while set B contains 2 elements, then total number of functions from A to B is 8.

FN0085

23. A parabola of the form $y = ax^2 + bx + c$ with $a > 0$ intersects the graph of $f(x) = \frac{1}{x^2 - 4}$. Number of possible distinct intersection(s) of these graph is
 (A) 0 (B) 2 (C) 3 (D) 4

FN0086

24. Let $f(x) = \begin{cases} x^2 & ; 0 < x < 2 \\ 2x - 3 & ; 2 \leq x < 3 \\ x + 2 & ; x \geq 3 \end{cases}$ Then :-

- (A) $f\left\{f\left(f\left(\frac{3}{2}\right)\right)\right\} = f\left(\frac{3}{2}\right)$ (B) $1 + f\left\{f\left(f\left(\frac{5}{2}\right)\right)\right\} = f\left(\frac{5}{2}\right)$
 (C) $f\{f(1)\} = f(1) = 1$ (D) none of these

FN0087

25. If $\{x\} = \frac{2}{3}$ & $\left[x + \left\{x + \left[x + \{x + \dots \dots 100 \text{ times}\}\right]\right\}\right] = 5$, then -

- (A) $x = \frac{14}{3}$ (B) $[x] = 5$ (C) $x = \frac{17}{3}$ (D) $[x] = 4$

(where $[.]$ & $\{.\}$ denotes greatest integer function & fractional part function respectively)

FN0088

26. Given $f(1) = 2$ and $f(n+1) = \frac{f(n)-1}{f(n)+1} \forall n \in \mathbb{N}$, then-

- (A) $f(2015) = -\frac{1}{2}$ (B) $(f(2012))^{f(2013)} = 9$
 (C) $f(1001) = 2$ (D) $f(2015) = -3$

FN0089

27. Which of the following is/are true?

- (A) $f(x) = e^x$ and $g(x) = \ln x$, then $f(g(x)) = x$ (wherever $f(g(x))$ is defined)
 (B) $f(x) = \frac{2+x}{3-x}$ and $g(x) = \frac{(3x-2)}{(x+1)}$, then $f(g(x)) = x$ (wherever $f(g(x))$ is defined)
 (C) $f(x) = 4x + 7$ and $g(x) = \frac{(x-7)}{4}$, then $f(g(x)) = x$ (wherever $f(g(x))$ is defined)
 (D) $f(x) = x^3 + 1$ and $g(x) = (x-1)^{1/3}$, then $f(g(x)) = x$ (wherever $f(g(x))$ is defined)

FN0090

28. If $g(x) = x^2 - x + 1$ and $f(x) = \sqrt{\frac{1}{x} - x}$, then -

- (A) Domain of $f(g(x))$ is $[0, 1]$ (B) Range of $f(g(x))$ is $\left(0, \frac{7}{2\sqrt{3}}\right]$
 (C) $f(g(x))$ is many-one function (D) $f(g(x))$ is unbounded function

FN0091

29. Let $f: \{1,2,3,4,5\} \rightarrow \{1,2,3,4,5\}$ is such that $f(x)$ is a one-one function satisfying following condition $f(x) = x + 1$ **if and only if** x is even (i.e. $f(3) \neq 4$, $f(4) = 5$ etc). Then $f^{-1}(2)$ can be-
- (A) 1 (B) 3 (C) 5 (D) 2

FN0092

30. If $f(x) = ax + b$ and $f(f(f(x))) = 27x + 13$ where a and b are real numbers, then-
- (A) $a + b = 3$ (B) $a + b = 4$ (C) $f'(x) = 3$ (D) $f'(x) = -3$

FN0093

Linked Comprehension Type

Paragraph for Question 31 to 33

$$f(x) = \frac{x^2 + 1}{ax} \quad (a \neq 0); \quad g(x) = 3 \sec x; \quad h(x) = \frac{x + 3}{x - 4}.$$

On the basis of above information, answer the following questions :

31. If range of $f(x)$ and $g(x)$ are equal sets then 'a' is equal to -
- (A) 3 (B) $-2/3$ (C) $3/2$ (D) $-3/2$

FN0094

32. $f(x)$ is one-one if-
- (A) $x \in (0, \infty)$ (B) $x \in (-\infty, 0)$ (C) $x \in (1, \infty)$ (D) $x \in (-\infty, 1) - \{0\}$

FN0094

33. Which the following is always false ?

- (A) $h(x)$ is one-one
 (B) $f(x)$ is one-one if $x > 10$
 (C) $g(x)$ is many-one if $x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, 3\right)$
 (D) The values of k for which $f(x) = k$ has exactly one solution is $k = 2$ or $k = -2$

FN0094

Matrix Match Type

- | 34. | Column-I | Column-II |
|-----|--|-----------------|
| | Number of integers in | |
| (A) | Domain of $f(x) = \ell n \{x\}$ | (P) 0 |
| (B) | Domain of $f(x) = \sqrt{\sec(\sin x)} + \sqrt{\left[x + \frac{1}{x}\right]} + \sqrt{10 - [x]^2}$ | (Q) 2 |
| (C) | Range of $f(x) = x^2 - 2x + 2$, $x \in [0, 2]$ | (R) 3 |
| (D) | Range of $f(x) = \sqrt{25 - [x]^2}$ | (S) less than 3 |
| | | (T) more than 3 |
| | (where $[.]$ and $\{.\}$ denote greatest integer function and fractional part function respectively) | |

FN0095

35. Match the function mentioned in column-I with the respective classification given in column-II. (where $[.]$ and $\{.\}$ denote greatest integer function and fractional part function respectively)

Column-I**Column-II**

- | | |
|---|---------------|
| (A) $f: \mathbb{R} \rightarrow \mathbb{R}^+ \quad f(x) = (e^{[x]})(e^{\{x\}})$ | (P) one-one |
| (B) $f: (-\infty, -2) \cup (0, \infty) \rightarrow \mathbb{R} \quad f(x) = \ln(x^2 + 2x)$ | (Q) many-one |
| (C) $f: [-2, 2] \rightarrow [-1, 1] \quad f(x) = \sin x$ | (R) onto |
| (D) $f: \mathbb{R} \rightarrow \mathbb{R} \quad f(x) = x^3 - 3x^2 + 3x - 7$ | (S) periodic |
| | (T) aperiodic |

FN0096

EXERCISE (S-1)

1. Find the domains of definitions of the following functions :

(Read the symbols $[*]$ and $\{*\}$ as greatest integers and fractional part functions respectively.)

(i) $f(x) = \sqrt{\cos 2x} + \sqrt{16 - x^2}$ FN0097

(ii) $f(x) = \log_7 \log_5 \log_3 \log_2 (2x^3 + 5x^2 - 14x)$ FN0098

(iii) $f(x) = \ln(\sqrt{x^2 - 5x - 24} - x - 2)$ FN0099

(iv) $f(x) = \sqrt{\frac{1 - 5^x}{7^{-x} - 7}}$ FN0100

(v) $y = \log_{10} \sin(x - 3) + \sqrt{16 - x^2}$ FN0101

(vi) $f(x) = \log_{100x} \left(\frac{2 \log_{10} x + 1}{-x} \right)$ FN0102

(vii) $f(x) = \sqrt{x^2 - |x|} + \frac{1}{\sqrt{9 - x^2}}$ FN0103

(viii) $f(x) = \sqrt{(x^2 - 3x - 10) \ln^2(x - 3)}$ FN0104

(ix) $f(x) = \sqrt{(5x - 6 - x^2) \left[\{ \ln \{x\} \} \right]} + \sqrt{(7x - 5 - 2x^2)} + \left(\ln \left(\frac{7}{2} - x \right) \right)^{-1}$ FN0105

(x) $f(x) = \log_{\left[\frac{x+1}{x} \right]} |x^2 - x - 6| + {}^{16-x}C_{2x-1} + {}^{20-3x}P_{2x-5}$ FN0106

2. Find the domain & range of the following functions.

(i) $y = \log_{\sqrt{5}} (\sqrt{2} (\sin x - \cos x) + 3)$ FN0107

(ii) $y = \frac{2x}{1+x^2}$ FN0108

(iii) $f(x) = \frac{x^2 - 3x + 2}{x^2 + x - 6}$ FN0109

(iv) $f(x) = \frac{x}{1+|x|}$ FN0110

(v) $y = \sqrt{2-x} + \sqrt{1+x}$ FN0111

(vi) $f(x) = \frac{\sqrt{x+4}-3}{x-5}$ FN0112

3. (i) Write explicitly, functions of y defined by the following equations and also find the domains of definition of the given implicit functions :

(a) $10^x + 10^y = 10$

(b) $x + |y| = 2y$

FN0113

- (ii) The function $f(x)$ is defined on the interval $[0, 1]$. Find the domain of definition of the functions.

(a) $f(\sin x)$

(b) $f(2x + 3)$

FN0114

- (iii) Given that $y = f(x)$ is a function whose domain is $[4, 7]$ and range is $[-1, 9]$. Find the range and domain of

(a) $g(x) = \frac{1}{3}f(x)$

(b) $h(x) = f(x - 7)$

FN0115

4. The sum of integral values of the elements in the domain of $f(x) = \sqrt{\log_{\frac{1}{2}} |3-x|}$ is

FN0116

5. Number of integers in range of $f(x) = x(x+2)(x+4)(x+6)+7$, $x \in [-4, 2]$ is

FN0117

6. The number of even integral value(s) in the range of the function $f(x) = \frac{\tan^2 x + 8 \tan x + 15}{1 + \tan^2 x}$ is

FN0118

7. (a) Draw graphs of the following function, where $[]$ denotes the greatest integer function.

(i) $f(x) = x + [x]$

(ii) $y = (x)^{[x]}$ where $x = [x] + \{x\}$ & $x > 0$ & $x \leq 3$

(iii) $y = \operatorname{sgn}[x]$

(iv) $\operatorname{sgn}(x - [x])$

FN0119

- (b) Identify the pair(s) of functions which are identical ?

(where $[x]$ denotes greatest integer and $\{x\}$ denotes fractional part function)

(i) $f(x) = \operatorname{sgn}(x^2 - 3x + 4)$ and $g(x) = e^{[x]}$ (ii) $f(x) = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$ and $g(x) = \tan x$

(iii) $f(x) = \ln(1+x) + \ln(1-x)$ and $g(x) = \ln(1-x^2)$ (iv) $f(x) = \frac{\cos x}{1 - \sin x}$ and $g(x) = \frac{1 + \sin x}{\cos x}$

FN0120

8. Solve the following problems from (a) to (d) on functional equation :

- (a) The function $f(x)$ defined on the real numbers has the property that $f(f(x)) \cdot (1+f(x)) = -f(x)$ for all x in the domain of f . If the number 3 is in the domain and range of f , compute the value of $f(3)$.

FN0121

- (b) Suppose f is a real function satisfying $f(x + f(x)) = 4f(x)$ and $f(1) = 4$. Find the value of $f(21)$.

FN0122

- (c) Let f be function defined from $\mathbb{R}^+ \rightarrow \mathbb{R}^+$. If $[f(xy)]^2 = x(f(y))^2$ for all positive numbers x and y and $f(2) = 6$, find the value of $f(50)$.

FN0123

- (d) Let f be a function such that $f(3) = 1$ and $f(3x) = x + f(3x - 3)$ for all x . Then find the value of $f(300)$.

FN0124

9. Classify the following functions $f(x)$ defined in $\mathbb{R} \rightarrow \mathbb{R}$ as injective, surjective, both or none.

(a) $f(x) = \frac{x^2 + 4x + 30}{x^2 - 8x + 18}$

FN0125

(b) $f(x) = x^3 - 6x^2 + 11x - 6$

FN0126

(c) $f(x) = (x^2 + x + 5)(x^2 + x - 3)$

FN0127

10. Suppose $f(x) = \sin x$ and $g(x) = 1 - \sqrt{x}$. Then find the domain and range of the following functions.

- (a) fog (b) gof (c) fof (d) gog

FN0128

11. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is such that $f\left(\frac{1-x}{1+x}\right) = x$ for all $x \neq -1$. Prove the following.

- (a) $f(f(x)) = x$ (b) $f(1/x) = -f(x)$, $x \neq 0$ (c) $f(-x - 2) = -f(x) - 2$.

FN0129

12. (a) Find the formula for the function fogoh, given $f(x) = \frac{x}{x+1}$; $g(x) = x^{10}$ and $h(x) = x + 3$. Find also the domain of this function. Also compute $(fogoh)(-1)$.

FN0130

- (b) If $f(x) = \max(x, 1/x)$ for $x > 0$ where $\max(a, b)$ denotes the greater of the two real numbers a and b . Define the function $g(x) = f(x) f(1/x)$ and plot its graph.

FN0131

13. $f(x) = \begin{cases} 1-x & \text{if } x \leq 0 \\ x^2 & \text{if } x > 0 \end{cases}$ and $g(x) = \begin{cases} -x & \text{if } x < 1 \\ 1-x & \text{if } x \geq 1 \end{cases}$ find $(fog)(x)$ and $(gof)(x)$.

FN0132

14. Compute the inverse of the functions :

(a) $f(x) = \ln(x + \sqrt{x^2 + 1})$ FN0133

(b) $f(x) = 2^{\frac{x}{x-1}}$ FN0134

(c) $y = \frac{10^x - 10^{-x}}{10^x + 10^{-x}}$ FN0135

15. Find the inverse of $f(x) = 2^{\log_{10} x} + 8$ and hence solve the equation $f(x) = f^{-1}(x)$.

FN0136

16. Let $f(x) = \frac{2x-1}{x+3}$. If $f^{-1} = \frac{ax+b}{c-x}$, then $a + b + c$ is

FN0137

17. Let f be a one-one function with domain $\{x, y, z\}$ and range $\{1, 2, 3\}$. It is given that exactly one of the following statements is true and the remaining two are false.

$f(x) = 1$; $f(y) \neq 1$; $f(z) \neq 2$. Determine $f^{-1}(1)$

FN0138

18. Find whether the following functions are even or odd or none :

(a) $f(x) = \log(x + \sqrt{1+x^2})$ FN0139

(b) $f(x) = \frac{x(a^x + 1)}{a^x - 1}$ FN0140

(c) $f(x) = \sin x + \cos x$ FN0141

(d) $f(x) = x \sin^2 x - x^3$ FN0142

(e) $f(x) = \sin x - \cos x$ FN0143

(f) $f(x) = \frac{(1+2^x)^2}{2^x}$ FN0144

(g) $f(x) = \frac{x}{e^x - 1} + \frac{x}{2} + 1$ FN0145

(h) $f(x) = [(x+1)^2]^{1/3} + [(x-1)^2]^{1/3}$ FN0146

19. (a) Function f & g are defined by $f(x) = \sin x$, $x \in \mathbb{R}$; $g(x) = \tan x$, $x \in \mathbb{R} - \left(K + \frac{1}{2}\right)\pi$ where $K \in \mathbb{Z}$. Find (i) periods of $fo g$ & gof (ii) range of the function $fo g$ & gof
FN0147
- (b) Suppose that f is an even, periodic function with period 2, and that $f(x) = x$ for all x in interval $[0, 1]$. Find the value of $f(3.14)$.
FN0148
20. Let $f(x)$ be a periodic function with period 'p' satisfying $f(x) + f(x+3) + f(x+6) + \dots + f(x+42) = \text{constant } \forall x \in \mathbb{R}$, then sum of digits of 'p' is
FN0149
21. Let ' f ' be an even periodic function with period '4' such that $f(x) = 2^x - 1$, $0 \leq x \leq 2$. The number of solutions of the equation $f(x) = 1$ in $[-10, 20]$ are
FN0150
22. If $f(x) = \begin{cases} 0 & x < 1 \\ 2x - 2 & x \geq 1 \end{cases}$; then the number of solutions of the equation $f(f(f(x))) = x$ is
FN0151
23. Let $P(x) = x^4 + ax^3 + bx^2 + cx + d$ be a polynomial such that $P(1) = 1$, $P(2) = 8$, $P(3) = 27$, $P(4) = 64$, then find $P(10)$.
FN0152

24. Column-I

 $f(x)$

(A) $\frac{\cos^2 x + \cos x + 2}{\cos^2 x + \cos x + 1}$

(B) $\left| \frac{(\sqrt{\cos x} - \sqrt{\sin x})(\sqrt{\cos x} + \sqrt{\sin x})}{3(\cos x + \sin x)} \right|$

(C) $\frac{7}{3(x^6 + 2x^4 + 3x^2 + 1)}$

(D) $\log_8(x^2 + 2x + 2)$

Column-II

Range

(P) $\left(0, \frac{7}{3}\right]$


(Q) $\left[\frac{4}{3}, \frac{7}{3}\right]$

(R) $\left[0, \frac{1}{3}\right]$

(S) $[0, \infty)$

FN0153

25. The graph of the function $y = f(x)$ is as follows :




Match the function mentioned in **Column-I** with the respective graph given in **Column-II**.


Column-I

Column-II


(A) $y = |f(x)|$




(B) $y = f(|x|)$



(C) $y = f(-|x|)$



(D) $y = \frac{1}{2} (|f(x)| - f(x))$



FN0154

26. If $f(x) = a \log\left(\frac{1+x}{1-x}\right) + bx^3 + c \sin x + 5$ and $f(\log_3 2) = 4$, then $f\left(\log_3\left(\frac{1}{2}\right)\right)$ is equal to

FN0155

EXERCISE (S-2)

1. Find the domains of definitions of the following functions :

(Read the symbols $[*]$ as greatest integers function)

(i) $f(x) = \sqrt{\log_x (\cos 2\pi x)}$

FN0156

(ii) $f(x) = \frac{[x]}{2x - [x]}$

FN0157

2. (a) Let $P(x) = x^6 + ax^5 + bx^4 + cx^3 + dx^2 + ex + f$ be a polynomial such that $P(1) = 1$; $P(2) = 2$; $P(3) = 3$; $P(4) = 4$; $P(5) = 5$ and $P(6) = 6$, then find the value of $P(7)$.

FN0158

- (b) Let a and b be real numbers and let $f(x) = a \sin x + b \sqrt[3]{x} + 4$, $\forall x \in \mathbb{R}$. If $f(\log_{10}(\log_3 10)) = 5$ then find the value of $f(\log_{10}(\log_3 3))$.

FN0159

3. Suppose $p(x)$ is a polynomial with integer coefficients. The remainder when $p(x)$ is divided by $x - 1$ is 1 and the remainder when $p(x)$ is divided by $x - 4$ is 10. If $r(x)$ is the remainder when $p(x)$ is divided by $(x - 1)(x - 4)$, find the value of $r(2006)$.

FN0160

4. A function f , defined for all $x, y \in \mathbb{R}$ is such that $f(1) = 2$ & $f(x + y) - kxy = f(x) + 2y^2$, where k is some constant. Find $f(x)$ & show that :

$$f(x + y)f\left(\frac{1}{x + y}\right) = k \text{ for } x + y \neq 0.$$

FN0161

5. If $f(x) = -1 + |x - 2|$, $0 \leq x \leq 4$
 $g(x) = 2 - |x|$, $-1 \leq x \leq 3$

Then find $\text{fog}(x)$ & $\text{gof}(x)$. Draw rough sketch of the graphs of $\text{fog}(x)$ & $\text{gof}(x)$.

FN0162

6. Let $f(x) = x^{135} + x^{125} - x^{115} + x^5 + 1$. If $f(x)$ is divided by $x^3 - x$ then the remainder is some function of x say $g(x)$. Find the value of $g(10)$.

FN0163

7. Let $\{x\}$ & $[x]$ denote the fractional and integral part of a real number x respectively. Solve $4\{x\} = x + [x]$.

FN0164

8. Let $f(x) = \frac{9^x}{9^x + 3}$ then find the value of the sum $f\left(\frac{1}{2006}\right) + f\left(\frac{2}{2006}\right) + f\left(\frac{3}{2006}\right) + \dots + f\left(\frac{2005}{2006}\right)$

FN0165

9. Let $f(x) = (x + 1)(x + 2)(x + 3)(x + 4) + 5$ where $x \in [-6, 6]$. If the range of the function is $[a, b]$ where $a, b \in \mathbb{N}$ then find the value of $(a + b)$.

FN0166

10. The set of real values of ' x ' satisfying the equality $\left[\frac{3}{x}\right] + \left[\frac{4}{x}\right] = 5$ (where $[]$ denotes the greatest integer function) belongs to the interval $\left(a, \frac{b}{c}\right]$ where $a, b, c \in \mathbb{N}$ and $\frac{b}{c}$ is in its lowest form. Find the value of $a + b + c + abc$.

FN0167

11. $f(x)$ and $g(x)$ are linear function such that for all x , $f(g(x))$ and $g(f(x))$ are Identity functions. If $f(0) = 4$ and $g(5) = 17$, compute $f(2006)$.

FN0168

Comprehension

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying $f(2 - x) = f(2 + x)$ and $f(20 - x) = f(x)$, $\forall x \in \mathbb{R}$.

On the basis of above information, answer the following questions :

12. If $f(0) = 5$, then minimum possible number of values of x satisfying $f(x) = 5$, for $x \in [0, 170]$ is-
 (A) 21 (B) 12 (C) 11 (D) 22

FN0169

13. Graph of $y = f(x)$ is -

- (A) symmetrical about $x = 18$ (B) symmetrical about $x = 5$
 (C) symmetrical about $x = 8$ (D) symmetrical about $x = 20$

FN0169

14. If $f(2) \neq f(6)$, then

- (A) fundamental period of $f(x)$ is 1 (B) fundamental period of $f(x)$ may be 1
 (C) period of $f(x)$ can't be 1 (D) fundamental period of $f(x)$ is 8

FN0169

15. The function $f(x)$ has the property that for each real number x in its domain, $1/x$ is also in its domain and $f(x) + f(1/x) = x$. Find the largest set of real numbers that can be in the domain of $f(x)$?

FN0170

EXERCISE (JM)

1. The domain of the function $f(x) = \frac{1}{\sqrt{|x| - x}}$ is :- [AIEEE - 2011]
 (1) $(-\infty, 0)$ (2) $(-\infty, \infty) - \{0\}$ (3) $(-\infty, \infty)$ (4) $(0, \infty)$

FN0171

2. Let f be a function defined by $f(x) = (x - 1)^2 + 1$, $(x \geq 1)$ [AIEEE - 2009, 2011]

Statement - 1 : The set $\{x : f(x) = f^{-1}(x)\} = \{1, 2\}$

Statement - 2 : f is bijection and $f^{-1}(x) = 1 + \sqrt{x - 1}$, $x \geq 1$.

- (1) Statement-1 is true, Statement-2 is false.
 (2) Statement-1 is false, Statement-2 is true.
 (3) Statement-1 is true, Statement-2 is true ; Statement-2 is a correct explanation for Statement-1.
 (4) Statement-1 is true, Statement-2 is true ; Statement-2 is not a correct explanation for statement-1.

FN0172

3. If $a \in \mathbb{R}$ and the equation $-3(x - [x])^2 + 2(x - [x]) + a^2 = 0$ (where $[x]$ denotes the greatest integer $\leq x$) has no integral solution, then all possible values of a lie in the interval :

[JEE(Main)-2014]

- (1) $(-1, 0) \cup (0, 1)$ (2) $(1, 2)$ (3) $(-2, -1)$ (4) $(-\infty, -2) \cup (2, \infty)$

FN0173

4. If $f(x) + 2f\left(\frac{1}{x}\right) = 3x$, $x \neq 0$, and

[JEE(Main)-2016]

$S = \{x \in \mathbb{R} : f(x) = f(-x)\}$; then S :

- (1) contains more than two elements. (2) is an empty set.
(3) contains exactly one element (4) contains exactly two elements

FN0174

5. The function $f : \mathbb{R} \rightarrow \left[-\frac{1}{2}, \frac{1}{2}\right]$ defined as $f(x) = \frac{x}{1+x^2}$, is :

[JEE(Main)-2017]

- (1) neither injective nor surjective. (2) invertible.
(3) injective but not surjective. (4) surjective but not injective

FN0175

6. Let $S = \{x \in \mathbb{R} : x \geq 0 \text{ and } 2|\sqrt{x} - 3| + \sqrt{x}(\sqrt{x} - 6) + 6 = 0\}$. Then S : [JEE(Main)-2018]

- (1) contains exactly one element. (2) contains exactly two elements.
(3) contains exactly four elements. (4) is an empty set.

FN0176

7. For $x \in \mathbb{R} - \{0, 1\}$, let $f_1(x) = \frac{1}{x}$, $f_2(x) = 1 - x$ and $f_3(x) = \frac{1}{1-x}$ be three given functions. If a function,

$J(x)$ satisfies $(f_2 \circ f_1)(x) = f_3(x)$ then $J(x)$ is equal to :-

[JEE(Main)-2019]

- (1) $f_3(x)$ (2) $f_1(x)$ (3) $f_2(x)$ (4) $\frac{1}{x} f_3(x)$

FN0177

8. Let \mathbb{N} be the set of natural numbers and two functions f and g be defined as $f, g : \mathbb{N} \rightarrow \mathbb{N}$

such that : $f(n) = \begin{cases} \frac{n+1}{2} & \text{if } n \text{ is odd} \\ \frac{n}{2} & \text{if } n \text{ is even} \end{cases}$ and $g(n) = n - (-1)^n$. The fog is :

[JEE(Main)-2019]

- (1) Both one-one and onto (2) One-one but not onto
(3) Neither one-one nor onto (4) onto but not one-one

FN0178

9. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \frac{x}{1+x^2}$, $x \in \mathbb{R}$. Then the range of f is : [JEE(Main)-2019]

- (1) $(-1, 1) - \{0\}$ (2) $\left[-\frac{1}{2}, \frac{1}{2}\right]$ (3) $\mathbb{R} - \left[-\frac{1}{2}, \frac{1}{2}\right]$ (4) $\mathbb{R} - [-1, 1]$

FN0179

10. The number of functions f from $\{1, 2, 3, \dots, 20\}$ onto $\{1, 2, 3, \dots, 20\}$ such that $f(k)$ is a multiple of 3, whenever k is a multiple of 4, is :- [JEE(Main)-2019]

- (1) $(15)! \times 6!$ (2) $5^6 \times 15$ (3) $5! \times 6!$ (4) $6^5 \times (15)!$

FN0180

11. If $f(x) = \log_e \left(\frac{1-x}{1+x} \right)$, $|x| < 1$, then $f\left(\frac{2x}{1+x^2}\right)$ is equal to : [JEE(Main)-2019]

- (1) $2f(x)$ (2) $2f(x^2)$ (3) $(f(x))^2$ (4) $-2f(x)$

FN0181

12. Let $\sum_{k=1}^{10} f(a+k) = 16(2^{10} - 1)$, where the function f satisfies $f(x+y) = f(x)f(y)$ for all natural numbers

x, y and $f(1) = 2$. then the natural number 'a' is [JEE(Main)-2019]

- (1) 4 (2) 3 (3) 16 (4) 2

FN0182

13. Let $f(x) = x^2$, $x \in \mathbb{R}$. For any $A \subseteq \mathbb{R}$, define $g(A) = \{x \in \mathbb{R}, f(x) \in A\}$. If $S = [0, 4]$, then which one of the following statements is not true ? [JEE(Main)-2019]

- (1) $f(g(S)) \neq f(S)$ (2) $f(g(S)) = S$ (3) $g(f(S)) = g(S)$ (4) $g(f(S)) \neq S$

FN0183

14. The number of real roots of the equation $5 + |2^x - 1| = 2^x (2^x - 2)$ is : [JEE(Main)-2019]

- (1) 2 (2) 3 (3) 4 (4) 1

FN0184

15. For $x \in \left(0, \frac{3}{2}\right)$, let $f(x) = \sqrt{x}$, $g(x) = \tan x$ and $h(x) = \frac{1-x^2}{1+x^2}$. If $\phi(x) = ((h \circ f) \circ g)(x)$, then $\phi = \left(\frac{\pi}{3}\right)$ is equal to : [JEE(Main)-2019]

- (1) $\tan \frac{\pi}{12}$ (2) $\tan \frac{7\pi}{12}$ (3) $\tan \frac{11\pi}{12}$ (4) $\tan \frac{5\pi}{12}$

FN0185

16. For $x \in \mathbb{R}$, let $[x]$ denote the greatest integer $\leq x$, then the sum of the series

$$\left[-\frac{1}{3}\right] + \left[-\frac{1}{3} - \frac{1}{100}\right] + \left[-\frac{1}{3} - \frac{2}{100}\right] + \dots + \left[-\frac{1}{3} - \frac{99}{100}\right] \text{ is} \quad [\text{JEE(Main)-2019}]$$

- (1) -153 (2) -133 (3) -131 (4) -135

FN0186

17. If $g(x) = x^2 + x - 1$ and $(g \circ f)(x) = 4x^2 - 10x + 5$, then $f\left(\frac{5}{4}\right)$ is equal to [JEE(Main)-2020]

- (1) $\frac{3}{2}$ (2) $-\frac{1}{2}$ (3) $-\frac{3}{2}$ (4) $\frac{1}{2}$

FN0187

18. Let $f : (1, 3) \rightarrow \mathbb{R}$ be a function defined by $f(x) = \frac{x[x]}{1+x^2}$, where $[x]$ denotes the greatest integer $\leq x$. Then the range of f is [JEE(Main)-2020]

- (1) $\left(\frac{3}{5}, \frac{4}{5}\right)$ (2) $\left(\frac{2}{5}, \frac{3}{5}\right] \cup \left(\frac{3}{4}, \frac{4}{5}\right)$ (3) $\left(\frac{2}{5}, \frac{4}{5}\right]$ (4) $\left(\frac{2}{5}, \frac{1}{2}\right) \cup \left(\frac{3}{5}, \frac{4}{5}\right]$

FN0188

19. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be such that for all $x \in \mathbb{R}$ ($2^{1+x} + 2^{1-x}$), $f(x)$ and $(3^x + 3^{-x})$ are in A.P., then the minimum value of $f(x)$ is [JEE(Main)-2020]

- (1) 0 (2) 3 (3) 2 (4) 4

FN0189

20. The inverse function of $f(x) = \frac{8^{2x} - 8^{-2x}}{8^{2x} + 8^{-2x}}$, $x \in (-1, 1)$, is [JEE(Main)-2020]

- (1) $\frac{1}{4}(\log_8 e) \log_e \left(\frac{1-x}{1+x}\right)$ (2) $\frac{1}{4} \log_e \left(\frac{1-x}{1+x}\right)$
 (3) $\frac{1}{4}(\log_8 e) \log_e \left(\frac{1+x}{1-x}\right)$ (4) $\frac{1}{4} \log_e \left(\frac{1+x}{1-x}\right)$

FN0190

EXERCISE (JA)

1. If functions $f(x)$ and $g(x)$ are defined on $\mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x) = \begin{cases} 0, & x \in \text{rational} \\ x, & x \in \text{irrational} \end{cases}, \quad g(x) = \begin{cases} 0, & x \in \text{irrational} \\ x, & x \in \text{rational} \end{cases}, \text{ then } (f - g)(x) \text{ is -}$$

- (A) one-one and onto (B) neither one-one nor onto
 (C) one-one but not onto (D) onto but not one-one [JEE 2005 (Scr.)]

FN0191

2. Let $f(x) = x^2$ and $g(x) = \sin x$ for all $x \in \mathbb{R}$. Then the set of all x satisfying $(f \circ g \circ g \circ f)(x) = (g \circ g \circ f)(x)$, where $(f \circ g)(x) = f(g(x))$, is- [JEE 2011, 3, (-1)]

- (A) $\pm\sqrt{n\pi}, n \in \{0, 1, 2, \dots\}$ (B) $\pm\sqrt{n\pi}, n \in \{1, 2, \dots\}$
(C) $\frac{\pi}{2} + 2n\pi, n \in \{\dots, -2, -1, 0, 1, 2, \dots\}$ (D) $2n\pi, n \in \{\dots, -2, -1, 0, 1, 2, \dots\}$

FN0192

3. The function $f : [0, 3] \rightarrow [1, 29]$, defined by $f(x) = 2x^3 - 15x^2 + 36x + 1$, is : [JEE 2012, 3, (-1)]

- (A) one-one and onto (B) onto but not one-one
(C) one-one but not onto (D) neither one-one nor onto

FN0193

4. Let $f : (-1, 1) \rightarrow \mathbb{R}$ be such that $f(\cos 4\theta) = \frac{2}{2 - \sec^2 \theta}$ for $\theta \in \left(0, \frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$. Then the value(s) of

$f\left(\frac{1}{3}\right)$ is (are) -

[JEE 2012, 4]

- (A) $1 - \sqrt{\frac{3}{2}}$ (B) $1 + \sqrt{\frac{3}{2}}$ (C) $1 - \sqrt{\frac{2}{3}}$ (D) $1 + \sqrt{\frac{2}{3}}$

FN0194

5. Let $f : \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}$ be given by $f(x) = (\log(\sec x + \tan x))^3$. Then :-

- (A) $f(x)$ is an odd function (B) $f(x)$ is a one-one function
(C) $f(x)$ is an onto function (D) $f(x)$ is an even function

[JEE(Advanced)-2014, 3]

FN0195

6. If the function $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = |x|(x - \sin x)$, then which of the following statements is TRUE ? [JEE(Advanced)-2020]

- (A) f is one-one, but NOT onto (B) f is onto, but NOT one-one
(C) f is BOTH one-one and onto (D) f is NEITHER one-one NOR onto

7. Let the function $f : [0, 1] \rightarrow \mathbb{R}$ be defined by $f(x) = \frac{4^x}{4^x + 2}$ [JEE(Advanced)-2020]

Then the value of $f\left(\frac{1}{40}\right) + f\left(\frac{2}{40}\right) + f\left(\frac{3}{40}\right) + \dots + f\left(\frac{39}{40}\right) - f\left(\frac{1}{2}\right)$ is _____


ANSWER KEY

FUNCTION

EXERCISE (O-1)

1. (i) $-\infty < x \leq 0$ (ii) $x \in \mathbb{R}$ (iii) $x \in \mathbb{R} - \{-1, 0, 1\}$
 (iv) $-\infty < x < 0$ & $4 < x < \infty$ (v) $-\infty < x \leq 1$ and $3 \leq x < \infty$ (vi) $-\infty < x < 1$ and $2 < x < \infty$
 (vii) $-1 \leq x \leq 1$ (viii) $0 < x < 1$ and $1 < x < \infty$. (ix) $-2 \leq x < 0$ and $0 < x < 1$
 (x) $\frac{3}{2} < x < 2$ and $2 < x < \infty$. (xi) $-1 < x < 0$ and $1 < x < 2$; $2 < x < \infty$
 (xii) $2k\pi < x < (2k+1)\pi$, where k is an integer. (xiii) $4 \leq x \leq 6$ (xiv) $2 < x < 3$
2. (i) $\mathbb{R} - \{1\}$ (ii) $\mathbb{R} - \{0\}$ (iii) $(0, 4/3]$ (iv) $[1/3, 3]$ (v) $[1, \infty)$ (vi) \mathbb{R}
 (vii) $[0, \infty)$ (viii) $[3, 7]$ (ix) $[-1, 1]$ (x) $[2, \infty)$ (xi) $[1/3, 1]$ (xii) $\left[\frac{1}{11}, \frac{1}{5}\right]$
3. C 4. B 5. C 6. B 7. C
8. (a) $y = \begin{cases} x, & 0 \leq x \leq 1 \\ 2-x, & 1 < x \leq 2 \end{cases}$ (b) $y = \begin{cases} 2, & 0 \leq x < 1 \text{ and } 2 \leq x < 3 \\ 0, & 1 \leq x < 2 \text{ and } 3 \leq x \leq 4 \end{cases}$
9. C 10. C 11. D 12. D 13. A 14. C 15. A
16. (a) -2, (b) -3, 1, (c) $[-3, 3]$, $[-2, 3]$, (d) $[-1, 3]$, (e) B, (f) A, D

17.



18. (a) -2, 4, (b) -2, 2, (c) -3, 4, (d) $[0, 4]$ (e) $[-4, 4]$, $[-2, 3]$ (f) $[-4, 3]$, $[0.5, 4]$
19. (a) $\left[\frac{4}{3}, \frac{5}{3}\right] \cup \{3\}$ (b) $[-2, -1) \cup [1, 3]$ (c) $[-3, -2] \cup [-1, 1]$ (d) $\left(\frac{5}{3}, 3\right) - \{2\}$
20. D 21. D 22. C 23. D 24. B 25. D 26. B
27. C 28. B 29. D 30. C 31. B, C 32. A, B, C, D
33. A, B, C 34. A, B, C 35. A, B, C 36. A, C 37. A, B, C
38. $(A) \rightarrow (P, R, T)$; $(B) \rightarrow (Q, R)$; $(C) \rightarrow (Q, S)$; $(D) \rightarrow (P, R, T)$ 39. A 40. C

EXERCISE (O-2)

1. B 2. B 3. A 4. C 5. C 6. D 7. D
8. A 9. B 10. A 11. B 12. B 13. D 14. C
15. D 16. B 17. C 18. A 19. A, C 20. B, C, D 21. A, C
22. C, D 23. B, C, D 24. A, B, C 25. A, D 26. A, B, C 27. A, B, C, D
28. A, C 29. B, C 30. B, C 31. B 32. C 33. C
34. $(A) \rightarrow (P, S)$; $(B) \rightarrow (R)$; $(C) \rightarrow (Q, S)$; $(D) \rightarrow (T)$ 35. $(A) \rightarrow (P, R, T)$; $(B) \rightarrow (Q, R, T)$; $(C) \rightarrow (Q, R, T)$; $(D) \rightarrow (P, R, T)$

EXERCISE (S-1)

1. (i) $\left[-\frac{5\pi}{4}, -\frac{3\pi}{4}\right] \cup \left[-\frac{\pi}{4}, \frac{\pi}{4}\right] \cup \left[\frac{3\pi}{4}, \frac{5\pi}{4}\right]$ (ii) $\left(-4, -\frac{1}{2}\right) \cup (2, \infty)$ (iii) $(-\infty, -3]$
 (iv) $(-\infty, -1) \cup [0, \infty)$ (v) $(3 - 2\pi < x < 3 - \pi) \cup (3 < x \leq 4)$
 (vi) $\left(0, \frac{1}{100}\right) \cup \left(\frac{1}{100}, \frac{1}{\sqrt{10}}\right)$
 (vii) $(-3, -1] \cup \{0\} \cup [1, 3)$ (viii) $\{4\} \cup [5, \infty)$ (ix) $(1, 2) \cup (2, 5/2)$ (x) $x \in \{4, 5\}$
2. (i) $D : x \in \mathbb{R} \quad \mathbb{R} : [0, 2]$
 (ii) $D = \mathbb{R}$; range $[-1, 1]$
 (iii) $D : \{x \mid x \in \mathbb{R}; x \neq -3; x \neq 2\} \quad \mathbb{R} : \{f(x) \mid f(x) \in \mathbb{R}, f(x) \neq 1/5; f(x) \neq 1\}$
 (iv) $D : \mathbb{R}, \mathbb{R} : (-1, 1)$
 (v) $D : -1 \leq x \leq 2 \quad \mathbb{R} : [\sqrt{3}, \sqrt{6}]$
 (vi) $D : [-4, \infty) - \{5\}; \mathbb{R} : \left(0, \frac{1}{6}\right) \cup \left(\frac{1}{6}, \frac{1}{3}\right]$
3. (i) (a) $y = \log(10 - 10^x), -\infty < x < 1$
 (b) $y = x/3$ when $-\infty < x < 0$ & $y = x$ when $0 \leq x < +\infty$
 (ii) (a) $2K\pi \leq x \leq 2K\pi + \pi$ where $K \in \mathbb{I}$ (b) $[-3/2, -1]$
 (iii) (a) Range : $[-1/3, 3]$, Domain = $[4, 7]$; (b) Range $[-1, 9]$ and domain $[11, 14]$
4. 6 5. 401 6. 9 7. (b) (i), (iii) are identical
8. (a) $-3/4$; (b) 64; (c) 30, (d) 5050
9. (a) neither surjective nor injective (b) surjective but not injective
 (c) neither injective nor surjective
10. (a) domain is $x \geq 0$; range $[-1, 1]$; (b) domain $2k\pi \leq x \leq 2k\pi + \pi$; range $[0, 1]$
 (c) domain $x \in \mathbb{R}$; range $[-\sin 1, \sin 1]$ (d) domain is $0 \leq x \leq 1$; range is $[0, 1]$
12. (a) $\frac{(x+3)^{10}}{(x+3)^{10}+1}$, domain is \mathbb{R} , $\frac{1024}{1025}$; (b) $g(x) = \begin{cases} \frac{1}{x^2} & \text{if } 0 < x \leq 1 \\ x^2 & \text{if } x > 1 \end{cases}$
13. $(\text{gof})(x) = \begin{cases} x & \text{if } x \leq 0 \\ -x^2 & \text{if } 0 < x < 1 \\ 1-x^2 & \text{if } x \geq 1 \end{cases}$; $(\text{fog})(x) = \begin{cases} x^2 & \text{if } x < 0 \\ 1+x & \text{if } 0 \leq x < 1 \\ x & \text{if } x \geq 1 \end{cases}$
14. (a) $\frac{e^x - e^{-x}}{2}$; (b) $\frac{\log_2 x}{\log_2 x - 1}$; (c) $\frac{1}{2} \log \frac{1+x}{1-x}$
15. $x = 10$; $f^{-1}(x) = 10^{\log_2(x-8)}$ 16. 6 17. $f^{-1}(1) = y$
18. (a) odd, (b) even, (c) neither odd nor even, (d) odd, (e) neither odd nor even,
 (f) even, (g) even, (h) even

19. (a) (i) Period of fog is π , period of gof is 2π (ii) range of fog is $[-1, 1]$, range of gof is $[-\tan 1, \tan 1]$
 (b) 0.86
 20. 9 21. 15 22. 2 23. 4024
 24. (A) \rightarrow (Q); (B) \rightarrow (R); (C) \rightarrow (P); (D) \rightarrow (S) 25. (A) S; (B) R; (C) P; (D) Q 26. 6

EXERCISE (S-2)

1. (i) $(0, 1/4) \cup (3/4, 1) \cup \{x : x \in \mathbb{N}, x \geq 2\}$ (ii) $\mathbb{R} - \left\{-\frac{1}{2}, 0\right\}$
 2. (a) 727, (b) 3 3. 6016 4. $f(x) = 2x^2$
 5. $\text{fog}(x) = \begin{cases} -(1+x), & -1 \leq x \leq 0 \\ x-1, & 0 < x \leq 2 \end{cases}; \text{gof}(x) = \begin{cases} x+1, & 0 \leq x < 1 \\ 3-x, & 1 \leq x \leq 2 \\ x-1, & 2 < x \leq 3 \\ 5-x, & 3 < x \leq 4 \end{cases}$
 $\text{fof}(x) = \begin{cases} x, & 0 \leq x \leq 1 \\ 4-x, & 1 < x \leq 2 \end{cases}; \text{gog}(x) = \begin{cases} -x, & -1 \leq x \leq 0 \\ x, & 0 < x \leq 2 \\ 4-x, & 2 < x \leq 3 \end{cases}$
 6. 21
 7. $x = 0$ या $5/3$ 8. 1002.5 9. 5049 10. 20
 11. 122 12. D 13. A 14. C 15. $\{-1, 1\}$

EXERCISE (JM)

1. 1 2. 3 3. 1 4. 4 5. 4 6. 2 7. 1 8. 4
 9. 2 10. 1 11. 1 12. 2 13. 3 14. 4 15. 3 16. 2
 17. 2 18. 4 19. 2 20. 3

EXERCISE (JA)

1. A 2. A 3. B
 4. (zero marks to all) Reason : (as we are getting two f images for one x in the domain) 5. A,B,C
 6. C 7. 19.00

INVERSE TRIGONOMETRIC FUNCTION

1. INTRODUCTION :

The inverse trigonometric functions, denoted by $\sin^{-1}x$ or $(\arcsin x)$, $\cos^{-1}x$ etc., denote the angles whose sine, cosine etc, is equal to x . The angles are usually the numerically smallest angles, except in the case of $\cot^{-1}x$ and if positive & negative angles have same numerical value, the positive angle has been chosen.

It is worthwhile noting that the functions $\sin x$, $\cos x$ etc are in general not invertible. Their inverse is defined by choosing an appropriate domain & co-domain so that they become invertible. For this reason the chosen value is usually the simplest and easy to remember.

2. DOMAIN & RANGE OF INVERSE TRIGONOMETRIC FUNCTIONS :

S.No	$f(x)$	Domain	Range
(1)	$\sin^{-1}x$	$ x \leq 1$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
(2)	$\cos^{-1}x$	$ x \leq 1$	$[0, \pi]$
(3)	$\tan^{-1}x$	$x \in \mathbb{R}$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
(4)	$\sec^{-1}x$	$ x \geq 1$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$ or $\left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$
(5)	$\operatorname{cosec}^{-1}x$	$ x \geq 1$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
(6)	$\cot^{-1}x$	$x \in \mathbb{R}$	$(0, \pi)$


3. GRAPH OF INVERSE TRIGONOMETRIC FUNCTIONS :

(a) $f: \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1]$


$f(x) = \sin x$

$f^{-1}: [-1, 1] \rightarrow [-\pi/2, \pi/2]$

$f^{-1}(x) = \sin^{-1}(x)$




(Taking image of $\sin x$ about $y = x$ to get $\sin^{-1}x$)



($y = \sin^{-1}x$)

(b) $f : [0, \pi] \rightarrow [-1, 1]$


$f(x) = \cos x$



(Taking image of $\cos x$ about $y = x$)

(c) $f : (-\pi/2, \pi/2) \rightarrow \mathbb{R}$


$f(x) = \tan x$



(Taking image of $\tan x$ about $y = x$)

(d) $f : (0, \pi) \rightarrow \mathbb{R}$

$f(x) = \cot x$



(Taking image of $\cot x$ about $y = x$)

(e) $f : [0, \pi/2) \cup (\pi/2, \pi] \rightarrow (-\infty, -1] \cup [1, \infty)$


$f(x) = \sec x$

$f^{-1} : (-\infty, -1] \cup [1, \infty) \rightarrow [0, \pi/2) \cup (\pi/2, \pi]$

$f^{-1}(x) = \sec^{-1} x$

$f^{-1} : [-1, 1] \rightarrow [0, \pi]$


$f^{-1}(x) = \cos^{-1} x$



$(y = \cos^{-1} x)$

$f^{-1} : \mathbb{R} \rightarrow (-\pi/2, \pi/2)$


$f^{-1}(x) = \tan^{-1} x$




$(y = \tan^{-1} x)$

$f^{-1} : \mathbb{R} \rightarrow (0, \pi)$

$f^{-1}(x) = \cot^{-1} x$



$(y = \cot^{-1} x)$




(f) $f : [-\pi/2, 0) \cup (0, \pi/2] \rightarrow (-\infty, -1] \cup [1, \infty)$

$f(x) = \operatorname{cosec} x$

$f^{-1} : (-\infty, -1] \cup [1, \infty) \rightarrow [-\pi/2, 0) \cup (0, \pi/2]$

$f^{-1}(x) = \operatorname{cosec}^{-1} x$



From the above discussions following IMPORTANT points can be concluded :

- All the inverse trigonometric functions represent an angle.
- If $x > 0$, then all six inverse trigonometric functions viz $\sin^{-1} x$, $\cos^{-1} x$, $\tan^{-1} x$, $\sec^{-1} x$, $\operatorname{cosec}^{-1} x$, $\cot^{-1} x$ represent an acute angle.
- If $x < 0$, then $\sin^{-1} x$, $\tan^{-1} x$ & $\operatorname{cosec}^{-1} x$ represent an angle from $-\pi/2$ to 0 (IVth quadrant)
- If $x < 0$, then $\cos^{-1} x$, $\cot^{-1} x$ & $\sec^{-1} x$ represent an obtuse angle. (IInd quadrant)
- IIIrd quadrant is never used in range of inverse trigonometric function.

Illustration 1 : The value of $\tan^{-1}(1) + \cos^{-1}\left(-\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{2}\right)$ is equal to

- (A) $\frac{\pi}{4}$ (B) $\frac{5\pi}{12}$ (C) $\frac{3\pi}{4}$ (D) $\frac{13\pi}{12}$

Solution : $\tan^{-1}(1) + \cos^{-1}\left(-\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{2}\right) = \frac{\pi}{4} + \frac{2\pi}{3} - \frac{\pi}{6} = \frac{\pi}{4} + \frac{\pi}{2} = \frac{3\pi}{4}$ **Ans.(C)**

Illustration 2 : If $\sum_{i=1}^{2n} \cos^{-1} x_i = 0$ then find the value of $\sum_{i=1}^{2n} x_i$

Solution : We know, $0 \leq \cos^{-1} x \leq \pi$
Hence, each value $\cos^{-1} x_1, \cos^{-1} x_2, \cos^{-1} x_3, \dots, \cos^{-1} x_{2n}$ are non-negative their sum is zero only when each value is zero.

i.e., $\cos^{-1} x_i = 0$ for all i

$\Rightarrow x_i = 1$ for all i

$\therefore \sum_{i=1}^{2n} x_i = x_1 + x_2 + x_3 + \dots + x_{2n} = \underbrace{\{1 + 1 + 1 \dots + 1\}}_{2n \text{ times}} = 2n$ {using (i)}

$\Rightarrow \sum_{i=1}^{2n} x_i = 2n$ **Ans.**


Do yourself - 1 :

- (i) If α, β are roots of the equation $6x^2 + 11x + 3 = 0$, then
 (A) both $\cos^{-1}\alpha$ and $\cos^{-1}\beta$ are real (B) both $\operatorname{cosec}^{-1}\alpha$ and $\operatorname{cosec}^{-1}\beta$ are real
 (C) both $\cot^{-1}\alpha$ and $\cot^{-1}\beta$ are real (D) none of these
- (ii) If $\sin^{-1}x + \sin^{-1}y = \pi$ and $x = ky$, then find the value of $39^{2k} + 5^k$.

3. PROPERTIES OF INVERSE CIRCULAR FUNCTIONS :


P-1 (i) $y = \sin(\sin^{-1}x) = x$

$$x \in [-1, 1], y \in [-1, 1]$$




(ii) $y = \cos(\cos^{-1}x) = x$

$$x \in [-1, 1], y \in [-1, 1]$$




(iii) $y = \tan(\tan^{-1}x) = x$

$$x \in \mathbb{R}, y \in \mathbb{R}$$




(iv) $y = \cot(\cot^{-1}x) = x,$

$$x \in \mathbb{R}; y \in \mathbb{R}$$




(v) $y = \operatorname{cosec}(\operatorname{cosec}^{-1}x) = x,$

$$|x| \geq 1, |y| \geq 1$$



(vi) $y = \sec(\sec^{-1} x) = x$

$|x| \geq 1 ; |y| \geq 1$



Note : All the above functions are aperiodic.

Illustration 3 : Evaluate the following :

(i) $\sin(\cos^{-1} 3/5)$ (ii) $\cos(\tan^{-1} 3/4)$ (iii) $\sin\left(\frac{\pi}{2} - \sin^{-1}\left(-\frac{1}{2}\right)\right)$

(iv) $\tan\left\{2 \tan^{-1} \frac{1}{5} - \frac{\pi}{4}\right\}$

Solution :

(i) Let $\cos^{-1} 3/5 = \theta$. Then,
 $\cos \theta = 3/5 \Rightarrow \sin \theta = 4/5$
 $\therefore \sin(\cos^{-1} 3/5) = \sin \theta = 4/5$

(ii) Let $\tan^{-1} 3/4 = \theta$. Then,
 $\tan \theta = 3/4$
 $\Rightarrow \cos \theta = \frac{4}{5}$ $\left\{ \because \text{as } \cos^2 \theta = \frac{1}{1 + \tan^2 \theta} \right\}$
 $\therefore \cos(\tan^{-1} 3/4) = \cos \theta = 4/5$

(iii) $\sin\left(\frac{\pi}{2} - \sin^{-1}\left(-\frac{1}{2}\right)\right) = \sin\left(\frac{\pi}{2} - \left(-\frac{\pi}{6}\right)\right) = \sin \frac{2\pi}{3} = \frac{\sqrt{3}}{2}$

(iv) Let $\tan^{-1}\left(\frac{1}{5}\right) = \theta \Rightarrow \tan \theta = \frac{1}{5}$

$\tan\left(2\theta - \frac{\pi}{4}\right) = \frac{\tan(2\theta) - 1}{1 + \tan 2\theta}$ and $\tan 2\theta = \frac{2 \times \frac{1}{5}}{1 - \frac{1}{25}} = \frac{5}{12}$ ($\because \tan \theta = \frac{1}{5}$)

$\Rightarrow \tan\left(2\theta - \frac{\pi}{4}\right) = \frac{\frac{5}{12} - 1}{1 + \frac{5}{12}} = -\frac{7}{17}$

Ans.

Illustration 4 : Prove that $\sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3) = 15$

Solution :

We have,

$$\sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3)$$

$$= \left\{ \sec(\tan^{-1} 2) \right\}^2 + \left\{ \operatorname{cosec}(\cot^{-1} 3) \right\}^2 = \left\{ \sec\left(\tan^{-1} \frac{2}{1}\right) \right\}^2 + \left\{ \operatorname{cosec}\left(\cot^{-1} \frac{3}{1}\right) \right\}^2$$

$$= \left\{ \sec(\sec^{-1} \sqrt{5}) \right\}^2 + \left\{ \operatorname{cosec}(\operatorname{cosec}^{-1} \sqrt{10}) \right\}^2 = (\sqrt{5})^2 + (\sqrt{10})^2 = 15$$


Do yourself - 2 :

Evaluate the following :

(i) $\tan\left(\cos^{-1}\left(\frac{8}{17}\right)\right)$ (ii) $\sin\left(\frac{1}{2}\cos^{-1}\left(\frac{4}{5}\right)\right)$ (iii) $\sin\left(\frac{\pi}{3} - \sin^{-1}\left(\frac{-1}{2}\right)\right)$ (iv) $\sin\left(\cos^{-1}\frac{3}{5}\right)$


P-2 (i) $y = \sin^{-1}(\sin x)$, $x \in \mathbb{R}$, $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ periodic with period 2π and it is an odd function.

$$\sin^{-1}(\sin x) = \begin{cases} -\pi - x & , -\pi \leq x \leq -\frac{\pi}{2} \\ x & , -\frac{\pi}{2} < x < \frac{\pi}{2} \\ \pi - x & , \frac{\pi}{2} \leq x \leq \pi \end{cases}$$



(ii) $y = \cos^{-1}(\cos x)$, $x \in \mathbb{R}$, $y \in [0, \pi]$, periodic with period 2π and it is an even function.


$$\cos^{-1}(\cos x) = \begin{cases} -x & , -\pi \leq x \leq 0 \\ x & , 0 < x \leq \pi \end{cases}$$




(iii) $y = \tan^{-1}(\tan x)$

$x \in \mathbb{R} - \left\{ (2n-1)\frac{\pi}{2}, n \in \mathbb{I} \right\}$, $y \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, periodic with period π and it is an odd function.


$$\tan^{-1}(\tan x) = \begin{cases} x + \pi & , -\frac{3\pi}{2} < x < -\frac{\pi}{2} \\ x & , -\frac{\pi}{2} < x < \frac{\pi}{2} \\ x - \pi & , \frac{\pi}{2} < x < \frac{3\pi}{2} \end{cases}$$



- (iv) $y = \cot^{-1}(\cot x)$, $x \in \mathbb{R} - \{n\pi, n \in \mathbb{I}\}$, $y \in (0, \pi)$, periodic with period π and neither even nor odd function.



- (v) $y = \operatorname{cosec}^{-1}(\operatorname{cosec} x)$, $x \in \mathbb{R} - \{n\pi, n \in \mathbb{I}\}$, $y \in \left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$, is periodic with period 2π and it is an odd function.



- (vi) $y = \sec^{-1}(\sec x)$, y is periodic with period 2π and it is an even function.

$$x \in \mathbb{R} - \left\{(2n-1)\frac{\pi}{2}, n \in \mathbb{I}\right\}, y \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$$




Illustration 5 : The value of $\sin^{-1}(-\sqrt{3}/2) + \cos^{-1}(\cos(7\pi/6))$ is -

- (A) $5\pi/6$ (B) $\pi/2$ (C) $3\pi/2$ (D) none of these

Solution : $\sin^{-1}(-\sqrt{3}/2) = -\sin^{-1}(\sqrt{3}/2) = -\pi/3$

and $\cos^{-1}(\cos(7\pi/6)) = \cos^{-1}(\cos(2\pi - 5\pi/6)) = \cos^{-1}(\cos(5\pi/6)) = 5\pi/6$

Hence $\sin^{-1}(-\sqrt{3}/2) + \cos^{-1}(\cos(7\pi/6)) = -\frac{\pi}{3} + \frac{5\pi}{6} = \frac{\pi}{2}$

Ans.(B)

Illustration 6 : Evaluate the following :

- (i) $\sin^{-1}(\sin 10)$ (ii) $\tan^{-1}(\tan(-6))$ (iii) $\cot^{-1}(\cot 4)$

Solution : (i) We know that $\sin^{-1}(\sin\theta) = \theta$, if $-\pi/2 \leq \theta \leq \pi/2$

Here, $\theta = 10$ radians which does not lie between $-\pi/2$ and $\pi/2$

But, $3\pi - \theta$ i.e., $3\pi - 10$ lie between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$

Also, $\sin(3\pi - 10) = \sin 10$

$\therefore \sin^{-1}(\sin 10) = \sin^{-1}(\sin(3\pi - 10)) = (3\pi - 10)$

(ii) We know that,

$\tan^{-1}(\tan\theta) = \theta$, if $-\pi/2 < \theta < \pi/2$. Here, $\theta = -6$, radians which does not lie between $-\pi/2$ and $\pi/2$. We find that $2\pi - 6$ lies between $-\pi/2$ and $\pi/2$ such that;

$\tan(2\pi - 6) = -\tan 6 = \tan(-6)$


$\therefore \tan^{-1}(\tan(-6)) = \tan^{-1}(\tan(2\pi - 6)) = (2\pi - 6)$

(iii) $\cot^{-1}(\cot 4) = \cot^{-1}(\cot(\pi + (4 - \pi))) = \cot^{-1}(\cot(4 - \pi)) = (4 - \pi)$

Ans.

Illustration 7 : Find the number of solutions of (x, y) which satisfy $|y| = \cos x$ and $y = \sin^{-1}(\sin x)$, where $|x| \leq 3\pi$.

Solution : Graphs of $y = \sin^{-1}(\sin x)$ and $|y| = \cos x$ meet exactly six times in $[-3\pi, 3\pi]$.



Do yourself - 3 :

Evaluate the following :

(i) $\cos^{-1}\left(\cos\frac{13\pi}{6}\right)$ (ii) $\tan^{-1}\left(\tan\left(\frac{7\pi}{6}\right)\right)$ (iii) $\sin^{-1}\left(\sin\left(\frac{5\pi}{6}\right)\right)$

P-3 (i) $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \quad |x| \leq 1$

(ii) $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2} \quad x \in \mathbb{R}$

(iii) $\operatorname{cosec}^{-1} x + \sec^{-1} x = \frac{\pi}{2} \quad |x| \geq 1$

- P-4** (i) $\sin^{-1}(-x) = -\sin^{-1} x$, $|x| \leq 1$
 (ii) $\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1} x$, $|x| \geq 1$
 (iii) $\tan^{-1}(-x) = -\tan^{-1} x$, $x \in \mathbb{R}$
 (iv) $\cot^{-1}(-x) = \pi - \cot^{-1} x$, $x \in \mathbb{R}$
 (v) $\cos^{-1}(-x) = \pi - \cos^{-1} x$, $|x| \leq 1$
 (vi) $\sec^{-1}(-x) = \pi - \sec^{-1} x$, $|x| \geq 1$

- P-5** (i) $\operatorname{cosec}^{-1} x = \sin^{-1} \frac{1}{x}$; $|x| \geq 1$
 (ii) $\sec^{-1} x = \cos^{-1} \frac{1}{x}$; $|x| \geq 1$
 (iii) $\cot^{-1} x = \begin{cases} \tan^{-1} \frac{1}{x} & ; x > 0 \\ \pi + \tan^{-1} \frac{1}{x} & ; x < 0 \end{cases}$

Illustration 8 : Find value of x if $\cos^{-1}(-x) + \tan^{-1}(-x) - 2\sin^{-1}(x) + \sec^{-1}\left(-\frac{1}{x}\right) = \frac{\pi}{4}$ for $|x| \leq 1$.

Solution :

$$\pi - \cos^{-1}(x) - \tan^{-1}(x) - 2\sin^{-1}(x) + \cos^{-1}(-x) = \frac{\pi}{4}$$

$$\pi - \cos^{-1}(x) - \tan^{-1}(x) - 2\sin^{-1}(x) + \pi - \cos^{-1}(x) = \frac{\pi}{4}$$

$$2\pi - 2(\sin^{-1}x + \cos^{-1}x) - \frac{\pi}{4} = \tan^{-1}x$$

$$2\pi - \pi - \frac{\pi}{4} = \tan^{-1}x \Rightarrow \tan^{-1}x = \frac{3\pi}{4} \text{ Hence no solution}$$

Ans.

Do yourself - 4 :

(i) Prove the following :

(a) $\cos^{-1}\left(\frac{5}{13}\right) = \tan^{-1}\left(\frac{12}{5}\right)$

(b) $\sin^{-1}\left(-\frac{4}{5}\right) = \tan^{-1}\left(-\frac{4}{3}\right) = \cos^{-1}\left(-\frac{3}{5}\right) - \pi$

(ii) Find the value of $\sin(\tan^{-1}a + \tan^{-1}\frac{1}{a})$; $a \neq 0$

P-6 (i) (a) $\tan^{-1} x + \tan^{-1} y = \begin{cases} \tan^{-1} \frac{x+y}{1-xy} & \text{where } x > 0, y > 0 \text{ \& } xy < 1 \\ \pi + \tan^{-1} \frac{x+y}{1-xy} & \text{where } x > 0, y > 0 \text{ \& } xy > 1 \\ \frac{\pi}{2}, & \text{where } x > 0, y > 0 \text{ \& } xy = 1 \end{cases}$

- (b) $\tan^{-1} x - \tan^{-1} y = \tan^{-1} \frac{x-y}{1+xy}$ where $x > 0, y > 0$
- (c) $\tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \tan^{-1} \left[\frac{x+y+z-xyz}{1-xy-yz-zx} \right]$ if $x > 0, y > 0, z > 0$ & $xy + yz + zx < 1$
- (ii) (a) $\sin^{-1} x + \sin^{-1} y = \begin{cases} \sin^{-1}[x\sqrt{1-y^2} + y\sqrt{1-x^2}] & \text{where } x > 0, y > 0 \text{ \& } (x^2 + y^2) \leq 1 \\ \pi - \sin^{-1}[x\sqrt{1-y^2} + y\sqrt{1-x^2}] & \text{where } x > 0, y > 0 \text{ \& } x^2 + y^2 > 1 \end{cases}$
- (b) $\sin^{-1} x - \sin^{-1} y = \sin^{-1} [x\sqrt{1-y^2} - y\sqrt{1-x^2}]$ where $x > 0, y > 0$
- (iii) (a) $\cos^{-1} x + \cos^{-1} y = \cos^{-1} [xy - \sqrt{1-x^2}\sqrt{1-y^2}]$ where $x > 0, y > 0$
- (b) $\cos^{-1} x - \cos^{-1} y = \begin{cases} \cos^{-1}(xy + \sqrt{1-x^2}\sqrt{1-y^2}) & ; \quad x < y, \quad x, y > 0 \\ -\cos^{-1}(xy + \sqrt{1-x^2}\sqrt{1-y^2}) & ; \quad x > y, \quad x, y > 0 \end{cases}$

Note : In the above results x & y are taken positive. In case if these are given as negative, we first apply **P-4** and then use above results.

Illustration 9 : Prove that

$$(i) \quad \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{13} = \tan^{-1} \frac{2}{9} \quad (ii) \quad \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{8} = \frac{\pi}{4}$$

Solution : (i) L.H.S. = $\tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{13}$

$$= \tan^{-1} \left\{ \frac{\frac{1}{7} + \frac{1}{13}}{1 - \frac{1}{7} \times \frac{1}{13}} \right\} \quad \left\{ \because \tan^{-1} x + \tan^{-1} y = \tan^{-1} \left(\frac{x+y}{1-xy} \right); \text{ if } xy < 1 \right\}$$

$$= \tan^{-1} \left(\frac{20}{90} \right) = \tan^{-1} \left(\frac{2}{9} \right) = \text{R.H.S.}$$

(ii) $\left(\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} \right) + \left(\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{8} \right)$

$$= \tan^{-1} \left(\frac{\frac{1}{5} + \frac{1}{7}}{1 - \frac{1}{5} \times \frac{1}{7}} \right) + \tan^{-1} \left(\frac{\frac{1}{3} + \frac{1}{8}}{1 - \frac{1}{3} \times \frac{1}{8}} \right) = \tan^{-1} \left(\frac{6}{17} \right) + \tan^{-1} \left(\frac{11}{23} \right)$$

$$= \tan^{-1} \left(\frac{\frac{6}{17} + \frac{11}{23}}{1 - \frac{6}{17} \times \frac{11}{23}} \right) = \tan^{-1} \left(\frac{325}{325} \right) = \tan^{-1}(1) = \frac{\pi}{4}$$

Ans.

Illustration 10 : Prove that $\sin^{-1} \frac{12}{13} + \cot^{-1} \frac{4}{3} + \tan^{-1} \frac{63}{16} = \pi$

Solution : We have,

$$\begin{aligned} & \sin^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{63}{16} \\ &= \tan^{-1} \frac{12}{5} + \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{63}{16} \quad \left[\because \sin^{-1} \frac{12}{13} = \tan^{-1} \frac{12}{5} \text{ and } \cos^{-1} \frac{4}{5} = \tan^{-1} \frac{3}{4} \right] \\ &= \pi + \tan^{-1} \left\{ \frac{\frac{12}{5} + \frac{3}{4}}{1 - \frac{12}{5} \times \frac{3}{4}} \right\} + \tan^{-1} \frac{63}{16} \quad \left[\because \tan^{-1} x + \tan^{-1} y = \pi + \tan^{-1} \left(\frac{x+y}{1-xy} \right), \text{ if } xy > 1 \right] \\ &= \pi + \tan^{-1} \left(\frac{63}{-16} \right) + \tan^{-1} \left(\frac{63}{16} \right) \\ &= \pi - \tan^{-1} \frac{63}{16} + \tan^{-1} \frac{63}{16} \quad \left[\because \tan^{-1}(-x) = -\tan^{-1} x \right] \\ &= \pi \end{aligned}$$

Illustration 11 : Prove that : $\cos^{-1} \frac{12}{13} + \sin^{-1} \frac{3}{5} = \sin^{-1} \frac{56}{65}$

Solution : We have, L.H.S. = $\cos^{-1} \left(\frac{12}{13} \right) + \sin^{-1} \left(\frac{3}{5} \right) = \tan^{-1} \left(\frac{5}{12} \right) + \tan^{-1} \left(\frac{3}{4} \right)$

$$\left[\because \cos^{-1} \left(\frac{12}{13} \right) = \tan^{-1} \left(\frac{5}{12} \right) \text{ \& } \sin^{-1} \left(\frac{3}{5} \right) = \tan^{-1} \left(\frac{3}{4} \right) \right]$$

$$\text{L.H.S.} = \tan^{-1} \left(\frac{\frac{5}{12} + \frac{3}{4}}{1 - \frac{5}{12} \times \frac{3}{4}} \right) = \tan^{-1} \left(\frac{56}{33} \right)$$

$$\text{R.H.S.} = \sin^{-1} \left(\frac{56}{65} \right) = \tan^{-1} \left(\frac{56}{33} \right)$$

L.H.S. = R.H.S. Hence Proved

Do yourself - 5 :

Prove the following :


$$(i) \quad \sin^{-1}\left(\frac{3}{5}\right) + \sin^{-1}\left(\frac{8}{17}\right) = \cos^{-1}\left(\frac{36}{85}\right)$$

$$(ii) \quad \tan^{-1}\left(\frac{3}{4}\right) + \tan^{-1}\left(\frac{3}{5}\right) - \tan^{-1}\left(\frac{8}{19}\right) = \frac{\pi}{4}$$


$$(iii) \quad \tan^{-1}\left(\frac{2}{11}\right) + \tan^{-1}\left(\frac{7}{24}\right) = \tan^{-1}\left(\frac{1}{2}\right)$$

4. SIMPLIFIED INVERSE TRIGONOMETRIC FUNCTIONS :


$$(a) \quad y = f(x) = \sin^{-1}\left(\frac{2x}{1+x^2}\right) = \begin{cases} 2 \tan^{-1} x & \text{if } |x| \leq 1 \\ \pi - 2 \tan^{-1} x & \text{if } x > 1 \\ -(\pi + 2 \tan^{-1} x) & \text{if } x < -1 \end{cases}$$



$$(b) \quad y = f(x) = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) = \begin{cases} 2 \tan^{-1} x & \text{if } x \geq 0 \\ -2 \tan^{-1} x & \text{if } x < 0 \end{cases}$$




$$(c) \quad y = f(x) = \tan^{-1} \frac{2x}{1-x^2} = \begin{cases} 2 \tan^{-1} x & \text{if } |x| < 1 \\ \pi + 2 \tan^{-1} x & \text{if } x < -1 \\ -(\pi - 2 \tan^{-1} x) & \text{if } x > 1 \end{cases}$$




$$(d) \quad y = f(x) = \sin^{-1}(3x - 4x^3)$$

$$= \begin{cases} -(\pi + 3 \sin^{-1} x) & \text{if } -1 \leq x \leq -\frac{1}{2} \\ 3 \sin^{-1} x & \text{if } -\frac{1}{2} \leq x \leq \frac{1}{2} \\ \pi - 3 \sin^{-1} x & \text{if } \frac{1}{2} \leq x \leq 1 \end{cases}$$




(e) $y = f(x) = \cos^{-1}(4x^3 - 3x)$

$$= \begin{cases} 3\cos^{-1}x - 2\pi & \text{if } -1 \leq x \leq -\frac{1}{2} \\ 2\pi - 3\cos^{-1}x & \text{if } -\frac{1}{2} \leq x \leq \frac{1}{2} \\ 3\cos^{-1}x & \text{if } \frac{1}{2} \leq x \leq 1 \end{cases}$$



(f) $\sin^{-1}(2x\sqrt{1-x^2}) = \begin{cases} -(\pi + 2\sin^{-1}x) & -1 \leq x \leq -\frac{1}{\sqrt{2}} \\ 2\sin^{-1}x & -\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}} \\ \pi - 2\sin^{-1}x & \frac{1}{\sqrt{2}} \leq x \leq 1 \end{cases}$



(g) $\cos^{-1}(2x^2 - 1) = \begin{cases} 2\cos^{-1}x & 0 \leq x \leq 1 \\ 2\pi - 2\cos^{-1}x & -1 \leq x \leq 0 \end{cases}$




Illustration 12 : Prove that : $2\tan^{-1}\frac{1}{2} + \tan^{-1}\frac{1}{7} = \tan^{-1}\frac{31}{17}$

Solution : We have, $2\tan^{-1}\frac{1}{2} + \tan^{-1}\frac{1}{7}$

$$= \tan^{-1}\left\{\frac{2 \times \frac{1}{2}}{1 - \left(\frac{1}{2}\right)^2}\right\} + \tan^{-1}\frac{1}{7}$$

$$\left[\because 2\tan^{-1}x = \tan^{-1}\left(\frac{2x}{1-x^2}\right), \text{ if } -1 < x < 1 \right]$$

$$= \tan^{-1}\frac{4}{3} + \tan^{-1}\frac{1}{7} = \tan^{-1}\left\{\frac{\frac{4}{3} + \frac{1}{7}}{1 - \frac{4}{3} \times \frac{1}{7}}\right\} = \tan^{-1}\frac{31}{17}$$

Illustration 13 : Prove that $\tan^{-1} \sqrt{x} = \frac{1}{2} \cos^{-1} \left(\frac{1-x}{1+x} \right), x \in [0, 1]$

Solution : We have, $\frac{1}{2} \cos^{-1} \left(\frac{1-x}{1+x} \right) = \frac{1}{2} \cos^{-1} \left\{ \frac{1-(\sqrt{x})^2}{1+(\sqrt{x})^2} \right\} = \frac{1}{2} \times 2 \tan^{-1} \sqrt{x} = \tan^{-1} \sqrt{x}$.

Alter : Putting $\sqrt{x} = \tan \theta$, we have $\Rightarrow \theta \in \left[0, \frac{\pi}{4} \right]$

$$\begin{aligned} \text{RHS} &= \frac{1}{2} \cos^{-1} \left(\frac{1-x}{1+x} \right) = \frac{1}{2} \cos^{-1} \left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta} \right) = \frac{1}{2} \cos^{-1} (\cos 2\theta) = \theta \quad \because \left(2\theta \in \left[0, \frac{\pi}{2} \right] \right) \\ &= \tan^{-1} \sqrt{x} = \text{LHS} \end{aligned}$$

Illustration 14 : Prove that : (i) $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99} = \frac{\pi}{4}$

$$(ii) \quad 2 \tan^{-1} \frac{1}{5} + \sec^{-1} \frac{5\sqrt{2}}{7} + 2 \tan^{-1} \frac{1}{8} = \frac{\pi}{4}$$

Solution : (i) $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99} = 2 \left\{ 2 \tan^{-1} \frac{1}{5} \right\} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99}$

$$= 2 \left\{ \tan^{-1} \frac{2 \times 1/5}{1 - (1/5)^2} \right\} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99} \quad \left[\begin{array}{l} \because 2 \tan^{-1} x \\ = \tan^{-1} \frac{2x}{1-x^2}, \text{ if } |x| < 1 \end{array} \right]$$

$$= 2 \tan^{-1} \frac{5}{12} - \left\{ \tan^{-1} \frac{1}{70} - \tan^{-1} \frac{1}{99} \right\} = \tan^{-1} \left\{ \frac{2 \times 5/12}{1 - (5/12)^2} \right\} - \tan^{-1} \left\{ \frac{\frac{1}{70} - \frac{1}{99}}{1 + \frac{1}{70} \times \frac{1}{99}} \right\}$$

$$= \tan^{-1} \frac{120}{119} - \tan^{-1} \frac{29}{6931} = \tan^{-1} \frac{120}{119} - \tan^{-1} \frac{1}{239} = \tan^{-1} \left\{ \frac{\frac{120}{119} - \frac{1}{239}}{1 + \frac{120}{119} \times \frac{1}{239}} \right\} = \tan^{-1} 1 = \frac{\pi}{4}$$

$$(ii) \quad 2 \tan^{-1} \frac{1}{5} + \sec^{-1} \frac{5\sqrt{2}}{7} + 2 \tan^{-1} \frac{1}{8} = 2 \left\{ \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{8} \right\} + \sec^{-1} \frac{5\sqrt{2}}{7}$$

$$\begin{aligned}
 &= 2 \tan^{-1} \left\{ \frac{\frac{1}{5} + \frac{1}{8}}{1 - \frac{1}{5} \times \frac{1}{8}} \right\} + \tan^{-1} \sqrt{\left(\frac{5\sqrt{2}}{7} \right)^2 - 1} \quad \left[\because \sec^{-1} x = \tan^{-1} \sqrt{x^2 - 1} \right] \\
 &= 2 \tan^{-1} \frac{13}{39} + \tan^{-1} \frac{1}{7} = 2 \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{7} \\
 &= \tan^{-1} \left\{ \frac{2 \times 1/3}{1 - (1/3)^2} \right\} + \tan^{-1} \frac{1}{7} \quad \left[\because 2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2}, \text{ if } |x| < 1 \right] \\
 &= \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{1}{7} = \tan^{-1} \left\{ \frac{\frac{3}{4} + \frac{1}{7}}{1 - \frac{3}{4} \times \frac{1}{7}} \right\} = \tan^{-1} 1 = \frac{\pi}{4}
 \end{aligned}$$

Do yourself - 6 :

Prove the following results :

$$\begin{aligned}
 \text{(i)} \quad & 2 \tan^{-1} \left(\frac{1}{5} \right) + \tan^{-1} \left(\frac{1}{8} \right) = \tan^{-1} \left(\frac{4}{7} \right) \qquad \text{(ii)} \quad 2 \sin^{-1} \left(\frac{3}{5} \right) - \tan^{-1} \left(\frac{17}{31} \right) = \frac{\pi}{4}
 \end{aligned}$$

6. EQUATIONS INVOLVING INVERSE TRIGONOMETRIC FUNCTIONS :

Illustration 15 : The equation $2\cos^{-1}x + \sin^{-1}x = \frac{11\pi}{6}$ has

- (A) no solution (B) only one solution (C) two solutions (D) three solutions

Solution : Given equation is $2 \cos^{-1} x + \sin^{-1} x = \frac{11\pi}{6}$

$$\Rightarrow \cos^{-1} x + (\cos^{-1} x + \sin^{-1} x) = \frac{11\pi}{6} \Rightarrow \cos^{-1} x + \frac{\pi}{2} = \frac{11\pi}{6} \Rightarrow \cos^{-1} x = 4\pi/3$$

which is not possible as $\cos^{-1} x \in [0, \pi]$

Ans.(A)

Illustration 16 : If $(\tan^{-1} x)^2 + (\cot^{-1} x)^2 = 5\pi^2/8$, then x is equal to-

- (A) -1 (B) 0 (C) 1 (D) none of these

Solution : The given equation can be written as $(\tan^{-1} x + \cot^{-1} x)^2 - 2 \tan^{-1} x \cot^{-1} x = 5\pi^2/8$

Since $\tan^{-1} x + \cot^{-1} x = \pi/2$ we have

$$(\pi/2)^2 - 2 \tan^{-1} x (\pi/2 - \tan^{-1} x) = 5\pi^2/8$$

$$\Rightarrow 2(\tan^{-1} x)^2 - 2(\pi/2) \tan^{-1} x - 3\pi^2/8 = 0 \Rightarrow \tan^{-1} x = -\pi/4 \Rightarrow x = -1 \quad \text{Ans. (A)}$$

Illustration 17 : Solve the equation : $\tan^{-1} \frac{x-1}{x-2} + \tan^{-1} \frac{x+1}{x+2} = \frac{\pi}{4}$

Solution : $\tan^{-1} \frac{x-1}{x-2} + \tan^{-1} \frac{x+1}{x+2} = \frac{\pi}{4}$

taking tangent on both sides

$$\Rightarrow \tan \left(\tan^{-1} \left(\frac{x-1}{x-2} \right) + \tan^{-1} \left(\frac{x+1}{x+2} \right) \right) = 1 \Rightarrow \frac{\tan \left(\tan^{-1} \left(\frac{x-1}{x-2} \right) \right) + \tan \left(\tan^{-1} \left(\frac{x+1}{x+2} \right) \right)}{1 - \tan \left(\tan^{-1} \left(\frac{x-1}{x-2} \right) \right) \tan \left(\tan^{-1} \left(\frac{x+1}{x+2} \right) \right)} = 1$$

$$\Rightarrow \frac{\frac{x-1}{x-2} + \frac{x+1}{x+2}}{1 - \frac{x-1}{x-2} \cdot \frac{x+1}{x+2}} = 1 \Rightarrow \frac{(x-1)(x+2) + (x-2)(x+1)}{x^2 - 4 - (x^2 - 1)} = 1 \Rightarrow 2x^2 - 4 = -3 \Rightarrow x = \pm \frac{1}{\sqrt{2}}$$

Now verify $x = \frac{1}{\sqrt{2}}$

$$= \tan^{-1} \left(\frac{\frac{1}{\sqrt{2}} - 1}{\frac{1}{\sqrt{2}} - 2} \right) + \tan^{-1} \left(\frac{\frac{1}{\sqrt{2}} + 1}{\frac{1}{\sqrt{2}} + 2} \right) = \tan^{-1} \left(\frac{\sqrt{2} - 1}{2\sqrt{2} - 1} \right) + \tan^{-1} \left(\frac{\sqrt{2} + 1}{2\sqrt{2} + 1} \right)$$

$$= \tan^{-1} \left(\frac{(2\sqrt{2} + 1)(\sqrt{2} - 1) + (2\sqrt{2} - 1)(\sqrt{2} + 1)}{(2\sqrt{2} - 1)(2\sqrt{2} + 1) - (\sqrt{2} - 1)(\sqrt{2} + 1)} \right) = \tan^{-1} \left(\frac{6}{6} \right) = \tan^{-1}(1) = \frac{\pi}{4}$$

$$x = -\frac{1}{\sqrt{2}}$$

$$= \tan^{-1} \left(\frac{-\frac{1}{\sqrt{2}} - 1}{-\frac{1}{\sqrt{2}} - 2} \right) + \tan^{-1} \left(\frac{-\frac{1}{\sqrt{2}} + 1}{-\frac{1}{\sqrt{2}} + 2} \right) = \tan^{-1} \left(\frac{\sqrt{2} + 1}{2\sqrt{2} + 1} \right) + \tan^{-1} \left(\frac{\sqrt{2} - 1}{2\sqrt{2} - 1} \right) \{ \text{same as above} \}$$

$$= \tan^{-1}(1) = \frac{\pi}{4}$$

$$\therefore x = \pm \frac{1}{\sqrt{2}} \text{ are solutions}$$

Ans.

Illustration 18 : Solve the equation : $2 \tan^{-1}(2x + 1) = \cos^{-1}x$.

Solution : Here, $2 \tan^{-1}(2x + 1) = \cos^{-1}x$

$$\text{or } \cos(2 \tan^{-1}(2x + 1)) = x \quad \left\{ \text{We know } \cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right\}$$

$$\therefore \frac{1 - (2x + 1)^2}{1 + (2x + 1)^2} = x \Rightarrow (1 - 2x - 1)(1 + 2x + 1) = x(4x^2 + 4x + 2)$$

$$\Rightarrow -2x \cdot 2(x+1) = 2x(2x^2 + 2x + 1) \Rightarrow 2x(2x^2 + 2x + 1 + 2x + 2) = 0$$

$$\Rightarrow 2x(2x^2 + 4x + 3) = 0$$

$$\Rightarrow x = 0 \text{ or } 2x^2 + 4x + 3 = 0 \quad \{\text{No solution}\}$$

Verify $x = 0$

$$2\tan^{-1}(1) = \cos^{-1}(1) \Rightarrow \frac{\pi}{2} = \frac{\pi}{2}$$

$\therefore x = 0$ is only the solution

Ans.

Do yourself - 7 :

Solve the following equation for x :

$$(i) \quad \sin \left[\sin^{-1} \left(\frac{1}{5} \right) + \cos^{-1} x \right] = 1 \quad (ii) \quad \cos^{-1} x + \sin^{-1} \frac{x}{2} = \frac{\pi}{6}$$

7. INEQUATIONS INVOLVING INVERSE TRIGONOMETRIC FUNCTION :

Illustration 19 : Find the complete solution set of $\sin^{-1}(\sin 5) > x^2 - 4x$.

Solution : $\sin^{-1}(\sin 5) > x^2 - 4x \Rightarrow \sin^{-1}[\sin(5 - 2\pi)] > x^2 - 4x$

$$\Rightarrow x^2 - 4x < 5 - 2\pi \Rightarrow x^2 - 4x + (2\pi - 5) < 0$$

$$\Rightarrow 2 - \sqrt{9 - 2\pi} < x < 2 + \sqrt{9 - 2\pi} \Rightarrow x \in (2 - \sqrt{9 - 2\pi}, 2 + \sqrt{9 - 2\pi})$$

Ans.

Illustration 20 : Find the complete solution set of $[\cot^{-1}x]^2 - 6[\cot^{-1}x] + 9 \leq 0$, where $[.]$ denotes the greatest integer function.

Solution : $[\cot^{-1}x]^2 - 6[\cot^{-1}x] + 9 \leq 0$

$$\Rightarrow ([\cot^{-1}x] - 3)^2 \leq 0 \Rightarrow [\cot^{-1}x] = 3 \Rightarrow 3 \leq \cot^{-1}x < 4 \Rightarrow x \in (-\infty, \cot 3]$$

Illustration 21 : If $\cot^{-1} \frac{n}{\pi} > \frac{\pi}{6}$, $n \in \mathbb{N}$, then the maximum value of n is -

- (A) 1 (B) 5 (C) 9 (D) none of these

Solution : $\cot^{-1} \frac{n}{\pi} > \frac{\pi}{6}$

$$\Rightarrow \cot \left(\cot^{-1} \left(\frac{n}{\pi} \right) \right) < \cot \left(\frac{\pi}{6} \right) \Rightarrow \frac{n}{\pi} < \sqrt{3}$$

$$\Rightarrow n < \pi\sqrt{3} \Rightarrow n < 5.5 \text{ (approx)}$$

$$\Rightarrow n = 5 \quad \because (n \in \mathbb{N})$$

Ans. (B)

Do yourself - 8 :

- (i) Solve the inequality $\tan^{-1}x > \cot^{-1}x$.
 (ii) Complete solution set of inequation $(\cos^{-1}x)^2 - (\sin^{-1}x)^2 > 0$, is

- (A) $\left[0, \frac{1}{\sqrt{2}}\right)$ (B) $\left[-1, \frac{1}{\sqrt{2}}\right)$ (C) $(-1, \sqrt{2})$ (D) none of these

8. SUMMATION OF SERIES :

Illustration 22 : Prove that :

$$\tan^{-1}\left(\frac{c_1x-y}{c_1y+x}\right) + \tan^{-1}\left(\frac{c_2-c_1}{1+c_2c_1}\right) + \tan^{-1}\left(\frac{c_3-c_2}{1+c_3c_2}\right) + \dots + \tan^{-1}\left(\frac{c_n-c_{n-1}}{1+c_nc_{n-1}}\right) + \tan^{-1}\left(\frac{1}{c_n}\right) = \tan^{-1}\left(\frac{x}{y}\right)$$

Solution : L.H.S.

$$\tan^{-1}\left(\frac{c_1x-y}{c_1y+x}\right) + \tan^{-1}\left(\frac{c_2-c_1}{1+c_2c_1}\right) + \tan^{-1}\left(\frac{c_3-c_2}{1+c_3c_2}\right) + \dots + \tan^{-1}\left(\frac{c_n-c_{n-1}}{1+c_nc_{n-1}}\right) + \tan^{-1}\left(\frac{1}{c_n}\right)$$

$$= \tan^{-1}\left(\frac{\frac{x}{y} - \frac{1}{c_1}}{1 + \frac{x}{y} \cdot \frac{1}{c_1}}\right) + (\tan^{-1}c_2 - \tan^{-1}c_1) + (\tan^{-1}c_3 - \tan^{-1}c_2) + \dots + (\tan^{-1}c_n - \tan^{-1}c_{n-1}) + \tan^{-1}\left(\frac{1}{c_n}\right)$$

$$= \tan^{-1}\left(\frac{x}{y}\right) - \tan^{-1}\left(\frac{1}{c_1}\right) - \tan^{-1}c_1 + \tan^{-1}c_n + \tan^{-1}\left(\frac{1}{c_n}\right)$$

$$= \tan^{-1}\left(\frac{x}{y}\right) - (\cot^{-1}c_1 + \tan^{-1}c_1) + (\tan^{-1}c_n + \cot^{-1}c_n)$$

$$= \tan^{-1}\left(\frac{x}{y}\right) - \frac{\pi}{2} + \frac{\pi}{2} = \tan^{-1}\left(\frac{x}{y}\right) = \text{R.H.S.}$$

Do yourself - 9 :

- (i) Evaluate : $\sum_{r=1}^{\infty} \tan^{-1}\left(\frac{2}{1+(2r+1)(2r-1)}\right)$

Miscellaneous Illustrations :

Illustration 23 : If $\tan^{-1} y = 4 \tan^{-1} x$, $\left(|x| < \tan \frac{\pi}{8} \right)$, find y as an algebraic function of x and hence prove

that $\tan \frac{\pi}{8}$ is a root of the equation $x^4 - 6x^2 + 1 = 0$.

Solution :

We have $\tan^{-1} y = 4 \tan^{-1} x$

$$\Rightarrow \tan^{-1} y = 2 \tan^{-1} \frac{2x}{1-x^2} \quad (\text{as } |x| < 1)$$

$$= \tan^{-1} \frac{\frac{4x}{1-x^2}}{1 - \frac{4x^2}{(1-x^2)^2}} = \tan^{-1} \frac{4x(1-x^2)}{x^4 - 6x^2 + 1} \quad \left(\text{as } \left| \frac{2x}{1-x^2} \right| < 1 \right)$$

$$\Rightarrow y = \frac{4x(1-x^2)}{x^4 - 6x^2 + 1}$$

$$\text{If } x = \tan \frac{\pi}{8}$$

$$\Rightarrow \tan^{-1} y = 4 \tan^{-1} x = \frac{\pi}{2} \Rightarrow y \text{ is not defined} \Rightarrow x^4 - 6x^2 + 1 = 0 \quad \text{Ans.}$$

Illustration 24 : If $A = 2 \tan^{-1}(2\sqrt{2}-1)$ and $B = 3 \sin^{-1}(1/3) + \sin^{-1}(3/5)$, then show $A > B$.

Solution :

We have, $A = 2 \tan^{-1}(2\sqrt{2}-1) = 2 \tan^{-1}(1.828)$

$$\Rightarrow A > 2 \tan^{-1}(\sqrt{3}) \Rightarrow A > \frac{2\pi}{3} \quad \dots\dots (i)$$

$$\text{also we have, } \sin^{-1}\left(\frac{1}{3}\right) < \sin^{-1}\left(\frac{1}{2}\right) \Rightarrow \sin^{-1}\left(\frac{1}{3}\right) < \frac{\pi}{6}$$

$$\Rightarrow 3 \sin^{-1}\left(\frac{1}{3}\right) < \frac{\pi}{2}$$

$$\text{also, } 3 \sin^{-1}\left(\frac{1}{3}\right) = \sin^{-1}\left(3 \cdot \frac{1}{3} - 4 \left(\frac{1}{3}\right)^3\right) = \sin^{-1}\left(\frac{23}{27}\right) = \sin^{-1}(0.852)$$

$$\Rightarrow 3 \sin^{-1}(1/3) < \sin^{-1}(\sqrt{3}/2) \Rightarrow 3 \sin^{-1}(1/3) < \pi/3$$

$$\text{also, } \sin^{-1}(3/5) = \sin^{-1}(0.6) < \sin^{-1}(\sqrt{3}/2) \Rightarrow \sin^{-1}(3/5) < \pi/3$$

$$\text{Hence, } B = 3 \sin^{-1}(1/3) + \sin^{-1}(3/5) < \frac{2\pi}{3} \quad \dots\dots (ii)$$

From (i) and (ii), we have $A > B$.

Illustration 25 : If $\theta = \tan^{-1}(2 \tan^2 \theta) - \frac{1}{2} \sin^{-1} \left(\frac{3 \sin 2\theta}{5 + 4 \cos 2\theta} \right)$ then find the sum of all possible values of $\tan \theta$.

Solution :

$$\theta = \tan^{-1}(2 \tan^2 \theta) - \frac{1}{2} \sin^{-1} \left(\frac{3 \sin 2\theta}{5 + 4 \cos 2\theta} \right) \Rightarrow \theta = \tan^{-1}(2 \tan^2 \theta) - \frac{1}{2} \sin^{-1} \left(\frac{6 \tan \theta}{9 + \tan^2 \theta} \right)$$

$$\Rightarrow \theta = \tan^{-1}(2 \tan^2 \theta) - \frac{1}{2} \sin^{-1} \left[\frac{2 \left(\frac{1}{3} \tan \theta \right)}{1 + \left(\frac{1}{3} \tan \theta \right)^2} \right] \Rightarrow \theta = \tan^{-1}(2 \tan^2 \theta) - \frac{2}{2} \tan^{-1} \left(\frac{1}{3} \tan \theta \right)$$

$$\Rightarrow \theta = \tan^{-1}(2 \tan^2 \theta) - \tan^{-1} \left(\frac{1}{3} \tan \theta \right) \quad \dots\dots\dots (i)$$

taking tangent on both sides

$$\Rightarrow \tan \theta = \frac{6 \tan^2 \theta - \tan \theta}{3 + 2 \tan^3 \theta} \Rightarrow 2 \tan^4 \theta - 6 \tan^2 \theta + 4 \tan \theta = 0$$

$$\Rightarrow 2 \tan \theta (\tan^3 \theta - 3 \tan \theta + 2) = 0 \Rightarrow 2 \tan \theta (\tan \theta - 1)^2 (\tan \theta + 2) = 0$$

$$\Rightarrow \tan \theta = 0, 1, -2 \quad \text{which satisfy equation (i)}$$

$$\therefore \text{sum} = 0 + 1 - 2 = -1$$

Ans.

ANSWERS FOR DO YOURSELF

- | | | | | | | |
|-----|------|---|------|-----------------------|-------|-----------------|
| 1 : | (i) | C | (ii) | 1526 | | |
| 2 : | (i) | $\frac{15}{8}$ | (ii) | $\frac{1}{\sqrt{10}}$ | (iii) | 1 |
| | | | | | (iv) | $\frac{4}{5}$ |
| 3 : | (i) | $\frac{\pi}{6}$ | (ii) | $\frac{\pi}{6}$ | (iii) | $\frac{\pi}{6}$ |
| 4 : | (ii) | $\begin{cases} 1, & \text{if } a > 0 \\ -1, & \text{if } a < 0 \end{cases}$ | | | | |
| 7 : | (i) | $\frac{1}{5}$ | (ii) | 1 | (iii) | $\sqrt{3}$ |
| 8 : | (i) | $(1, \infty)$ | (ii) | B | | |
| 9 : | (i) | $\pi/4$ | | | | |

EXERCISE (O-1)

Straight Objective Type

- The domain of the function $\sin^{-1}\left(\log_2\left(\frac{x}{3}\right)\right)$ is-
 (A) $\left[\frac{1}{2}, 3\right]$ (B) $\left[\frac{1}{2}, 3\right]$ (C) $\left[\frac{3}{2}, 6\right]$ (D) $\left[\frac{1}{2}, 2\right]$
IT0001
- Domain of the function $f(x) = \log_e \cos^{-1}\{\sqrt{x}\}$ is, where $\{.\}$ represents fractional part function -
 (A) $x \in \mathbb{R}$ (B) $x \in [0, \infty)$ (C) $x \in (0, \infty)$ (D) $x \in \mathbb{R} - \{x \mid x \in I\}$
IT0002
- The value of $\tan^2(\sec^{-1}3) + \cot^2(\operatorname{cosec}^{-1}4)$ is -
 (A) 9 (B) 16 (C) 25 (D) 23
IT0003
- $\tan^{-1}\left(1 - x^2 - \frac{1}{x^2}\right) + \sin^{-1}\left(x^2 + \frac{1}{x^2} - 1\right)$ (where $x \neq 0$) is equal to
 (A) $\frac{\pi}{2}$ (B) $\frac{\pi}{4}$ (C) $\frac{3\pi}{4}$ (D) π
IT0004
- If $2 \leq a < 3$, then the value of $\cos^{-1} \cos[a] + \operatorname{cosec}^{-1} \operatorname{cosec}[a] + \cot^{-1} \cot[a]$, (where $[.]$ denotes greatest integer less than equal to x) is equal to
 (A) $2 - \pi$ (B) $2 + \pi$ (C) π (D) 6
IT0005
- If $x > 0$ $\cos^{-1}\left(\frac{12}{x}\right) = \frac{\pi}{2} - \cos^{-1}\left(\frac{16}{x}\right)$ then x is -
 (A) 12 (B) 16 (C) 20 (D) 320
IT0006
- Number of integral ordered pairs (a, b) for which $\sin^{-1}(1 + b + b^2 + \dots \infty) + \cos^{-1}\left(a - \frac{a^2}{3} + \frac{a^3}{9} - \dots \infty\right) = \frac{\pi}{2}$ is-
 (A) 0 (B) 4 (C) 9 (D) Infinitely many
IT0007
- If $\cos^{-1}(2x^2 - 1) = 2\pi - 2\cos^{-1}x$, then -
 (A) $x \in [-1, 0]$ (B) $x \in [0, 1]$ (C) $x \in \left[0, \frac{1}{\sqrt{2}}\right]$ (D) $x \in \left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right]$
IT0008

9. $\lim_{n \rightarrow \infty} \sum_{r=1}^n \tan^{-1} \frac{2r+1}{r^4 + 2r^3 + r^2 + 1}$ is equal to -

(A) $\frac{\pi}{4}$

(B) $\frac{3\pi}{4}$

(C) $\frac{\pi}{2}$

(D) $-\frac{\pi}{8}$

IT0009

Multiple Correct Answer Type

10. If $\sin\left(2\cos^{-1}\left(\frac{1}{\sqrt{5}}\right)\right) + \cos\left(2\tan^{-1}\left(\frac{1}{3}\right)\right) = \frac{p}{q}$, where p & q are relatively prime then digit at units place of $(p-q)^{2k+1}$, $k \in \mathbb{N}$, can be -

(A) 1

(B) 3

(C) 7

(D) 9

IT0010

11. Let $f(x) = \sin^{-1}(\tan x) + \cos^{-1}(\cot x)$ then

(A) $f(x) = \frac{\pi}{2}$ wherever defined

(B) domain of $f(x)$ is $x = n\pi \pm \frac{\pi}{4}$, $n \in \mathbb{I}$

(C) period of $f(x)$ is $\frac{\pi}{2}$

(D) $f(x)$ is many one function

IT0011

12. Which of the following is/are correct ?

(A) $\cot^{-1}(x) = \tan^{-1}\left(\frac{1}{x}\right) \forall x \in \mathbb{R} - \{0\}$

(B) If $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = \operatorname{sgn}(e^x)$ then $f(x)$ is an into function.

(C) If $f: \mathbb{R}^+ \rightarrow \mathbb{R}$ such that $f(x) = \sin x + x$ then $f(x)$ is an odd function.

(D) If $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = \frac{e^x}{e^{[x]}}$ then $f(x)$ is a periodic function .

(where $[.]$ represents greatest integer function)

IT0012

13. Consider the function $f(x) = e^x$ and $g(x) = \sin^{-1}x$, then which of the following is/are necessarily true.

(A) Domain of $g \circ f$ = Domain of f

(B) Range of $g \circ f \subset$ Range of g

(C) Domain of $g \circ f$ is $(-\infty, 0]$

(D) Range of $g \circ f$ is $\left[-\frac{\pi}{2}, 0\right]$

IT0013

14. Let $f(x) = e^{x^3 - x^2 + x}$ be an invertible function such that $f^{-1} = g$, then -

(A) $g(e) = 0$

(B) Domain of 'g' is \mathbb{R}^+

(C) Range of 'g' is \mathbb{R}

(D) $f(g(e)) = e$

IT0014

15. Value of $3 \tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{2}\right) + \sin^{-1}\left(\frac{1}{\sqrt{5}}\right) + \cos^{-1}\left(\frac{2}{\sqrt{5}}\right)$ is greater than

(A) $\frac{\pi}{2}$

(B) $\frac{2\pi}{3}$

(C) $\frac{3\pi}{4}$

(D) $\frac{5\pi}{6}$

IT0015

EXERCISE (O-2)

Straight Objective Type

1. The range of the function $f(x) = \sin^{-1}(\log_2(-x^2 + 2x + 3))$ is -

(A) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

(B) $\left[-\frac{\pi}{2}, 0\right]$

(C) $\left[0, \frac{\pi}{2}\right]$

(D) $[-1, 1]$

IT0016

2. Range of $f(x) = \cot^{-1}(\log_e(1 - x^2))$ is -

(A) $(0, \pi)$

(B) $\left(0, \frac{\pi}{2}\right)$

(C) $\left[\frac{\pi}{2}, \pi\right)$

(D) $\left(0, \frac{\pi}{2}\right]$

IT0017

3. The value of $\sin^{-1}\left\{\cot\left(\sin^{-1}\sqrt{\frac{2-\sqrt{3}}{4}} + \cos^{-1}\frac{\sqrt{12}}{4} + \sec^{-1}\sqrt{2}\right)\right\}$ is -

(A) 5

(B) 6

(C) 0

(D) 10

IT0018

4. Number of solution(s) of the equation $\cos^{-1}\sqrt{x} - \sin^{-1}\sqrt{x-1} + \cos^{-1}\sqrt{1-x} - \sin^{-1}\frac{1}{\sqrt{x}} = \frac{\pi}{2}$ is -

(A) 0

(B) 1

(C) 2

(D) 4

IT0019

5. $\sum_{r=0}^{\infty} \tan^{-1}\left(\frac{r((r+1)!)}{(r+1) + ((r+1)!)^2}\right)$ is equal to -

(A) $\frac{\pi}{2}$

(B) $\frac{\pi}{4}$

(C) $\cot^{-1}3$

(D) $\tan^{-1}2$

IT0020

Multiple Correct Answer Type

6. Let $f(x) = \sqrt{\cos^{-1} \sqrt{1-x^2} - \sin^{-1} x}$ then which of the following statement/s is/are correct -

(A) Domain of $f(x)$ is $[-1, 1]$ (B) Domain of $f(x)$ is $[0, 1]$
 (C) Range of $f(x)$ is $\{0\}$ (D) Range of $f(x)$ is $[0, \sqrt{\pi}]$

IT0021

7. If $\alpha = 2 \tan^{-1}(\sqrt{3-2\sqrt{2}}) + \sin^{-1}\left(\frac{1}{\sqrt{6}-\sqrt{2}}\right)$, $\beta = \cot^{-1}(\sqrt{3}-2) + \frac{1}{8} \sec^{-1}(-2)$ & $\gamma = \tan^{-1} \frac{1}{\sqrt{2}} + \cos^{-1} \frac{1}{\sqrt{3}}$, then

(A) $\alpha = \beta$ (B) $\alpha + \beta = 3\gamma$ (C) $4(\beta - \gamma) = \alpha$ (D) $\beta = \gamma$

IT0022

8. If α is only real root of the equation $x^3 + (\cos 1)x^2 + (\sin 1)x + 1 = 0$, then $\left(\tan^{-1} \alpha + \tan^{-1} \frac{1}{\alpha}\right)$ cannot be equal to-

(A) 0 (B) $\frac{\pi}{2}$ (C) $-\frac{\pi}{2}$ (D) π

IT0023

9. Let $f(x) = \begin{cases} x^2 - 4 & \text{if } |x| \leq 3 \\ 5 \operatorname{sgn}|x-3| & \text{if } |x| > 3 \end{cases}$ and $g(x) = 2 \tan^{-1}(e^x) - \frac{\pi}{2}$ for all $x \in \mathbb{R}$, then which of the following is wrong ?

(where $\operatorname{sgn}(x)$ denotes signum function of x)

(A) $f(x)$ is an even function (B) $g \circ f(x)$ is an even function
 (C) $g(x)$ is an odd function (D) $f \circ g(x)$ is an odd function

IT0024

Linked Comprehension Type**Paragraph for Question 10 to 12**

Consider a continuous function such that each image has atmost three preimage & atleast one image has exactly three preimages. This type of function is to be called as three-one function.

On the basis of above information, answer the following questions :

10. Which of the following function is a three-one function ?

(A) $|\ell n|x||$ (B) $e^{|x|}$ (C) $x^3 + 3x^2 - 7x + 6$ (D) $\cos(\cos^{-1}x)$

IT0025

11. If $f(x) = \sin^{-1}(\sin x)$ is a three-one function, then possible interval of x is -

(A) $[-\pi, \pi]$ (B) $\left[-\frac{\pi}{2}, \frac{3\pi}{2}\right]$ (C) $(-2\pi, 0]$ (D) $\left[\frac{-3\pi}{2}, \frac{\pi}{2}\right]$

IT0025

12. If $f(x)$ is a three-one function such that $f(a) = f(b)$ (where $a \neq b$), then number of maximum possible values of b is -

(A) 1 (B) 2 (C) 3 (D) 4

IT0025

Subjective Type Questions

13. If $\sin^{-1}\sin\left(\frac{10\pi}{3}\right) + \cos^{-1}\cos\frac{22\pi}{3} + \tan^{-1}\tan 10 = a\pi + b$, then $\left|\frac{3ab}{80}\right|$ is equal to

IT0026

14. Number of integral solutions of the equation $2\sin^{-1}\sqrt{x^2 - x + 1} + \cos^{-1}\sqrt{x^2 - x} = \frac{3\pi}{2}$ is

IT0027

15. The product of all real values of x satisfying the equation

$$\sin^{-1}\cos\left(\frac{2x^2 + 10|x| + 4}{x^2 + 5|x| + 3}\right) = \cot\left(\cot^{-1}\left(\frac{2 - 18|x|}{9|x|}\right)\right) + \frac{\pi}{2}$$

IT0028

16. Least integral value of x for which inequality $\sin^{-1}\left(\sin\left(\frac{2e^x + 3}{e^x + 1}\right)\right) > \pi - \frac{5}{2}$ holds, is


IT0029

EXERCISE (S-1)

1. Given is a partial graph of an even periodic function f whose period is 8.

If $[*]$ denotes greatest integer function then find the value of the expression.

$$f(-3) + 2|f(-1)| + \left[f\left(\frac{7}{8}\right)\right] + f(0) + \arccos(f(-2)) + f(-7) + f(20)$$



IT0030

2. (a) Find the following :

(i) $\tan\left[\cos^{-1}\frac{1}{2} + \tan^{-1}\left(\frac{-1}{\sqrt{3}}\right)\right]$

(ii) $\cos^{-1}\left(\cos\frac{7\pi}{6}\right)$

(iii) $\cos\left(\tan^{-1}\frac{3}{4}\right)$

(iv) $\tan\left(\sin^{-1}\frac{3}{5} + \cot^{-1}\frac{3}{2}\right)$

IT0031

(b) Find the following :

(i) $\sin\left[\frac{\pi}{2} - \sin^{-1}\left(\frac{-\sqrt{3}}{2}\right)\right]$

(ii) $\cos\left[\cos^{-1}\left(\frac{-\sqrt{3}}{2}\right) + \frac{\pi}{6}\right]$

(iii) $\tan^{-1}\left(\tan\frac{3\pi}{4}\right)$

(iv) $\sin\left(\frac{1}{4}\arcsin\frac{\sqrt{63}}{8}\right)$

IT0032

3. Find the domain of definition the following functions.

(Read the symbols $[*]$ and $\{*\}$ as greatest integers and fractional part functions respectively)

(i) $f(x) = \arccos \frac{2x}{1+x}$ IT0033

(ii) $f(x) = \sqrt{\cos(\sin x)} + \sin^{-1} \frac{1+x^2}{2x}$ IT0034

(iii) $f(x) = \sin^{-1} \left(\frac{x-3}{2} \right) - \log_{10}(4-x)$ IT0035

(iv) $f(x) = \sin^{-1}(2x + x^2)$ IT0036

(v) $f(x) = \frac{\sqrt{1-\sin x}}{\log_5(1-4x^2)} + \cos^{-1}(1-\{x\})$, where $\{x\}$ is the fractional part of x . IT0037

(vi) $f(x) = \sqrt{3-x} + \cos^{-1} \left(\frac{3-2x}{5} \right) + \log_6(2|x|-3) + \sin^{-1}(\log_2 x)$ IT0038

(vii) $f(x) = e^{\sin^{-1} \left(\frac{x}{2} \right)} + \tan^{-1} \left[\frac{x}{2} - 1 \right] + \ln(\sqrt{x} - [x])$ IT0039

4. Identify the pair(s) of functions which are identical. Also plot the graphs in each case.

(a) $y = \tan(\cos^{-1} x); y = \frac{\sqrt{1-x^2}}{x}$ (b) $y = \tan(\cot^{-1} x); y = \frac{1}{x}$

(c) $y = \sin(\arctan x); y = \frac{x}{\sqrt{1+x^2}}$ (d) $y = \cos(\arctan x); y = \sin(\arccot x)$

IT0040

5. Let $y = \sin^{-1}(\sin 8) - \tan^{-1}(\tan 10) + \cos^{-1}(\cos 12) - \sec^{-1}(\sec 9) + \cot^{-1}(\cot 6) - \operatorname{cosec}^{-1}(\operatorname{cosec} 7)$. If y simplifies to $a\pi + b$, then find $(a-b)$.

IT0041

6. Show that : $\sin^{-1} \left(\sin \frac{33\pi}{7} \right) + \cos^{-1} \left(\cos \frac{46\pi}{7} \right) + \tan^{-1} \left(-\tan \frac{13\pi}{8} \right) + \cot^{-1} \left(\cot \left(-\frac{19\pi}{8} \right) \right) = \frac{13\pi}{7}$

IT0042

7. Prove that : (a) $2\cos^{-1} \frac{3}{\sqrt{13}} + \cot^{-1} \frac{16}{63} + \frac{1}{2}\cos^{-1} \frac{7}{25} = \pi$ (b) $\arccos \sqrt{\frac{2}{3}} - \arccos \frac{\sqrt{6}+1}{2\sqrt{3}} = \frac{\pi}{6}$

IT0043

8. If α and β are the roots of the equation $x^2 + 5x - 49 = 0$, then find the value of $\cot(\cot^{-1}\alpha + \cot^{-1}\beta)$.

IT0044

9. If $a > b > c > 0$, then find the value of : $\cot^{-1} \left(\frac{1+ab}{a-b} \right) + \cot^{-1} \left(\frac{1+bc}{b-c} \right) + \cot^{-1} \left(\frac{1+ca}{c-a} \right)$.

IT0045

10. Find all values of k for which there is a triangle whose angles have measure $\tan^{-1}\left(\frac{1}{2}\right)$, $\tan^{-1}\left(\frac{1}{2} + k\right)$ and $\tan^{-1}\left(\frac{1}{2} + 2k\right)$.

IT0046

11. Find the simplest value of

(a) $f(x) = \arccos x + \arccos\left(\frac{x}{2} + \frac{1}{2}\sqrt{3-3x^2}\right)$, $x \in \left(\frac{1}{2}, 1\right)$

IT0047

(b) $f(x) = \tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$, $x \in \mathbb{R} - \{0\}$

IT0048

12. Prove that the identities :

(a) $\sin^{-1}\cos(\sin^{-1}x) + \cos^{-1}\sin(\cos^{-1}x) = \frac{\pi}{2}$, $|x| \leq 1$

IT0049

(b) $\tan(\tan^{-1}x + \tan^{-1}y + \tan^{-1}z) = \cot(\cot^{-1}x + \cot^{-1}y + \cot^{-1}z)$

IT0050

13. (a) Solve the inequality : $(\arccos x)^2 - 6(\arccos x) + 8 > 0$

IT0051

(b) If $\sin^2 x + \sin^2 y < 1$; $x, y \in \mathbb{R}$ then prove that $\sin^{-1}(\tan x \cdot \tan y) \in (-\pi/2, \pi/2)$.

IT0052

14. Solve the following :

(a) $\sin^{-1} x + \sin^{-1} 2x = \frac{\pi}{3}$

IT0053

(b) $\tan^{-1} \frac{1}{1+2x} + \tan^{-1} \frac{1}{1+4x} = \tan^{-1} \frac{2}{x^2}$

IT0054

(c) $\tan^{-1}(x-1) + \tan^{-1}(x) + \tan^{-1}(x+1) = \tan^{-1}(3x)$

IT0055

(d) $3\cos^{-1} x = \sin^{-1}(\sqrt{1-x^2}(4x^2-1))$

IT0056

(e) $\sin^{-1} x + \sin^{-1} y = \frac{2\pi}{3}$ & $\cos^{-1} x - \cos^{-1} y = \frac{\pi}{3}$

IT0057

15. Find the sum of the series :

(a) $\cot^{-1}7 + \cot^{-1}13 + \cot^{-1}21 + \cot^{-1}31 + \dots$ to n terms.

IT0058

(b) $\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{2}{9} + \dots + \tan^{-1} \frac{2^{n-1}}{1+2^{2n-1}} + \dots \infty$

IT0059

(c) $\tan^{-1} \frac{1}{x^2+x+1} + \tan^{-1} \frac{1}{x^2+3x+3} + \tan^{-1} \frac{1}{x^2+5x+7} + \tan^{-1} \frac{1}{x^2+7x+13}$ to n terms.

IT0060

(d) $\sin^{-1} \frac{1}{\sqrt{5}} + \sin^{-1} \frac{1}{\sqrt{65}} + \sin^{-1} \frac{1}{\sqrt{325}} + \dots + \sin^{-1} \frac{1}{\sqrt{4n^4+1}} + \dots \infty$ terms.

IT0061

EXERCISE (S-2)

1. If $\alpha = 2 \arctan\left(\frac{1+x}{1-x}\right)$ & $\beta = \arcsin\left(\frac{1-x^2}{1+x^2}\right)$ for $0 < x < 1$, then prove that $\alpha + \beta = \pi$, what the value of $\alpha + \beta$ will be if $x > 1$. IT0062
2. Solve the following : IT0063
 - (a) $\tan^{-1}\frac{x-1}{x+1} + \tan^{-1}\frac{2x-1}{2x+1} = \tan^{-1}\frac{23}{36}$ IT0064
 - (b) $2 \tan^{-1} x = \cos^{-1}\frac{1-a^2}{1+a^2} - \cos^{-1}\frac{1-b^2}{1+b^2}$ ($a > 0, b > 0$) IT0065
3. If α and β are the roots of the equation $x^2 - 4x + 1 = 0$ ($\alpha > \beta$) then find the value of $f(\alpha, \beta) = \frac{\beta^3}{2} \operatorname{cosec}^2\left(\frac{1}{2} \tan^{-1} \frac{\beta}{\alpha}\right) + \frac{\alpha^3}{2} \sec^2\left(\frac{1}{2} \tan^{-1} \frac{\alpha}{\beta}\right)$. IT0066
4. Find all the positive integral solutions of, $\tan^{-1}x + \cos^{-1}\frac{y}{\sqrt{1+y^2}} = \sin^{-1}\frac{3}{\sqrt{10}}$. IT0067
5. Prove that the equation, $(\sin^{-1}x)^3 + (\cos^{-1}x)^3 = \alpha\pi^3$ has no roots for $\alpha < \frac{1}{32}$ and $\alpha > \frac{7}{8}$. IT0068
6. Solve the following inequalities : IT0069
 - (a) $\arccot^2 x - 5 \arccot x + 6 > 0$ IT0070
 - (b) $\arcsin x > \arccos x$
 - (c) $\tan^2(\arcsin x) > 1$
7. Solve the following system of inequalities : IT0071

$$4 \arctan^2 x - 8 \arctan x + 3 < 0 \quad \& \quad 4 \arccot x - \arccot^2 x - 3 \geq 0$$
8. If the sum $\sum_{n=1}^{10} \sum_{m=1}^{10} \tan^{-1}\left(\frac{m}{n}\right) = k\pi$, find the value of k . IT0072
9. Show that the roots r, s and t of the cubic $x(x-2)(3x-7) = 2$, are real and positive. Also compute the value of $\tan^{-1}(r) + \tan^{-1}(s) + \tan^{-1}(t)$. IT0073
10. Solve for x : $\sin^{-1}\left(\sin\left(\frac{2x^2+4}{1+x^2}\right)\right) < \pi - 3$. IT0074
11. Find the set of values of 'a' for which the equation $2\cos^{-1}x = a + a^2(\cos^{-1}x)^{-1}$ posses a solution. IT0075

EXERCISE (JM)

1. Let $f : (-1, 1) \rightarrow B$, be a function defined by $f(x) = \tan^{-1} \frac{2x}{1-x^2}$, then f is both one-one and onto when

B is the interval-

[AIEEE-2005]

- (1) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (2) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (3) $\left(0, \frac{\pi}{2}\right)$ (4) $\left[0, \frac{\pi}{2}\right)$

IT0076

2. If $\cos^{-1}x - \cos^{-1}\frac{y}{2} = \alpha$, then $4x^2 - 4xy \cos \alpha + y^2$ is equal to -

[AIEEE-2005]

- (1) $2 \sin 2\alpha$ (2) 4 (3) $4 \sin^2 \alpha$ (4) $-4 \sin^2 \alpha$

IT0077

3. If $\sin^{-1}\left(\frac{x}{5}\right) + \operatorname{cosec}^{-1}\left(\frac{5}{4}\right) = \frac{\pi}{2}$, then a value of x is-

[AIEEE-2007]

- (1) 1 (2) 3 (3) 4 (4) 5

IT0078

4. The value of $\cot\left(\operatorname{cosec}^{-1}\frac{5}{3} + \tan^{-1}\frac{2}{3}\right)$ is equal to-

[AIEEE-2008]

- (1) $\frac{6}{17}$ (2) $\frac{3}{17}$ (3) $\frac{4}{17}$ (4) $\frac{5}{17}$

IT0079

5. If x, y, z are in A.P. and $\tan^{-1}x, \tan^{-1}y$ and $\tan^{-1}z$ are also in A.P., then

[JEE (Main)-2013]

- (1) $x = y = z$ (2) $2x = 3y = 6z$ (3) $6x = 3y = 2z$ (4) $6x = 4y = 3z$

IT0080

6. Let $\tan^{-1}y = \tan^{-1}x + \tan^{-1}\left(\frac{2x}{1-x^2}\right)$, where $|x| < \frac{1}{\sqrt{3}}$. Then a value of y is : [JEE (Main)-2015]

- (1) $\frac{3x-x^3}{1+3x^2}$ (2) $\frac{3x+x^3}{1+3x^2}$ (3) $\frac{3x-x^3}{1-3x^2}$ (4) $\frac{3x+x^3}{1-3x^2}$

IT0081

7. If $\cos^{-1}\left(\frac{2}{3x}\right) + \cos^{-1}\left(\frac{3}{4x}\right) = \frac{\pi}{2}$ ($x > \frac{3}{4}$), then x is equal to :

[JEE(Main)-2019]

- (1) $\frac{\sqrt{145}}{12}$ (2) $\frac{\sqrt{145}}{10}$ (3) $\frac{\sqrt{146}}{12}$ (4) $\frac{\sqrt{145}}{11}$

IT0082

8. If $x = \sin^{-1}(\sin 10)$ and $y = \cos^{-1}(\cos 10)$, then $y - x$ is equal to: [JEE(Main)-2019]
- (1) π (2) 7π (3) 0 (4) 10

IT0083

9. The value of $\cot\left(\sum_{n=1}^{19} \cot^{-1}\left(1 + \sum_{p=1}^n 2p\right)\right)$ is: [JEE(Main)-2019]
- (1) $\frac{22}{23}$ (2) $\frac{23}{22}$ (3) $\frac{21}{19}$ (4) $\frac{19}{21}$

IT0084

10. All x satisfying the inequality $(\cot^{-1} x)^2 - 7(\cot^{-1} x) + 10 > 0$, lie in the interval :- [JEE(Main)-2019]
- (1) $(-\infty, \cot 5) \cup (\cot 4, \cot 2)$ (2) $(\cot 5, \cot 4)$
 (3) $(\cot 2, \infty)$ (4) $(-\infty, \cot 5) \cup (\cot 2, \infty)$

IT0085

11. Considering only the principal values of inverse functions, the set $A = \left\{x \geq 0 : \tan^{-1}(2x) + \tan^{-1}(3x) = \frac{\pi}{4}\right\}$ [JEE(Main)-2019]
- (1) is an empty set (2) Contains more than two elements
 (3) Contains two elements (4) is a singleton

IT0086

12. If $\alpha = \cos^{-1}\left(\frac{3}{5}\right)$, $\beta = \tan^{-1}\left(\frac{1}{3}\right)$, where $0 < \alpha, \beta < \frac{\pi}{2}$, then $\alpha - \beta$ is equal to: [JEE(Main)-2019]
- (1) $\sin^{-1}\left(\frac{9}{5\sqrt{10}}\right)$ (2) $\tan^{-1}\left(\frac{9}{14}\right)$ (3) $\cos^{-1}\left(\frac{9}{5\sqrt{10}}\right)$ (4) $\tan^{-1}\left(\frac{9}{5\sqrt{10}}\right)$

IT0087

13. If $\cos^{-1}x - \cos^{-1}\frac{y}{2} = \alpha$, where $-1 \leq x \leq 1$, $-2 \leq y \leq 2$, $x \leq \frac{y}{2}$, then for all x, y , $4x^2 - 4xy \cos \alpha + y^2$ is equal to [JEE(Main)-2019]
- (1) $4 \sin^2 \alpha - 2x^2y^2$ (2) $4 \cos^2 \alpha + 2x^2y^2$ (3) $4 \sin^2 \alpha$ (4) $2 \sin^2 \alpha$

IT0088

14. The value of $\sin^{-1}\left(\frac{12}{13}\right) - \sin^{-1}\left(\frac{3}{5}\right)$ is equal to : [JEE(Main)-2019]

(1) $\pi - \sin^{-1}\left(\frac{63}{65}\right)$ (2) $\pi - \cos^{-1}\left(\frac{33}{65}\right)$ (3) $\frac{\pi}{2} - \sin^{-1}\left(\frac{56}{65}\right)$ (4) $\frac{\pi}{2} - \cos^{-1}\left(\frac{9}{65}\right)$

IT0089

EXERCISE (JA)

1. Let (x, y) be such that $\sin^{-1}(ax) + \cos^{-1}(y) + \cos^{-1}(bxy) = \frac{\pi}{2}$ [JEE 2007, 6]

Match the statements in column-I with statements in column-II and indicate your answer by darkening the appropriate bubbles in the 4×4 matrix given in the ORS.

Column-I

- (A) If $a = 1$ and $b = 0$, then (x, y)
(B) If $a = 1$ and $b = 1$, then (x, y)
(C) If $a = 1$ and $b = 2$, then (x, y)
(D) If $a = 2$ and $b = 2$, then (x, y)

Column-II

- (p) lies on the circle $x^2 + y^2 = 1$
(q) lies on $(x^2 - 1)(y^2 - 1) = 0$
(r) lies on $y = x$
(s) lies on $(4x^2 - 1)(y^2 - 1) = 0$

IT0090

2. If $0 < x < 1$, then $\sqrt{1+x^2} [\{x \cos(\cot^{-1}x) + \sin(\cot^{-1}x)\}^2 - 1]^{1/2} =$ [JEE 2008, 3]

(A) $\frac{x}{\sqrt{1+x^2}}$ (B) x (C) $x\sqrt{1+x^2}$ (D) $\sqrt{1+x^2}$

IT0091

3. The value of $\cot\left(\sum_{n=1}^{23} \cot^{-1}\left(1 + \sum_{k=1}^n 2k\right)\right)$ is [JEE(Advanced) 2013, 2]

(A) $\frac{23}{25}$ (B) $\frac{25}{23}$ (C) $\frac{23}{24}$ (D) $\frac{24}{23}$

IT0092

4. Match List-I with List-II and select the correct answer using the code given below the lists.

List-I**List-II**

P. $\left(\frac{1}{y^2} \left(\frac{\cos(\tan^{-1} y) + y \sin(\tan^{-1} y)}{\cot(\sin^{-1} y) + \tan(\sin^{-1} y)} \right)^2 + y^4 \right)^{1/2}$ takes value

1. $\frac{1}{2}\sqrt{\frac{5}{3}}$

Q. If $\cos x + \cos y + \cos z = 0 = \sin x + \sin y + \sin z$,

2. $\sqrt{2}$

then possible value of $\cos \frac{x-y}{2}$ is

R. If $\cos\left(\frac{\pi}{4} - x\right) \cos 2x + \sin x \sin 2x \sec x$

3. $\frac{1}{2}$

$= \cos x \sin 2x \sec x + \cos\left(\frac{\pi}{4} + x\right) \cos 2x$,

then possible value of $\sec x$ is

S. If $\cot\left(\sin^{-1} \sqrt{1-x^2}\right) = \sin\left(\tan^{-1}(x\sqrt{6})\right)$, $x \neq 0$,

4. 1

then possible value of x is

Codes :

	P	Q	R	S
(A)	4	3	1	2
(B)	4	3	2	1
(C)	3	4	2	1
(D)	3	4	1	2

[JEE(Advanced) 2013, 3, (-1)]

IT0093

5. Let $f : [0, 4\pi] \rightarrow [0, \pi]$ be defined by $f(x) = \cos^{-1}(\cos x)$. The number of points $x \in [0, 4\pi]$ satisfying

the equation $f(x) = \frac{10-x}{10}$ is

[JEE(Advanced)-2014, 3]

IT0094

6. If $\alpha = 3 \sin^{-1}\left(\frac{6}{11}\right)$ and $\beta = 3 \cos^{-1}\left(\frac{4}{9}\right)$ where the inverse trigonometric functions take only the principal values, then the correct option(s) is(are)

[JEE(Advanced)-2015, 4]

(A) $\cos \beta > 0$ (B) $\sin \beta < 0$ (C) $\cos(\alpha + \beta) > 0$ (D) $\cos \alpha < 0$

IT0095

7. The number of real solutions of the equation

$$\sin^{-1} \left(\sum_{i=1}^{\infty} x^{i+1} - x \sum_{i=1}^{\infty} \left(\frac{x}{2} \right)^i \right) = \frac{\pi}{2} - \cos^{-1} \left(\sum_{i=1}^{\infty} \left(-\frac{x}{2} \right)^i - \sum_{i=1}^{\infty} (-x)^i \right) \text{ lying in the interval } \left(-\frac{1}{2}, \frac{1}{2} \right) \text{ is } \underline{\hspace{2cm}}$$

(Here, the inverse trigonometric functions $\sin^{-1}x$ and $\cos^{-1}x$ assume value in $\left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$ and $[0, \pi]$, respectively.)

[JEE(Advanced)-2018]
IT0096

8. Let $E_1 = \left\{ x \in \mathbb{R} : x \neq 1 \text{ and } \frac{x}{x-1} > 0 \right\}$ and $E_2 = \left\{ x \in E_1 : \sin^{-1} \left(\log_e \left(\frac{x}{x-1} \right) \right) \text{ is a real number} \right\}$.

(Here, the inverse trigonometric function $\sin^{-1}x$ assumes values in $\left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$.)

Let $f : E_1 \rightarrow \mathbb{R}$ be the function defined by $f(x) = \log_e \left(\frac{x}{x-1} \right)$

and $g : E_2 \rightarrow \mathbb{R}$ be the function defined by $g(x) = \sin^{-1} \left(\log_e \left(\frac{x}{x-1} \right) \right)$.

LIST-I

- P. The range of f is
Q. The range of g contains
R. The domain of f contains
S. The domain of g is

LIST-II

1. $\left(-\infty, \frac{1}{1-e} \right] \cup \left[\frac{e}{e-1}, \infty \right)$
2. $(0, 1)$
3. $\left[-\frac{1}{2}, \frac{1}{2} \right]$
4. $(-\infty, 0) \cup (0, \infty)$
5. $\left(-\infty, \frac{e}{e-1} \right]$
6. $(-\infty, 0) \cup \left(\frac{1}{2}, \frac{e}{e-1} \right]$

The correct option is :

- (A) $P \rightarrow 4; Q \rightarrow 2; R \rightarrow 1; S \rightarrow 1$
(C) $P \rightarrow 4; Q \rightarrow 2; R \rightarrow 1; S \rightarrow 6$

- (B) $P \rightarrow 3; Q \rightarrow 3; R \rightarrow 6; S \rightarrow 5$
(D) $P \rightarrow 4; Q \rightarrow 3; R \rightarrow 6; S \rightarrow 5$

[JEE(Advanced)-2018]

IT0097

9. The value of $\sec^{-1} \left(\frac{1}{4} \sum_{k=0}^{10} \sec \left(\frac{7\pi}{12} + \frac{k\pi}{2} \right) \sec \left(\frac{7\pi}{12} + \frac{(k+1)\pi}{2} \right) \right)$ in the interval $\left[-\frac{\pi}{4}, \frac{3\pi}{4} \right]$ equals

[JEE(Advanced)-2019, 3(0)]
IT0098

ANSWER KEY

INVERSE TRIGONOMETRIC FUNCTION

EXERCISE (O-1)





1. C 2. B 3. D 4. B 5. B 6. C 7. A
 8. A 9. A 10. B,C 11. A,B,C,D 12. B,D 13. B,C
 14. B,C,D 15. A,B

EXERCISE (O-2)

1. A 2. C 3. C 4. A 5. B 6. A,D 7. A,C
 8. A,B,D 9. D 10. C 11. A 12. B 13. 1 14. 2
 15. 9 16. 1

EXERCISE (S-1)

1. 5
 2. (a) (i) $\frac{1}{\sqrt{3}}$, (ii) $\frac{5\pi}{6}$, (iii) $\frac{4}{5}$, (iv) $\frac{17}{6}$; (b) (i) $\frac{1}{2}$, (ii) -1 , (iii) $-\frac{\pi}{4}$, (iv) $\frac{\sqrt{2}}{4}$
 3. (i) $-1/3 \leq x \leq 1$ (ii) $\{1, -1\}$ (iii) $1 \leq x < 4$ (iv) $[-(1+\sqrt{2}), (\sqrt{2}-1)]$
 (v) $x \in (-1/2, 1/2), x \neq 0$ (vi) $(3/2, 2]$ (vii) $(-2, 2) - \{-1, 0, 1\}$



5. 53 8. 10 9. π 10. $k = \frac{11}{4}$ 11. (a) $\frac{\pi}{3}$; (b) $\frac{1}{2} \tan^{-1} x$
 13. (a) $(-\infty, \sec 2) \cup [1, \infty)$
 14. (a) $x = \frac{1}{2} \sqrt{\frac{3}{7}}$; (b) $x = 3$; (c) $x = 0, \frac{1}{2}, -\frac{1}{2}$; (d) $\left[\frac{\sqrt{3}}{2}, 1\right]$; (e) $x = \frac{1}{2}, y = 1$
 15. (a) $\operatorname{arccot} \left[\frac{2n+5}{n} \right]$, (b) $\frac{\pi}{4}$, (c) $\arctan(x+n) - \arctan x$, (d) $\frac{\pi}{4}$

EXERCISE (S-2)

1. $-\pi$
2. (a) $x = \frac{4}{3}$; (b) $x = \frac{a-b}{1+ab}$
3. 56
4. $x = 1; y = 2$ & $x = 2; y = 7$
6. (a) $(\cot 2, \infty) \cup (-\infty, \cot 3)$ (b) $\left(\frac{\sqrt{2}}{2}, 1\right]$ (c) $\left(\frac{\sqrt{2}}{2}, 1\right) \cup \left(-1, -\frac{\sqrt{2}}{2}\right)$
7. $\left(\tan \frac{1}{2}, \cot 1\right]$
8. $k = 25$
9. $\frac{3\pi}{4}$
10. $x \in (-1, 1)$
11. $a \in [-2\pi, \pi] - \{0\}$

EXERCISE (JM)

1. 2
2. 3
3. 2
4. 1
5. 1
6. 3
7. 1
8. 1
9. 3
10. 3
11. 4
12. 1
13. 3
14. 3

EXERCISE (JA)

1. (A) P; (B) Q; (C) P; (D) S
2. C
3. B
4. B
5. 3
6. B, C, D
7. 2
8. A
9. 0.00