

Dr. Vainav Patel 'Scientist F' Viral Immunopathogenesis Lab Email: patelv@nirrh.res.in



Volume 111, Issue 6 June 2022

JOURNAL ARTICLE


Opposing roles for sMAdCAM and IL-15 in COVID-19 associated cellular immune pathology Get access >

Amit Kumar Singh, Nandini Kasarpalkar, Shilpa Bhowmick, Gaurav Paradkar, Mayur Talreja, Karan Shah, Abhishek Tiwari, Harsha Palav, Snehal Kaginkar, Rajiv Kulkarni, Ashwini Patil, Varsha Kalsurkar, Sachee Agrawal, Jayanthi Shastri, Rajesh Dere, Ramesh Bharmal, Smita D Mahale, Vikrant M Bhor 丞, Vainav Patel ☒

Journal of Leukocyte Biology, Volume 111, Issue 6, June 2022, Pages 1287–1295,

https://doi.org/10.1002/JLB.3COVBCR0621-300R


Published: 24 January 2022 Article history ▼



- Lymphopenia negatively correlated with sMAdCAM and positively with IL-15
- Elevated intermediate monocyte frequency (pan disease signature)
- Sparing of Treg and CD8 Effector Memory subsets
- Lower IL-15 corresponds to shorter hospitalization


Integrated viral and immune monitoring in a prospective COVID-19 cohort from India






Findings

- 51 participants received at least one indigenous vaccine dose (Covishield/Covaxin)
- Non-survivors did not restore circulating B and NK cells and were largely unvaccinated
- Delta variants of SARS-CoV-2 clearly demonstrated increased pathogenicity





Non-suvivors




Fig 3 : **Distribution of absolute counts** different immune cell subsets study participants: (A-E) among variation across asymptomatic (AS), mild symptomatic (MS), HP-S, and HP-NS individuals at day 0 (D0); (F–J) longitudinal follow-up of immune subsets from D0 to D90 at specified timepoints. Timepoints in HP-NS individuals were curtailed due to mortality in this group by D15. (K–O) Variation across AS,MS, HP-S, and HP-NS individuals at D7 when highest apparent divergence and maximum availability of follow-up data across available. Statistical groups were significance calculated was by nonparametric1-way analysis variance (Kruskal-Wallis test) followed by post hoc Dunn's test.*P < 0.05; **P < 0.01,***P < 0.001; ****P < 0.001.




Fig 5: Comparative analysis of absolute no. of different immune cell subsets between groups infected by Delta vs non-Delta subtypes: (A–F) Variation of mentioned immune subsets at day 0. (G–L) Variation of mentioned immune subsets at day 7. Black datapoints indicate survivors and red datapoints indicate non survivors. Statistical significance was calculated by Mann-Whitney U test.*P < 0.05; **P < 0.01; ***P < 0.001.



RESPONSE to NIPAH 2023 outbreak

15th **September, 2023 (Morning):** Request from Dr. Nivedita Gupta, Chief, ECD, ICMR and Dr. Pragya Yadav, Group Leader, Maximum Containment Laboratory, ICMR- NIV, Pune regarding need to understand the viral pathology and associated immune signatures

Decision taken to conduct immune monitoring studies on samples from Nipah exposed/infected individuals at ICMR-NIV (MCC), Pune BSL-4 Facility by Dr. Vainav Patel and team (Ms. Harsha Palav (Ph.D. Scholar); Mr. Nandan Mohite (Ph.D. Scholar); Mr. Pranay Gurav (Research Assistant); Ms. Denna Sarah (FACS Operator) in collaboration with Dr. Pragya Yadav and team

17th **September, 2023:** The team from ICMR-NIRRCH left from Mumbai to ICMR-NIV, Pune. Protocol for sample collection, safe transport, sample processing and storage created and finalized including **PAPR** training with all protective apparel was conducted.

18th **September, 2023:** Commencement of Immune monitoring: Panels used: **Absolute count panel (T, B, NK, Mono, Grans)**

T-cell Activation & Treg panel Plasma blast & homing panel





PAPR: Powered Air Purifying Respirator training



ICMR- NIRRCH NIPAH TEAM

19th **September 2023:** Blood samples from 2 infected individuals were received and processed in BLS-4 facility and successfully subjected to ex-vivo immunophenotyping; PBMC and plasma was stored for further analysis.



received and processed alongside a healthy sample.

27th **September 2023 & 13**th **October 2023:** Follow-up samples received and processed along with healthy samples

Work flow Whole blood collection 3 - 8ml in EDTA III. PBMC I. Ex-vivo whole blood isolation immune monitoring II. Plasma separation **Cryopreserved in** Liq. Nitro Plasmablast **Absolute** T cell activation and B cell panel count and Treg panel Aliquoted and Stored at -80°C

Preliminary results

Increased Low Density Granulocytes (LDGs) & Plasmablasts (PB) observed in patients




Figure S1. Representative flow cytometric gating strategy for lymphocytes, monocytes, neutrophils, eosinophils, NK cells, NKT like cells and subsets within these cells.




Figure S2. Representative flow cytometric gating strategy for CD4+ and CD8+ T cell subsets.






Figure S3. Representative flow cytometric gating strategy for detection of plasmablast and expression of integrin Beta-7 within different cell population.



JOURNAL OF MEDICAL VIROLOGY



Volume 96, Issue 9 September 2024 e29920

Comparative immune profiling in survivors of the 2023 Nipah outbreak in Kerala state, India

Rima R. Sahay, Harsha C. Palav, Anita M. Shete, Deepak Y. Patil, Sreelekshmy Mohandas, Chandni Radhakrishnan, P. Shihabudheen, Anoop Kumar, Anitha Puduvail Moorkoth, Nandan Mohite, Pranay Gurav, Niyas K. Pullor, Rajlaxmi Jain, Yash Joshi, Lathika Velichapat Ramakrishnan, Nivedita Gupta, Vainav Patel , Pragya D. Yadav ... See fewer authors

First published: 16 September 2024 | https://doi.org/10.1002/jmv.29920