| Name: | DOB: | Embrace No. | NHS No. | Weight: |
|---|---|---|---|---|

| Drug | | Dose range | Dose prescribed | Route | Prescribers signature | Date/time given | Checked/ given by | Batch No. |
|---|---|---|---|---|---|---|---|---|
| **INDUCTION AGENTS** | | | | | | | | |
| | **Dilute to:** | | | | | | | |
| **Ketamine** | 10 mg/ml | 1 - 2 mg/kg | | | | | | |
| **Rocuronium** | 10 mg/ml | 1 mg/kg | | | | | | |
| Fentanyl | 10 micrograms/ml | 2 - 5 micrograms/kg | | | | | | |
| Atropine | 600 micrograms/ml | 20 micrograms/kg<br>(min 100 micrograms, **MAX 600 micrograms**) | | | | | | |
| Propofol | 1% = 10 mg/ml | 1 - 4 mg/kg<br>(**CAUTION** with cardiovascular instability) | | | | | | |
| Thiopentone | 25 mg/ml | 1 - 5 mg/kg | | | | | | |
| **REVERSAL AGENTS** | | | | | | | | |
| Naloxone **Under 12 years**<br>(emergency opiod overdose) | | 100 micrograms/kg (**MAX 2 mg**) | | | | | | |
| Naloxone **12 years+**<br>(emergency opiod overdose) | | 400 micrograms then 800 micrograms at 1 minute intervals for up to 2 doses then 2 mg for 1 dose | | | | | | |
| Sugammadex<br>(for rocuronium) | | 16 mg/kg | | | | | | |
| Neostigmine/Glycopyrolate<br>(for atracurium/rocuronium) | | 0.02 ml/kg<br>**(MAX 2 ml per course)** | | | | | | |
| **RESUSCITATION DRUGS** | | | | | | | | |
| Adrenaline 1:10,000 | | 0.1 ml/kg = 10 micrograms/kg **MAX 1 mg** | | | | | | |
| Calcium Gluconate 10% | | 0.5 ml/kg **MAX 20 ml** | | | | | | |
| Sodium Bicarbonate 8.4% | | 1 ml/kg (1 mmol/kg) | | | | | | |
| Glucose 10% | | 2 ml/kg | | | | | | |
| Lorazepam | | 0.1 mg/kg **MAX 4 mg** | | | | | | |
| 2.7% Sodium Chloride | | 3 - 5 ml/kg over 30 mins | | | | | | |
| Mannitol 20% | | 2.5 ml/kg (500 mg/kg) over 30 mins | | | | | | |
| Amiodarone (for cardiac arrest) | | 5 mg/kg over 3 minutes<br>**MAX 300 mg -** use dextrose flush | | | | | | |
| Atropine | | 20 micrograms/kg<br>(min 100 micrograms, **MAX 600 micrograms)** | | | | | | |

Embrace
Yorkshire & Humber Infant & Children's Transport Service

# Embrace Paediatric Prescription Chart

Sheffield Children's NHS
NHS Foundation Trust

**Patient name:**

**DOB:** **Weight:**

**Age:**

**Corrected gestational age:**
**(If relevant)**

**NHS number:**

**Embrace number:**

**ALLERGY and ADVERSE drugs reaction status** **Unconfirmed allergy status ☐**

| Medicine/ Substance | Description of allergy/ reaction | Signature | Print Name | Date |
|---|---|---|---|---|
| | | | | |

**Prescriber information/stamp:**

**IF PATIENT HAS BMI >98th CENTILE PLEASE CONSIDER:**
**IDEAL BODY WEIGHT PRESCRIBING FOR:** ADRENALINE, PROPOFOL, THIOPENTONE, KETAMINE, MIDAZOLAM, FENTANYL, MORPHINE
**ADJUSTED BODY WEIGHT PRESCRIBING FOR:** ROCURONIUM AND ATRACURIUM

| Date / Time | Drug / Fluid bolus | Dose | Dose/Kg | Route | Over time | Signature | Date/time given | Checked/given by | Batch No. |
|---|---|---|---|---|---|---|---|---|---|
| | **Paracetamol<br>(1 month/44 wks CGA - 18 yrs)** | | **15 mg/kg<br>MAX 1 gram** | **IV** | **15 mins** | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |
| | | | | | | | | | |

| Date / Time | Infusion fluid | Batch No. | Total volume | Rate of infusion | Drugs added | | | Signature | Date/time started | Checked/ given by |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Drug(s) | Dose added | Batch No. | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |

| Name: | DOB: | Embrace No. | NHS No. | Weight: |
|---|---|---|---|---|
| | | | | |

| Sedatives/Paralytics and miscellaneous | Concentration | Amount in syringe | Dose range | Infusion rate | Prescribers signature | Date/time started | Checked/ given by | Batch No. |
|---|---|---|---|---|---|---|---|---|
| Dinoprostone | 6 micrograms/ml | 300 micrograms in 50ml total Glu 5% | 5 - 100 nanograms/kg/min | 0.1 ml/kg/hr = 10 nanograms/kg/min | | | | |
| | *FOLLOW ODN GUIDELINE OR CARDIOLOGY ADVICE* (use neonatal profile on pump) | | | | | | | |
| Morphine **Under 50kg** | 1 mg/kg (max 50 mg) | =____**mg** in 50 ml total Glu 5% or NaCl 0.9% | 20 - 40 micrograms/kg/hr | 1 ml/hr = 20 micrograms/kg/hr | | | | |
| Morphine **50kg+** | 1 mg/ml | 50 mg in 50 ml total Glu 5% or NaCl 0.9% | 20 – 40 micrograms/kg/hr | 0.02 ml/kg/hr = 20 micrograms/kg/hr | | | | |
| Midazolam **Under 40kg** | 6 mg/kg | =____**mg** in 50 ml total Glu 5% or NaCl 0.9% | 120 - 360 micrograms/kg/hr | 1 ml/hr = 120 micrograms/kg/hr | | | | |
| Midazolam **40kg+** | 3 mg/kg | =____**mg** in 50 ml total Glu 5% or NaCl 0.9% | 120 - 360 micrograms/kg/hr | 2 ml/hr = 120 micrograms/kg/hr | | | | |
| Ketamine | 1 mg/ml | 50 mg in 50 ml total Glu 5% or NaCl 0.9% | 10 - 45 micrograms/kg/min | 0.6 ml/kg/hr = 10 micrograms/kg/min | | | | |
| Propofol 1% | 10 mg/ml **NEAT** | 500 mg in 50 ml | 0.3 - 4 mg/kg/hr | 0.1 ml/kg/hr = 1 mg/kg/hr | | | | |
| Rocuronium **Under 10kg** | 2 mg/ml | 50 mg in **25 ml** total Glu 5% or NaCl 0.9% | 300 - 600 micrograms/kg/hr | 0.15 ml/kg/hr = 300 micrograms/kg/hr | | | | |
| Rocuronium **10kg+** | 10 mg/ml **NEAT** | 250 mg in **25 ml** | 300 - 600 micrograms/kg/hr | 0.03 ml/kg/hr = 300 micrograms/kg/hr | | | | |
| Atracurium **Under 10kg** | 5 mg/ml | 250 mg in 50 ml total Glu 5% or NaCl 0.9% | 0.5 - 1 mg/kg/hr | 0.1 ml/kg/hr = 0.5 mg/kg/hr | | | | |
| Atracurium **10kg+** | 10 mg/ml **NEAT** | 250 mg in **25 ml** | 0.5 - 1 mg/kg/hr | 0.1 ml/kg/hr = 1 mg/kg/hr | | | | |
| Heparin (arterial line) | 1 unit/ml | 50 units in 50 ml total NaCl 0.9% | 0.5 - 2 unit/hr | 0.5 - 2 ml/hr | | | | |

| Age | Weight (kg) | ETT size | At lips (cm) | At nose (cm) | Heart rate | Resp. rate | Systolic BP 5th centile | Systolic BP 50th centile |
|---|---|---|---|---|---|---|---|---|
| **1 month** | 4.5 | 3.5 | 9 | 11 | 120 - 170 | 25 - 50 | 65 - 75 | 80 - 90 |
| **3 month** | 6 | 3.5 - 4.0 | 10 | 12 | 115 - 165 | 25 - 45 | 65 - 75 | 80 - 90 |
| **6 month** | 8 | 4.0 | 12 | 14 | 110 - 160 | 20 - 40 | 65 - 75 | 80 - 90 |
| **1 years** | 9.5 | 4.0 - 4.5 | 13 | 15 | 110 - 160 | 20 - 40 | 70 - 75 | 85 - 95 |
| **2 years** | 12 | 4.5 | 13 | 15 | 100 - 150 | 20 - 30 | 70 - 80 | 85 - 100 |
| **3 years** | 14 | 5.0 | 14 | 16 | 90 - 140 | 20 - 30 | 70 - 80 | 85 - 100 |
| **4 years** | 16 | 5.0 | 14 | 17 | 80 - 135 | 20 - 30 | 80 - 90 | 85 - 100 |
| **6 years** | 20 | 5.5 | 15 | 19 | 80 - 130 | 20 - 30 | 80 - 90 | 90 - 110 |
| **8 years** | 25 | 6.0 | 16 | 20 | 70 - 120 | 15 - 25 | 80 - 90 | 90 - 110 |
| **10 years** | 32 | 6.5 | 17 | 21 | 70 - 120 | 15 - 25 | 80 - 90 | 90 - 110 |
| **12 years** | 43 | 7.0 - 7.5 | 18 | 22 | 65 - 115 | 12 - 24 | 90 - 105 | 100 - 120 |
| **14 years** | 50 | 7.5 - 8.0 | 21 | 25 | 60 - 110 | 12 - 24 | 90 - 105 | 100 - 120 |
| **Adult** | 60 - 70 | 8.0 - 9.0 | 21 - 24 | 25 - 28 | 60 - 110 | 12 - 24 | 90 - 105 | 100 - 120 |

| Maintenance fluids | |
|---|---|
| 100ml/kg for 1st 10 kg | |
| 50ml/kg for 2nd 10kg | |
| 20ml/kg for each subsequent kg | |
| Total required at 100% for 24hrs | |
| Total in 24hrs in ml/hr | |
| | |
| Amount to be given % | |
| Hourly rate required (ml) | |

**>65kg IV Fluids = 2400ml/day**
**Maximum Fluid Bolus per dose = 1000ml**
**Usually include infusion volumes in overall total**
**Consider neonatal fluid calculation for < 4 weeks old**

**ANAPHYLAXIS**
**IM ADRENALINE 1:1000**

| Age | Dose |
|---|---|
| Less than 6 months | 100 - 150 micrograms (0.1 - 0.15ml) |
| 6 months to 6 years | 150 micrograms (0.15ml) |
| 6 to 12 years | 300 micrograms (0.3ml) |
| More than 12 years | 500 micrograms (0.5ml) |

| Name: | DOB: | Embrace No. | NHS No. | Weight: |
|---|---|---|---|---|
| | | | | |

| PUSH PRESSORS (PERIPHERAL OR CENTRAL) Clearly label as dilute | Drug | Concentration | Amount in syringe | Dose range | Dose required (ml) | Prescribers signature | Date/time started | Checked/ given by | Batch No. | Total volume given |
|---|---|---|---|---|---|---|---|---|---|---|
| | Adrenaline **(dilute bolus)** | 10 micrograms/ml | 100 micrograms (1 ml of 1:10,000) in 10ml total NaCl 0.9% | 1 microgram/kg (0.1ml/kg) **MAX 20 micrograms** | | | | | | |
| | Metaraminol | 100 micrograms/ml | 1 mg in 10ml total NaCl 0.9% | 10 micrograms/kg (0.1ml/kg) **MAX 500 micrograms** | | | | | | |

| Vasoactive agents | Concentration | Amount in syringe | Dose range | Infusion rate | Prescribers signature | Date/time started | Checked/ given by | Batch No. |
|---|---|---|---|---|---|---|---|---|
| Adrenaline **PERIPHERAL** | 20 micrograms/ml | 1 mg in 50 ml total NaCl 0.9% | 0.1 – 1.5 micrograms/kg/min | 0.3 ml/kg/hr = 0.1 micrograms/kg/min | | | | |
| Adrenaline **CENTRAL Under 50kg** | 300 micrograms/kg | =____**mg** in 50 ml total Glu 5% or NaCl 0.9% | 0.1 – 1.5 micrograms/kg/min | 1 ml/hr = 0.1 micrograms/kg/min | | | | |
| Adrenaline **CENTRAL 50kg+** | 320 micrograms/ml | 16 mg in 50 ml total Glu 5% or NaCl 0.9% | 0.1 – 1.5 micrograms/kg/min | 0.018 ml/kg/hr = 0.1 micrograms/kg/min | | | | |
| Dobutamine **PERIPHERAL** | 3 mg/kg (**MAX** 250 mg in 50 ml) | =____**mg** in 50 ml total Glu 5% or NaCl 0.9% | 2 – 20 micrograms/kg/min | 2 ml/hr = 2 micrograms/kg/min | | | | |
| Dobutamine **CENTRAL Under 40kg** | 15 mg/kg | =____**mg** in 50 ml total Glu 5% or NaCl 0.9% | 2 – 20 micrograms/kg/min | 1 ml/hr = 5 micrograms/kg/min | | | | |
| Dobutamine **CENTRAL 40kg+** | 12.5 mg/ml | 625 mg in 50 ml Neat | 2 – 20 micrograms/kg/min | 0.024 ml/kg/hr = 5 micrograms/kg/min | | | | |
| Noradrenaline **PERIPHERAL** | 20 micrograms/ml | 1 mg in 50 ml total Glu 5% | 0.02 – 1 micrograms/kg/min | 0.3 ml/kg/hr = 0.1 micrograms/kg/min | | | | |
| Noradrenaline **CENTRAL Under 50kg** | 300 micrograms/kg | =____**mg** in 50 ml total Glu 5% | 0.02 – 1 micrograms/kg/min | 1 ml/hr = 0.1 micrograms/kg/min | | | | |
| Noradrenaline **CENTRAL 50kg+** | 320 micrograms/ml | 16 mg in 50 ml total Glu 5% | 0.02 – 1 micrograms/kg/min | 0.018 ml/kg/hr = 0.1 micrograms/kg/min | | | | |
| Milrinone **PERIPHERAL** | 200 micrograms/ml | 10 mg in 50 ml total Glu 5% or NaCl 0.9% | 0.3 – 0.75 micrograms/kg/min | 0.09 ml/kg/hr = 0.3 micrograms/kg/min | | | | |
| Milrinone **CENTRAL** | 1 mg/ml | 50 mg in 50 ml Neat | 0.3 – 0.75 micrograms/kg/min | 0.018 ml/kg/hr = 0.3 micrograms/kg/min | | | | |
| Vasopressin (Argipressin) **CENTRAL Under 10kg** | 200 milliunits/ml | 10 units in 50 ml total Glu 5% or NaCl 0.9% | 0.15 – 2 milliunits/kg/min | 0.03 ml/kg/hr = 0.1 milliunits/kg/min | | | | |
| Vasopressin (Argipressin) **CENTRAL 10kg+** | 400 milliunits/ml | 20 units in 50 ml total Glu 5% or NaCl 0.9% | 0.15 – 2 milliunits/kg/min | 0.015 ml/kg/hr = 0.1 milliunits/kg/min | | | | |