

RIVERSIDE MULTISPECIALTY HOSPITAL

Department of Endocrinology

Follow-up Consultation Note

Patient Name:	John Doe	Medical Record #:	MRN-2026-001247
Age / Sex:	54 years / Male	Date of Visit:	2026-04-15
Consultant:	Dr. Sarah Smith, MD (Endocrinology)		

REASON FOR FOLLOW-UP

Three-month follow-up evaluation after initial diagnosis of Type 2 Diabetes Mellitus and initiation of pharmacologic therapy.

INTERVAL HISTORY

Mr. John Doe returns for his scheduled follow-up visit. Since the previous consultation, he reports significant improvement in his overall well-being. He states that excessive thirst and frequent urination have almost completely resolved. Fatigue has noticeably improved, and episodes of blurred vision are now infrequent.

The patient reports good adherence to Metformin therapy without significant adverse effects. He has attended dietary counselling sessions and has substantially reduced intake of refined carbohydrates and sugary beverages. He reports walking approximately 35–40 minutes five days per week. Home blood glucose records demonstrate improving fasting glucose levels averaging 110–130 mg/dL.

CURRENT MEDICATIONS

Medication	Dose	Status
Metformin	500 mg orally twice daily	Continue

PHYSICAL EXAMINATION & VITALS

Parameter	Current Visit	Previous Visit
Blood Pressure	132/84 mmHg	138/86 mmHg
Heart Rate	76 bpm	82 bpm
Respiratory Rate	16 / min	16 / min
Temperature	98.2 °F	98.4 °F
Weight	88 kg (-4 kg)	92 kg
Height	173 cm	173 cm
BMI	29.4 kg/m²	30.7 kg/m²

LABORATORY RESULTS

Test Description	Current Result	Previous Result
HbA1c	7.5%	8.6%
Fasting Plasma Glucose	124 mg/dL	186 mg/dL
Random Blood Glucose	168 mg/dL	241 mg/dL
Serum Creatinine	0.9 mg/dL	0.9 mg/dL
eGFR	>90 mL/min/1.73m²	>90 mL/min/1.73m²

ASSESSMENT

- **Type 2 Diabetes Mellitus:** Showing significant improvement following initiation of Metformin and lifestyle modification.
- **Weight Reduction:** Decreased by approximately 4 kg since previous visit; BMI improved from 30.7 to 29.4.
- **Improved Glycemic Control:** Demonstrated by a solid reduction in HbA1c from 8.6% to 7.5%.
- **Renal Function:** Remains within normal limits and stable.
- **Adherence:** Patient demonstrates excellent medication adherence and high lifestyle motivation.

TREATMENT PLAN

- **Pharmacotherapy:** Continue Metformin 500 mg orally twice daily.
- **Dietary:** Continue diabetic diet guidelines.
- **Physical Activity:** Maintain regular physical activity (minimum 150 minutes/week).
- **Monitoring:** Continue home blood glucose monitoring.
- **Goals:** Reinforce ongoing weight-loss goals.

- **Follow-up Labs:** Repeat HbA1c in 3 months.

FOLLOW-UP

Patient to return in approximately 3 months for reassessment. If glycemic improvement continues, current therapy will be maintained.

Electronically Signed By:

Dr. Sarah Smith, MD

Consultant Endocrinologist

Riverside Multispecialty Hospital