

Medical necessity/Clinical guidelines 1. 95249 Short Term Interstitial CGM Use of an U.S. FDA approved short term interstitial CGM, minimum of 72 hours may be considered medically necessary for type I or type II diabetics when ANY ONE of the following criteria are met: • History of poorly controlled diabetes (i.e., unexplained hypoglycemic episodes, hypoglycemic unawareness, suspected postprandial hyperglycemia, and recurrent diabetic ketoacidosis); or • Pregnant female that requires insulin therapy; or • Pregnant female who develops gestational diabetes; or • An individual who requires determination of basal insulin level measurements prior to insulin pump initiation; or • An individual who has documentation of ALL of the following: o Inability to recognize or communicate their symptoms of hypoglycemia; and o Pediatric individuals (age 2 (two) to 18 years of age) must be managed by endocrinologist; and o The parent or caregiver must have the ability to understand the technology and be willing to use the monitor, read and interpret the glucose data; and o Received diabetes self-management education and instruction from a health care professional with expertise in the management of diabetes; and o Is on an intensive insulin regimen, requiring three (3) or more insulin administrations per day, or utilizes an insulin pump; and o Has ONE or MORE of the following while on an intensive insulin regimen: ▪ Unexplained large fluctuations in daily glucose values before meals; or ▪ Unexplained frequent hypoglycemic attacks (e.g., greater than one (1) per week or three (3) per month); or ▪ Episodes of ketoacidosis or hospitalizations for significantly elevated glucose levels. For short-term diagnostic use, no more than four (4) CGM periods may be considered medically necessary within a 12-month period. Short-term interstitial CGM not meeting the criteria as indicated in this policy is considered experimental/investigational and therefore non-covered because the safety and/or effectiveness of this service cannot be established by the available published peer-reviewed literature. Procedure Codes 95249 95250 95251 Long Term Interstitial CGM Systems Use of an U.S. FDA approved long term interstitial CGM system as an adjunct and non adjunct (i.e., greater than 72 hours) may be considered medically necessary for type I or type II diabetics when ANY ONE of the following criteria are met: • Pregnant female with poorly controlled diabetes (i.e., unexplained hypoglycemic episodes, hypoglycemic unawareness, suspected postprandial hyperglycemia, and recurrent diabetic ketoacidosis); or • Recurrent, unexplained, severe (generally blood glucose levels less than 50 mg/dL) hypoglycemia that puts the individual or others at risk; or • History of poorly controlled diabetes (i.e., unexplained hypoglycemic episodes, hypoglycemic unawareness, suspected postprandial hyperglycemia, and recurrent diabetic ketoacidosis); and • Documentation of ALL of the following: o The CGM is prescribed by a professional provider; and o Is on an intensive insulin regimen, requiring three (3) or more insulin administrations per day, or utilizes an insulin pump; and o The individual (parent or caregiver if individual is a child) has been instructed by a health care professional in the management of diabetes including care of the CGM, alarm use and problem solving and use of real time CGM application in diabetic care; and o Pediatric individuals (age 2 (two) -12 years old: The CGM is recommended and managed by a pediatric endocrinologist: ▪ Note: CGM may be approved for those under two (2) years of age with exceptional circumstances requiring individual consideration; and o The individual (parent or caregiver if the individual is a child) must have the ability to understand the technology and be willing to use the monitor (i.e., hear alarms, read and interpret glucose data and can take action based on the data interpretation; and o The responsible individual demonstrates mastery of the fundamentals of diabetes self-management, which includes BOTH of the following: ▪ Routine, regular testing of blood glucose levels; and ▪ Maintaining accurate records of blood glucose testing. Long-term interstitial CGM system not meeting the criteria as indicated in this policy is considered experimental/investigational and therefore non-covered because the safety and/or effectiveness of this service cannot be established by the available published peer-reviewed literature. Receiver/Reader kits are limited to one (1) per Benefit Period. Sensor kits are limited to one (1) unit every thirty (30) days, unless otherwise required due to manufacturer packaging restrictions.

Transmitter kits are limited to (1) unit every ninety (90) days unless otherwise required due to manufacturer packaging restrictions. Procedure Codes 95249 A4238 E2103 95250 A4239 G0564 95251 A4271 G0565 0446T A9276 0447T A9277 0448T A9278 A4226 E2102