

Name: Prathapani Satwika

Reg: 20BCD7160

Title of the experiment:-

Determination of energy band gap of a semiconductor by four probe method.

Objective:-

To determine the band gap of a semiconductor by measuring the resistivity as a function of temperature using four probe method.

Equipment list:-

- 1) Spring loaded four probes
- 2) Germanium (semiconductor) crystal
- 3) oven (upto 150°C)
- 4) Thermometer
- 5) Constant current source
- 6) Voltmeter


Formula:-

$$\sigma(T) = \sigma_0 \exp\left(-\frac{E_g}{2kT}\right)$$

$$\rho(T) = \rho_0 \exp\left(+\frac{E_g}{2kT}\right)$$

σ_0 - Residual conductivity

ρ_0 - Residual resistivity



E_g - Band gap of the intrinsic semiconductor

k - Boltzman constant

T - absolute temperature

$$\rho = CF \left(\frac{V}{I} \right) \Rightarrow \rho = 0.213 \times \left(\frac{V}{I} \right)$$

I - current passing through the semiconductor

V - measured voltage

Band gap Energy:-
 $E_g = 2km$

$m \rightarrow$ slope of the graph $\log_e \rho \text{ vs } 1/T$

Report:- $k \rightarrow$ Boltzman constant $= 8.617 \times 10^{-5} \text{ eV/K}$

S.No	Temperature (K)	voltage (V)	Resistivity (ρ)	$1/T$ (K^{-1})	$\log_e \rho$
1	300	0.303	18.65 Ωm	3.33×10^{-3}	2.91
2	313	0.323	19.65 Ωm	3.19×10^{-3}	2.97
3	323	0.331	20.14 Ωm	3.09×10^{-3}	3.00
4	333	0.332	20.20 Ωm	3.00×10^{-3}	3.00
5	343	0.315	19.17 Ωm	2.91×10^{-3}	2.95
6	353	0.300	18.25 Ωm	2.83×10^{-3}	2.90
7	363	0.223	13.57 Ωm	2.75×10^{-3}	2.60

calculations:-

$$C = 0.213 \text{ cm}$$

$$I = 3.5 \times 10^{-3}$$

obs1:-

$$P = 0.213 \times \frac{V}{I} = 0.213 \times \frac{0.303}{3.5 \times 10^{-3}} \\ = 18.439 \text{ } \mu\text{m}$$

obs2:

$$P = 0.213 \times \frac{V}{I} = \frac{0.213 \times 0.323}{3.5 \times 10^{-3}} = 19.65 \text{ } \mu\text{m}$$

obs3:

$$P = 0.213 \times \frac{V}{I} = \frac{0.213 \times 0.331}{3.5 \times 10^{-3}} = 20.14 \text{ } \mu\text{m}$$

obs4:

$$P = 0.213 \times \frac{V}{I} = \frac{0.213 \times 0.332}{3.5 \times 10^{-3}} = 20.20 \text{ } \mu\text{m}$$

obs5:

$$P = 0.213 \times \frac{V}{I} = \frac{0.213 \times 0.315}{3.5 \times 10^{-3}} = 19.17 \text{ } \mu\text{m}$$

obs6:

$$P = 0.213 \times \frac{V}{I} = \frac{0.213 \times 0.300}{3.5 \times 10^{-3}} = 18.25 \text{ } \mu\text{m}$$

obs7:

$$P = 0.213 \times \frac{V}{I} = \frac{0.213 \times 0.223}{3.5 \times 10^{-3}} = 13.57 \text{ } \mu\text{m}$$

From graph,

$$\rho(T) = \rho_0 \exp\left(\frac{E_g}{2kT}\right)$$

$$\log \rho(T) = \log \rho_0 + \frac{E_g}{2kT}$$

$$M = \frac{E_g}{2k} \quad [k \text{ is Boltzmann's constant}]$$

$$k = 8.617 \times 10^{-5} \text{ eV/K}$$

$$E_g = 2k \times M$$

$$E_g = 2 \times 8.617 \times 10^{-5} \times 3.75 \times 10^3$$

$$\boxed{E_g = 0.64 \text{ eV}}$$

Result:-

The band gap energy (E_g) of the germanium crystal is 0.64 eV

← (log P) →

scale.
 $x = 0.25 \text{ unit} = 0.0 \times 10^{-3} \text{ K}^{-1}$
 $y = 0.015 \text{ unit} = 0.01$

