

Name: - L Prathyusha

In [5]:

```
import numpy as np
a = np.array([0, 1, 2, 3])
a
```

Out[5]:

```
array([0, 1, 2, 3])
```

In [6]:

```
import numpy as np
from scipy.spatial.distance import pdist, squareform

# Create the following array where each row is a point in 2D space:
# [[0 1]
#  [1 0]
#  [2 0]]
x = np.array([[0, 1], [1, 0], [2, 0]])
print(x)

# Compute the Euclidean distance between all rows of x.
# d[i, j] is the Euclidean distance between x[i, :] and x[j, :],
# and d is the following array:
# [[ 0.          1.41421356  2.23606798]
#  [ 1.41421356  0.          1.          ]
#  [ 2.23606798  1.          0.          ]]
d = squareform(pdist(x, 'euclidean'))
print(d)
```


```
[[0 1]
 [1 0]
 [2 0]]
[[0.          1.41421356  2.23606798]
 [1.41421356  0.          1.          ]
 [2.23606798  1.          0.          ]]
```

In [7]:

```
import numpy as np
import matplotlib.pyplot as plt

# Compute the x and y coordinates for points on sine and cosine curves
x = np.arange(0, 3 * np.pi, 0.1)
y_sin = np.sin(x)
y_cos = np.cos(x)

# Plot the points using matplotlib
plt.plot(x, y_sin)
plt.plot(x, y_cos)
plt.xlabel('x axis label')
plt.ylabel('y axis label')
plt.title('Sine and Cosine')
plt.legend(['Sine', 'Cosine'])
plt.show()
```



In []: