13 group (Supplementary Fig. 2). The mITT and safety populations included 1167 and 1169 patients, respectively. Demographics and baseline characteristics were similar across the treatment groups (Table 1). Patients had an average age of 58 years, were predominantly Caucasian (~90%), were overweight or obese (BMI ~32 kg/m2), and had a mean duration of diabetes of approximately 9 years. Primary Efficacy Endpoint Baseline HbA1c was 8.1% (65 mmol/mol) in all three groups. Mean HbA1c levels achieved at Week 30 were 6.5% (48 mmol/mol) for iGlarLixi, 6.8% (51 mmol/mol) for iGlar, and 7.3% (56 mmol/mol) for Lixi (Fig. 1A). The least squares (LS) mean changes from baseline to Week 30 in HbA1c were -2.70% for iGlarLixi, -1.34% for iGlar, and -0.85% for Lixi (Table 2; Fig. 1B). Statistical superiority of iGlarLixi over Lixi was demonstrated for the change in HbA1c from baseline to Week 30 (LS mean difference versus Lixi -0.8% [-8.5 mmol/mol] [95% CI: -0.9, -0.7 (-9.8 mmol/mol, -7.3 mmol/mol); p<0.0001]). The LS mean HbA1c difference at Week 30 between iGlarLixi and iGlar (-0.3% [-3.2 mmol/mol] [95% CI: -0.4, -0.2 (-4.2 mmol/mol, -2.1 mmol/mol); p<0.0001]) met non-inferiority of iGlarLixi compared with iGlar and also demonstrated superiority for this primary efficacy endpoint (p<0.0001) based on the step-down testing procedure. Secondary Efficacy Endpoints A significantly higher proportion of patients in the iGlarLixi group (74%) reached the HbA1c target of <7% (53 mmol/mol) compared with patients receiving iGlar (59%) or Lixi (33%), or the HbA1c target of ?6.5% (48 mmol/mol) (p<0.0001 for all comparisons) (Table 2). Body weight increased in the iGlar group (+1.1 kg) and