

## T-Test

- ① A T-test is a statistical test used in hypothesis testing to compare the mean of two samples or to compare a sample mean to a known population mean.
- ② It is used when the population standard deviation is unknown and the sample size is small.
- ③ There are three main type of t-test —

### ① one sample t-test

The one sample t-test is used to compare the mean of a single sample to a known population mean. Here our Null hypothesis states that there is no significant ~~different~~ difference between the sample mean and the population mean.

### ② Assumption :

- ① Normality
- ② Independence (the observation of sample must be independent)
- ③ Random sampling
- ④ Unknown population std.

(Q) Suppose a manufacturer claims that the average weight of their new chocolate bars is 50 grams, we highly doubt that and want to check this so we drew out a sample of 25 chocolate bars and measured their weight, the sample mean came out to be 49.7 gm. and the sample SD was 1.2 gm. Consider the significance level to be 0.05.

Null Hypothesis ( $H_0$ ): ~~there is no significant difference between~~  
Average weight of chocolate bar is 50 gm.  
 $\mu = 50$

Alternative Hypothesis ( $H_a$ ):  $\mu \neq 50$

We have population mean ( $\mu$ ) = 50, sample mean ( $\bar{x}$ ) = 49.7  
sample s.d. ( $s$ ) = 1.2,  $N = 25$

Now, we need to find the T-statistic.

Now,

$$T\text{-statistic} = \frac{\text{Sample Mean} - \text{population mean}}{\frac{\text{sample SD}}{\sqrt{\text{sample size}}}}$$

Here

$$T\text{-statistic} = \frac{49.7 - 50}{1.2/\sqrt{25}} = -1.25$$

(Here degree of freedom (df) =  $(N-1) = (25-1) = 24$ )

Now, we are going to find out the p-value.



So, our p-value =  $(0.11167 \times 2) = 0.22334$ .

As our p-value > significance level (0.05)

Then we can't reject our Null hypothesis.

so, the average weight of chocolate bar is 50 gm. (proved)

## Independent two-sample t-test

The independent two sample t-test is used to compare the means of two independent samples. The null hypothesis states that there is no significance difference between the means of two samples.

It is also known as unpaired t-test.

### Assumption

① Independence of observations (Two sample must be independent)      ② Normality      ③ Equal Variance (Homoscedasticity)

④ Random sampling

Suppose a Website owner claims that there is no difference in the average time spend on their website between desktop and mobile users. To test this claim, we collect data for 30 desktop users and 30 mobile users regarding the time spend on the website in minutes. The sample statistics are as following —

desktop users = [12, 15, 18, 16, 20, 17, 14, 22, 19, 21, 23, 18, 25, 17, 16, 24, 20, 19, 22, 18, 15, 14, 23, 16, 12, 21, 19, 17, 20, 14]

Mobile users = [10, 12, 14, 13, 16, 15, 11, 17, 14, 16, 18, 14, 20, 15, 14, 19, 16, 15, 17, 14, 12, 11, 18, 15, 10, 16, 13, 16, 11]

and desktop SD = 9.5 and Mobile SD = 2.7.

First let construct our null hypothesis as well as alternative hypothesis —

Null hypothesis ( $H_0$ ): ~~there is~~ Desktop User Mean = Mobile user Mean

Alternative hypothesis ( $H_a$ ): Desktop user mean  $\neq$  Mobile user Mean

Now, we need to find out the t-statistic.

The formula of T-statistic be —

$$t = \frac{\text{First sample mean} - \text{Second sample mean}}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

where  $s_1$  = First sample SD  
 $s_2$  = Second sample SD


$n_1$  = First sample size

$n_2$  = Second sample size

$$\text{Here } t\text{-statistic} = \frac{18.5 - 11.3}{\sqrt{\frac{3.5^2}{30} + \frac{2.7^2}{30}}} = \frac{7.2}{0.81} = 8.89$$

$$\begin{aligned}\text{Now, degree of freedom} &= (n_1 - 1) + (n_2 - 1) \\ &= 29 + 29 \\ &= 58\end{aligned}$$

Now, corresponding this t-statistic and df (58) we have p-value = ~~2.7146~~  
 $= 2.7 \times 10^{-6}$



so, clearly p-value < level of significance (0.05).

so, Null hypothesis can be rejected.

so, the average time spend on desktop and average time spend on mobile are different. (Proved)

## ② Pair-2 sample t-test

A paired two sample t-test, also known as a dependent or paired samples t-test, is a statistical test used to compare the means of two related or dependent groups.

### Assumptions

① paired observations ② Normality ③ Independence of pairs (Each pair of observations should be independent).

④ Let's assume that a fitness center is evaluating the effectiveness of a new 8-week weight loss program. They enroll 15 participants in the program and measure their weights before and after the program.

The goal is to test whether the new weight loss program leads to a significant reduction in the participant weights.

Before the program —

[80, 92, 75, 68, 85, 78, 73, 90, 70, 88, 76, 84, 82, 77, 91]

After the program —

[78, 93, 81, 67, 88, 76, 74, 91, 69, 88, 77, 81, 80, 79, 88]

① Null hypothesis :  $(H_0)$  :  $\text{Mean}_{\text{before}} = \text{Mean}_{\text{after}}$

Alternative hypothesis  $(H_a)$  :  $\text{Mean}_{\text{before}} > \text{Mean}_{\text{after}}$

basically it is —


$$t\text{-statistic} = \frac{\bar{X}_{diff}}{S.D_{diff} / \sqrt{n}}$$

so, here t-statistic should be —

$$= \frac{-0.0667}{2.379 / \sqrt{15}}$$

$$= -0.108$$

$$\approx -0.11$$




( $\bar{X}_{diff}$  = Mean of the difference columns of before and after)

Means first we calculate the difference of before column and after column then we calculate the mean of that ~~no~~ new column.

~~S.D.~~ and  $S.D_{diff}$  = S.D of that new column

Here —

Before	After	Difference column
80	78	2
92	93	-1
75	81	-6
68	67	1
85	88	-3
78	76	2
73	74	-1
90	91	-1
70	91	-1
88	69	1
76	88	0
84	77	-1
82	81	3
77	80	2
91	79	-2
	88	3



so, with respect to the  $t$ -statistic  $(-0.11)$  and degree of freedom  $(14)$   
We got  $p\text{-value} = 0.5424$

so,  $p\text{-value} > \text{level of significance } (0.05)$

so, we can't reject our null hypothesis. so that mean we haven't  
enough evidence to prove that the mean average of weight  
before program is greater than the mean average of weight after  
program. (Proved)

---

⑦ Common scenario where paired t-test is used

- ① Match or correlated group    ② Before and After studies problems
-