

Problem Submissions Leaderboard

RATE THIS CHALLENGE

Java Loops II *

We use the integers a, b, and n to create the following series:

$$(a+2^0\cdot b), (a+2^0\cdot b+2^1\cdot b), \ldots, (a+2^0\cdot b+2^1\cdot b+\ldots +2^{n-1}\cdot b)$$

Editorial

You are given q queries in the form of a,b, and n. For each query, print the series corresponding to the given a,b, and n values as a single line of n space-separated integers.

Input Format

The first line contains an integer, q, denoting the number of queries.

Each line i of the q subsequent lines contains three space-separated integers describing the respective a_i , b_i , and n_i values for that query.

Constraints

- $0 \le q \le 500$
- $0 \le a, b \le 50$
- $1 \le n \le 15$

Output Format

For each query, print the corresponding series on a new line. Each series must be printed in order as a single line of n space-separated integers.

Sample Input

2 0 2 10

5 3 5

Sample Output

2 6 14 30 62 126 254 510 1022 2046 8 14 26 50 98

Explanation

We have two queries:

1. We use a=0, b=2, and n=10 to produce some series $s_0, s_1, \ldots, s_{n-1}$:

- $\circ \ s_0=0+1\cdot 2=2$
- $s_1 = 0 + 1 \cdot 2 + 2 \cdot 2 = 6$
- $\circ \ \ s_2 = 0 + 1 \cdot 2 + 2 \cdot 2 + 4 \cdot 2 = 14$
- ... and so on.

Once we hit n = 10, we print the first ten terms as a single line of space-separated integers.

2. We use a=5, b=3, and n=5 to produce some series $s_0, s_1, \ldots, s_{n-1}$:

- $s_0 = 5 + 1 \cdot 3 = 8$
- $s_1 = 5 + 1 \cdot 3 + 2 \cdot 3 = 14$
- \circ $s_2 = 5 + 1 \cdot 3 + 2 \cdot 3 + 4 \cdot 3 = 26$
- $s_3 = 5 + 1 \cdot 3 + 2 \cdot 3 + 4 \cdot 3 + 8 \cdot 3 = 50$
- $s_4 = 5 + 1 \cdot 3 + 2 \cdot 3 + 4 \cdot 3 + 8 \cdot 3 + 16 \cdot 3 = 98$



We then print each element of our series as a single line of space-separated values.

