

Measles Exposure - Female (Adult)

Algorithm 1474 | Adult (18 years and older) | Gender at birth: Female

Definition

- Measles contact/exposure assessment decomposed from NHS.UK's published measles guidance's contact-tracing recommendations.
- Adult patient.
- Female.

Initial assessment

1. [OPEN_TEXT] What is the exposed person's exact age and sex, are they pregnant or immunocompromised, when/where/how close was contact, who confirmed measles, and what vaccination or prior-immunity evidence exists? Include infant and safeguarding concerns.
2. [YES_NO] Has the person had 2 doses of MMR vaccine, or had measles before?
3. [YES_NO] Is the person pregnant, under 1 year old, or immunocompromised?

Triage and redirect questions

1. Redirect → Rash or Redness - Widespread

Is a widespread rash the main concern rather than contact with a person who may have measles?

A widespread rash belongs in the Rash or Redness - Widespread pathway.

1. Disposition level 70

Has the person had close contact with someone with measles and is unvaccinated or only partially vaccinated, pregnant, under 1 year old, or has a weakened immune system?

NHS.UK measles guidance recommends urgent clinical contact for these groups after measles exposure - measles in pregnancy can cause miscarriage, stillbirth, or premature birth, and infants/immunocompromised people are at higher risk of severe disease.

1. Disposition level 15

Has the person had 2 doses of MMR vaccine or measles before, with none of the higher-risk factors above?

Fully vaccinated or previously infected individuals are generally protected and can monitor at home for symptoms.

Advice

Urgent measles exposure follow-up

- Contact the Qatar urgent clinical review pathway promptly - do not simply walk into a clinic, as measles is highly contagious and the clinic may need to prepare to see you safely.
- Call Back If: fever or rash develops.

Governed Qatar measles-exposure monitoring

- Do not attend a clinic unannounced. Avoid contact with infants, pregnant people, and immunocompromised people, and contact the Qatar public-health/clinical route approved by governance to verify exposure, immunity, vaccination records, and whether time-sensitive post-exposure action applies. Monitor for fever, cough, runny nose, red eyes, and spreading rash; call ahead before any in-person care. Vaccine or immunoglobulin eligibility/timing must not be invented by UAT content.
- Call Back If: fever, cough, red eyes, or rash develops; breathing difficulty, dehydration, confusion, seizure, or severe illness.

First aid

References

- NHS.UK, "Measles", <https://www.nhs.uk/conditions/measles/> (page last reviewed 31 July 2025) - contact-tracing/exposure guidance section

Provenance

Shaped to mirror the licensed After-Hours Telehealth Triage Guidelines database structure. This is IST open-source guideline content derived from cited public sources; it is not licensed STCC content. Question.DispositionLevel=null marks a See More Appropriate Guideline redirect. Demographic variants contain only source-supported differences; identical thresholds are documented rather than fabricated. Local destinations and TelemedicineEligible values require IST clinical governance approval. UAT DATA ONLY - NOT FOR REAL-PATIENT CARE OR PRODUCTION USE.

Manifest hash: 3b5bb4cc015ba5558566fd4d2c511e2bb134d4e5c1c3c1aba7cb79e09c184470 | Status: BLOCKED_BY_RESEARCH_GAP