

Motion Sickness - Female (Child)

Algorithm 1212 | Child (0-17 years) | Gender at birth: Female

Definition

- Motion sickness assessment decomposed from NHS.UK's published guidance.
- Child patient.
- Female.

Initial assessment

1. [OPEN_TEXT] What is the person's exact age and sex, are they pregnant, what travel triggered symptoms, and are symptoms typical and limited to travel? Ask about medicines, neurological/ear symptoms, dehydration, poisoning, injury, and driving/aviation duties.
2. [DURATION] How long has this lasted?
3. [YES_NO] Is the person able to keep fluids down?
4. [CAREGIVER_OBSERVATION] What has a safe parent or caregiver observed about alertness, breathing, pain, swelling, fever, eating or drinking, sleep, play, and normal activity?

Triage and redirect questions

1. Redirect → Nausea

Is nausea occurring without travel, movement, or motion exposure?

Nausea not triggered by motion belongs in the general Nausea pathway.

1. Disposition level 100

Does this sound like a life-threatening emergency to the triager (e.g. severe dehydration or symptoms unrelated to travel)?

Universal emergency rule-out added ahead of the motion sickness guidance itself - motion sickness is not itself an emergency, but persistent vomiting could cause dehydration or mask another condition.

1. Disposition level 15

Is this typical motion sickness with none of the features above?

NHS.UK guidance provides prevention and self-care measures for typical motion sickness.

Advice

Qatar emergency vomiting/dizziness response

- Stop travel safely and call Qatar emergency services on 999; do not drive when symptoms include reduced consciousness, seizure, stroke signs, severe headache, chest pain, abnormal breathing, severe dehydration, blood/green vomit, major injury, or suspected poisoning. Do not give oral medicine or fluids if drowsy or unsafe to swallow. Call Qatar 999 now, follow the call-taker's ambulance and emergency-transport instructions, and do not self-drive. Do not delay emergency care.
- Call Back If: confusion, seizure, neurological change, or collapse; severe dehydration or inability to keep fluids down; blood/green vomit or severe pain.

Governed motion-sickness prevention

- If safe, look toward the horizon, get fresh air, avoid reading/screens, and take breaks. Medicine, patches, and complementary products require age/weight, pregnancy/breastfeeding, glaucoma/urinary/cardiac conditions, sedation, driving/aviation duties, and current-medicine checks under the approved Qatar pathway; no age cutoff or product is authorized by this UAT content.
- Call Back If: unable to keep fluids down; symptoms persist after travel or are unlike prior motion sickness.

First aid

- Stop travel safely and call Qatar emergency services on 999; do not drive when symptoms include reduced consciousness, seizure, stroke signs, severe headache, chest pain, abnormal breathing, severe dehydration, blood/green vomit, major injury, or suspected poisoning. Do not give oral medicine or fluids if drowsy or unsafe to swallow. Call Qatar 999 now, follow the call-taker's ambulance and emergency-transport instructions, and do not self-drive. Do not delay emergency care.

References

- NHS.UK, "Motion sickness", <https://www.nhs.uk/conditions/motion-sickness/> (page last reviewed 19 June 2023)

Provenance

Shaped to mirror the licensed After-Hours Telehealth Triage Guidelines database structure. This is IST open-source guideline content derived from cited public sources; it is not licensed STCC content. Question.DispositionLevel=null marks a See More Appropriate Guideline redirect. Demographic variants contain only source-supported differences; identical thresholds are documented rather than fabricated. Local destinations and TelemedicineEligible values require IST clinical governance approval. UAT DATA ONLY - NOT FOR REAL-PATIENT CARE OR PRODUCTION USE.

Manifest hash: 67a6bfb5c5d7de82b65bbd9c20416fcd82a566963f5c102e1a4f82e18e13723ce | Status: BLOCKED_BY_RESEARCH_GAP