

Ringworm - Male (Adult)

Algorithm 1337 | Adult (18 years and older) | Gender at birth: Male

Definition

- Ringworm assessment decomposed from NHS.UK's published ringworm guidance.
- Adult patient.
- Male.

Initial assessment

1. [OPEN_TEXT] What is the person's exact age and sex, are they pregnant or immunocompromised, where is the rash, and does it involve scalp, beard, nails, face, genitals, widespread skin, infection, household contacts, animals, or safeguarding concerns?
2. [DURATION] How long has it been present?
3. [YES_NO] Has it been treated with an antifungal already?

Triage and redirect questions

1. Redirect → Impetigo (Infected Sore)

Is this mainly a weeping, blistered, or honey-crusted sore rather than a ring-shaped scaly rash?

A weeping or honey-crusted infected sore belongs in the Impetigo pathway.

1. Disposition level 70

Has the ringworm not improved after antifungal treatment recommended by a pharmacist, does it affect the scalp, or does the person have a weakened immune system (from chemotherapy, steroids, or diabetes)?

NHS.UK ringworm guidance recommends clinical review for these situations - scalp ringworm typically needs prescription treatment.

1. Disposition level 15

Is this a typical ring-shaped rash with none of the features above?

NHS.UK guidance recommends pharmacist-guided antifungal treatment as the first step for typical ringworm.

Advice

Ringworm needing clinical review

- Arrange a primary-care appointment for prescription-strength antifungal treatment.
- Call Back If: spreads rapidly; signs of secondary bacterial infection develop.

Governed ringworm care

- Keep the area clean/dry, avoid scratching and sharing towels, clothing, combs, or bedding, wash fabrics regularly, and consider affected household members/pets. Antifungal product, duration, scalp/nail treatment, pregnancy/breastfeeding, pediatric age, immunocompromise, and medicine interactions require approved Qatar pharmacist/clinician guidance. Do not use steroid cream alone on an undiagnosed fungal rash.
- Call Back If: scalp, beard, nail, widespread, painful, infected, or rapidly spreading involvement; no improvement or immunocompromised patient.

First aid

References

- NHS.UK, "Ringworm", <https://www.nhs.uk/conditions/ringworm/> (page last reviewed 03 August 2023)

Provenance

Shaped to mirror the licensed After-Hours Telehealth Triage Guidelines database structure. This is IST open-source guideline content derived from cited public sources; it is not licensed STCC content. Question.DispositionLevel=null marks a See More Appropriate Guideline redirect. Demographic variants contain only source-supported differences; identical thresholds are documented rather than fabricated. Local destinations and TelemedicineEligible values require IST clinical governance approval. UAT DATA ONLY - NOT FOR REAL-PATIENT CARE OR PRODUCTION USE.

Manifest hash: e7d10461ab8c6909864a2bd4ed385c067166faaa33f50ce35037da746abe4284 | Status: BLOCKED_BY_RESEARCH_GAP