

# Postpartum - Urination Pain - Female (Child)

Algorithm 1438 | Child (12-17 years) | Gender at birth: Female

## Definition

- Postpartum painful urination assessment, combining NHS.UK's C-section recovery urinary-symptom warning with the cystitis guidance already used for Urination Pain - Female (batch14).
- Child patient.
- Female.

## Initial assessment

1. [DURATION] How many days/weeks since delivery?
2. [TEMPERATURE] What is the temperature, if measured?
3. [YES\_NO] Any back/flank pain or blood in the urine?
4. [YES\_NO] If the patient is under 18, can they speak privately now, and do they say they feel safe with the accompanying adult?
5. [CAREGIVER\_OBSERVATION] What has a safe parent or caregiver observed about alertness, breathing, pain, swelling, fever, eating or drinking, sleep, play, and normal activity?

## Triage and redirect questions

### **1. Redirect → Flank Pain**

Is pain in the side or back below the ribs the primary concern rather than pain during urination?

Primary side or back pain below the ribs belongs in the Flank Pain pathway.

### **1. Disposition level 100**

Along with urinary symptoms, is there confusion, trouble breathing or very fast breathing, fainting, cold or clammy skin, or rapidly worsening severe illness?

These are sepsis red flags, and postpartum infection can progress quickly - always treated as an emergency.

### **1. Disposition level 70**

Is there pain when urinating, leaking urine, back or flank pain, fever, or blood in the urine?

NHS.UK guidance lists painful or leaking urination after a C-section as needing prompt review, and the same applies to any postpartum delivery given the shared infection risk.

## Advice

### **Emergency postpartum infection precautions**

- Call Qatar emergency services on 999 now. Do not let the patient drive. Call Qatar 999 now, follow the call-taker's ambulance and emergency-transport instructions, and do not self-drive. Do not delay emergency care.
- Call Back If: worsening confusion; worsening breathing.

### **Urgent postpartum urinary symptom review**

- Arrange prompt clinical assessment; the exact non-emergency Qatar destination requires local approval. Postpartum urinary symptoms may need examination and testing.
- Call Back If: fever develops; back pain worsens.

## First aid

- Call Qatar emergency services on 999 now. Do not let the patient drive. Call Qatar 999 now, follow the call-taker's ambulance and emergency-transport instructions, and do not self-drive. Do not delay emergency care.

## References

- NHS.UK, "Caesarean section - Recovery" (urinary symptom warning), <https://www.nhs.uk/conditions/caesarean-section/recovery/> (page last reviewed 04 January 2023)
- NHS.UK, "Cystitis" (already cited for Urination Pain - Female, batch14), <https://www.nhs.uk/conditions/cystitis/>

## Provenance

Shaped to mirror the licensed After-Hours Telehealth Triage Guidelines database structure. This is IST open-source guideline content derived from cited public sources; it is not licensed STCC content. Question.DispositionLevel=null marks a See More Appropriate Guideline redirect. Demographic variants contain only source-supported differences; identical thresholds are documented rather than fabricated. Local destinations and TelemedicineEligible values require IST clinical governance approval. UAT DATA ONLY - NOT FOR REAL-PATIENT CARE OR PRODUCTION USE.

Manifest hash: b867a025eb432951ccdbbcf4a486d4348e4d6e416ef55a6beeb419c54252fe5e | Status: BLOCKED\_BY\_RESEARCH\_GAP