

Presentation of STCC content within a nurse triage application

1. In this software mock-up, calls are answered by a non-clinical person and populated into the Pathways Call Center Software for the nurse to pick their next call from the Call Queue.

Pathways Call Center Software

Call Queue

New Encounter

Call Queue

Nurse Triage Calls

Kathy Johnson RN

Pascal Health System

Acuity	Reason for Call	Age	Gender	Last Name	Minutes in Queue	Status	Action
2	Hit my head and now am vomiting. Denies LOC.	19	M	Smith	0	In Call	None
3	Diarrhea and stomach pains.	49	M	Prunitico	0	In Call	None
3	Caller playing basketball, jumped and fell on ankle.	24	M	Anderson	16	Waiting	Call Back
3	Mother calling about 5 yo who is complaining his ears hurt.	5	M	Ivanhoe	10	Waiting	Call Back
3	Patient was punched in the face by someone in a bar 2 hours ago. No loss of consciousness.	18	F	Munoz	7	Waiting	Call Back
4	Runny nose and cough.	10	M	Rogers	22	Waiting	Call Back
4	Earache for two days	85	M	Reed	20	Waiting	Call Back
4	What is the right dose of Advil	1	F	Rogers	17	Waiting	Call Back
5	My knee hurts	63	M	Rogers	3	Waiting	Call Back

2. The nurse identifies the Reason for Call and searches the database using the search word of Ankle. Corresponding topics with a search word of ankle are presented to the nurse to choose from. Based on dialog with the caller, the nurse selects “Ankle and Foot Injury” and continues with triage.

Pathways Call Center Software

Patient

Profile

Encounter History

Triage Process

Reason for Call

Triage Questions

Care Advice

Encounter Summary

Resources

Background

First Aid

Powered by:
Schmitt-Thompson
Clinical Content

Reason for Call

Kathy Johnson RN

Patient Name: John Anderson

Caller Name: Self

Encounter Date: 6/1/2021

Age: 24

Relationship: Self

Gender at Birth: M

PCP: Marcus Welby MD

Reason for Call

Caller playing basketball, jumped and fell on ankle.

Guideline Search

ankle

Enter a search word

Search

Guidelines Found

Ankle and Foot Injury

Injuries to a bone, muscle, joint, or ligament of the ankle and foot. Associated skin and soft tissue injuries are also included

Ankle Pain

Pain in the ankle. Not due to a traumatic injury

Ankle Swelling

Swelling in or around an ankle joint. Looks different than the other ankle. Not due to a traumatic injury.

Leg Injury

Injuries to a bone, muscle, joint or ligament of the leg. Associated skin and soft tissue injuries are also included.

Leg Swelling and Edema

Swelling of the leg(s) or generalized edema of body. Pedal edema (bilateral swelling of feet and ankles). Not due to a traumatic injury.

BACK

- The nurse begins asking triage questions with the most urgent/acute questions first. Once 911 is ruled out, there's a section called See More Appropriate Guideline for the nurse to review to make sure he/she is in the most appropriate guideline. Also note the Resource section for Background and First Aid. This is all content in the database that should be part of the software user interface and presented to the nurse to review.

Pathways Call Center Software

Patient

Profile

Encounter History

Triage Process

Reason for Call

Triage Questions

Care Advice

Encounter Summary

Resources

Background

First Aid

Powered by:

Schmitt-Thompson Clinical Content

Triage Assessment Questions

Kathy Johnson RN

Patient Name: John Anderson

Caller Name: Self

Encounter Date: 6/1/2021

Age: 24

Relationship: Self

Gender at Birth: M

PCP: Marcus Welby MD

Protocol: Ankle and Foot Injury (Adult)

Call EMS 911 Now

No To All

Yes No

○

○

Serious injury with multiple fractures (broken bones)

○

○

[1] Major bleeding (e.g., actively dripping or spurting) AND [2] can't be stopped

FIRST AID: Apply direct pressure to the entire wound with a clean cloth.

○

○

Amputation

FIRST AID: Apply direct pressure to the entire wound with a clean cloth.

○

○

Looks like a dislocated joint (very crooked or deformed)

Reason: If dislocated, adult will be unable to walk at all. Possible vascular compromise. Needs reduction.

○

○

Sounds like a life-threatening emergency to the triager

See More Appropriate Guideline

No To All

Yes No

○

○

Wound looks infected

Go to Guideline: Wound Infection (Adult)

○

○

Caused by an animal bite

Go to Guideline: Animal Bite (Adult)

BACK

NEXT

Schmitt-Thompson
Clinical Content

Proprietary – Do not distribute

- Pathways Call Center Software

Patient

Profile

Encounter History

Triage Process

Reason for Call

Triage Questions

Care Advice

Encounter Summary

Resources

Background

First Aid

Triage Assessment Questions

Kathy Johnson RN

Patient Name: John Anderson

Caller Name: Self

Encounter Date: 6/1/2021

Age: 24

Relationship: Self

Gender at Birth: M

PCP: Marcus Welby MD

Protocol: Ankle and Foot Injury (Adult)

See PCP within 24 hours

No To All

Yes

No

[1] Limp when walking AND [2] due to a twisted ankle or foot

R/O: sprain, minor fracture

[1] Limp when walking AND [2] due to a direct blow or crushing injury

R/O: contusion, minor fracture

Large swelling or bruise (> 2 inches or 5 cm)

R/O: minor fracture, muscle hematoma

Diabetes (Exception: small cut or scrape)

Reason: diabetic neuropathy reduces pain of fracture

[1] High-risk adult (e.g., age > 60, osteoporosis, chronic steroid use) AND [2] limping

Reason: there is greater risk of fracture in patients with osteoporosis (bone thinning)

[1] No prior tetanus shots (or is not fully vaccinated) AND [2] any wound (e.g., cut, scrape)

Note: A full tetanus vaccination series consists of 3 shots. Nearly all adults born in North America have received a full 3-tetanus shot series in childhood.

[1] HIV positive or severe immunodeficiency (severely weak immune system) AND [2] DIRTY cut

Reason: may need Tetanus Immune Globulin (TIG). Referral to the emergency department may likely be required as doctors' offices usually do not stock TIG.

BACK

NEXT

5. The nurse completes the call by providing Care Advice. The appropriate care advice for each disposition is already mapped in the database.

Pathways Call Center Software

Patient

Profile

Encounter History

Triage Process

Reason for Call

Triage Questions

Care Advice

Encounter Summary

Resources

Background

First Aid

Powered by:

Schmitt-Thompson Clinical Content

Care Advice

Kathy Johnson RN

Patient Name: John Anderson

Caller Name: Self

Encounter Date: 6/1/2021

Age: 24

Relationship: Self

Gender at Birth: M

PCP: Marcus Welby MD

Protocol: Ankle and Foot Injury (Adult)

Question: Large swelling or bruise(> 2 inches or 5 cm)

Given Now

Send Later

Apply a Cold Pack

- Apply a cold pack or an ice bag (wrapped in a moist towel) to the area for 20 minutes. Repeat in 1 hour, then every 4 hours while awake.
- Continue this for the first 48 hours after an injury.
- This will help decrease pain and swelling.

Pain Medicines

- For pain relief, you can take either acetaminophen, ibuprofen, or naproxen.
- They are over-the-counter (OTC) pain drugs. You can buy them at the drugstore.
- Acetaminophen - Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.
- Naproxen (e.g., Aleve):** Take 220 mg (one 220 mg pill) by mouth every 8 hours as needed. You may take 440 mg (two 220 mg pills) for your first dose. The most you should take each day is 660 mg (three 220 mg pills a day), unless your doctor has told you to take more.

Pain Medicines - Extra Notes and Warnings

- Use the lowest amount of medicine that makes your pain better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can

BACK

DONE

6. An Encounter Summary of the call has been prepared recapping the salient points of the discussion. This can then be saved to the patient record.

Pathways Call Center Software

Patient

Profile

Encounter History

Triage Process

Reason for Call

Triage Questions

Care Advice

Encounter Summary

Resources

Background

First Aid

Powered by:

Schmitt-Thompson

Clinical Content

Encounter Summary

Kathy Johnson RN

Patient Name: John Anderson

Age: 24

Gender at Birth: M

Caller Name: Self

Relationship: Self

PCP: Marcus Welby MD

Encounter Date: 6/1/2021

Date

6/1/2021

Reason for Call

Caller playing basketball, jumped and fell on ankle.

Guideline Used

Ankle and Foot Injury

Disposition

See PCP within 24 Hours

Positive Triage Assessment Question

Large swelling or bruise(> 2 inches or 5 cm)

Care Advice Provided

Apply a Cold Pack

Pain Medicines

Call Back If:

- Pain lasts longer than 2 hours
- You become worse

BACK

SAVE

PRINT

- Image of the transaction being recorded in the caller's patient record or history.

Pathways Call Center Software

Call Queue

New Encounter

Patient Profiles

History

John Anderson

AGE: 24 GENDER: Male

Date	Encounter Type	Encounter Description
6/1/2021	Phone	Caller playing basketball, jumped and fell on ankle.
7/22/2020	Phone	Diarrhea and stomach ache for two days.
1/19/2020	Office Visit	Annual well visit.
10/1/2019	ER	Stung by bees 2 hours ago and have hives, itching, and swelling all over body.