Questions for your oncologist

Synthetic demo data

Discuss these findings with your care team.

This is a discussion aid for you and your care team. It is not medical advice, and it cannot tell you that you qualify for a trial. Only your oncologist can do that.

- **9** strong matches
- **277** trials reviewed
- **198** other trials to ask about, **70** unlikely

01 **Quaratusugene Ozeplasmid (Reqorsa) and Osimertinib in Patients With Advanced Lung Cancer Who Progressed on Osimertinib**

Phase 1/2 Canton, OH, 103 miles

NCT04486833 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *The NSCLC must be epidermal growth factor receptor (EGFR) mutation positive-positive [sic] based on results from most recent tissue biopsy or most recent evaluation of circulating tumor DNA.*

Requires stage IIIA, IIIB, IIIC, IV, IVA, or IVB: chart shows stage IV.

Trial's own words

> *Stage III or IV NSCLC or recurrent NSCLC that is not potentially curable by radiotherapy or surgery.*

Requires ECOG 0 or 1: chart shows ECOG 1.

Trial's own words

> *Eastern Cooperative Oncology Group performance status (ECOG PS) score from 0 to 1.*

ANC ≥ 1500/µL: chart shows 3200/µL.

Trial's own words

> *Absolute neutrophil count (ANC) >1500/mm3, platelet count >100,000/mm3 within ≤28 days.*

Platelets ≥ 100000/µL: chart shows 210000/µL.

Trial's own words

> *Absolute neutrophil count (ANC) >1500/mm3, platelet count >100,000/mm3 within ≤28 days.*

Creatinine ≤ 1.5 mg/dL: chart shows 0.9 mg/dL.

Trial's own words

> *Adequate renal function documented by serum creatinine of ≤1.5 mg/dL or calculated creatinine clearance >50 ml/min within ≤28 days.*

Total bilirubin ≤ 1.5 × ULN: chart shows 0.6 mg/dL.

Trial's own words

> *Adequate hepatic function as documented by serum bilirubin <1.5 mg/dL and aspartate aminotransferase (AST) and alanine aminotransferase (ALT) ≤2.5 X upper limit of normal (ULN) within ≤28 days.*

▶

Not met (quoted)

▶

Discuss with your care team

02 **A Study to Assess Adverse Events and Change in Disease Activity of Intravenous (IV) Telisotuzumab Adizutecan, Monotherapy or in Combination With Osimertinib, Compared to...**

Phase 2/3 Fort Wayne, IN, 138 miles

NCT07155187 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

Ask your oncologist

- Have you had prior EGFR inhibitor treatment? The trial mentions it.
- Have you had prior EGFR inhibitor treatment? The trial mentions it.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Histologically or cytologically confirmed diagnosis of locally advanced or metastatic non-squamous non-small cell lung cancer (NSCLC) with documented EGFR Exon 19 deletion or Exon21 L858R mutation.*

Requires ECOG 0 or 1: chart shows ECOG 1.

Trial's own words

> *Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 to 1.*

▶

Not met (quoted)

▶

Discuss with your care team

03 **Study of Izalontamab Brengitecan (BMS-986507) Versus Platinum-Pemetrexed for EGFR-mutated Non-small Cell Lung Cancer After Failure of EGFR TKI Therapy (IZABRIGHT-Lung01)**

Phase 2/3 Columbus, OH, 0 miles

NCT07100080 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Documented evidence of EGFR mutation (exon 19 deletion, L858R mutation).*

▶

Not met (quoted)

▶

Discuss with your care team

04 **First in Human Study of BAY2927088 in Participants Who Have Advanced Non-small Cell Lung Cancer (NSCLC) With Mutations in the Genes of Epidermal Growth Factor Receptor (EGFR...**

Phase 1/2 Detroit, MI, 164 miles

NCT05099172 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

Ask your oncologist

- Have you had prior at least one prior systemic therapy for advanced disease? The trial excludes it.
- Have you had prior an EGFR tyrosine kinase inhibitor (TKI) ≤ 8 days or 5x the terminal phase? The trial mentions this therapy.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Documented activating EGFR and/or HER2 mutation assessed by a Clinical Laboratory Improvement Amendments (CLIA)-certified (United States [US] sites) or an equally accredited (outside of the US) local laboratory.*

Requires ECOG 0 or 1: chart shows ECOG 1.

Trial's own words

> *Eastern Cooperative Oncology Group (ECOG) performance status 0 or 1.*

Hemoglobin ≥ 9 g/dL: chart shows 12.4 g/dL.

Trial's own words

> *Hemoglobin ≥ 9.0 g/dL. Criteria must be met without erythropoietin dependency and without packed red blood cell (pRBC) transfusion within 2 weeks prior to testing.*

▶

Not met (quoted)

▶

Discuss with your care team

05 **A Clinical Study to Test if an Investigational Treatment Called BNT326 is Safe and Potentially Beneficial When Used Alone or in Combination With Other Investigational Treatments Such as...**

Phase 1/2 Cleveland, OH, 126 miles

NCT07070232 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Have documented positive test results for an EGFR-sensitizing mutation (EGFR-sensitizing mutation Exon 21-L858R and 19del).*

Requires ECOG 0 or 1: chart shows ECOG 1.

Trial's own words

> *Have ECOG performance status of 0 or 1.*

▶

Not met (quoted)

▶

Discuss with your care team

06 **Study of NXP900 With Osimertinib in Subjects With Advanced, EGFR-Mutated Non-Small Cell Lung Cancer**

Phase 1 Pittsburgh, PA, 162 miles

NCT07315113 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

Ask your oncologist

- Have you had prior osimertinib? The trial mentions this therapy.
- Have you had prior osimertinib? The trial mentions this therapy.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *EGFR-Mutated*

Requires ECOG 0 or 1: chart shows ECOG 1.

Trial's own words

> *Eastern Cooperative Oncology Group (ECOG) performance status of 0-1.*

▶

Not met (quoted)

▶

Discuss with your care team

07 **A Study of BH-30643 in Subjects With Locally Advanced or Metastatic NSCLC Harboring EGFR and/or HER2 Mutations**

Phase 1/2 Detroit, MI, 164 miles

NCT06706076 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Harboring EGFR*

Requires ECOG 0 or 1: chart shows ECOG 1.

Trial's own words

> *Eastern Cooperative Oncology Group Performance Status ≤ 1.*

▶

Not met (quoted)

▶

Discuss with your care team

08 **Comparing Combinations of Targeted Drugs for Advanced Non-Small Cell Lung Cancer That Has EGFR and MET Gene Changes (A Lung-MAP Treatment Trial)**

Phase 2 Columbus, OH, 0 miles

NCT05642572 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

Ask your oncologist

- Have you had prior strong inducers of CYP3A4? The trial excludes it.
- Have you had prior an anti-VEGF or VEGFR inhibitor or MET inhibitor? The trial excludes it.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Participants must have documentation of NSCLC with a sensitizing EGFR mutation*

eGFR ≥ 50 mL/min: chart shows 85 mL/min.

Trial's own words

> *Participants must have a serum creatinine =< the IULN OR calculated creatinine clearance >= 50 mL/min using the following Cockcroft-Gault Formula.*

▶

Not met (quoted)

▶

Discuss with your care team

09 **Prospective External Control Cohort In Patients With Newly Diagnosed Advanced or Metastatic EGFR Mutant Non-Small Cell Lung Cancer**

Elk Grove Village, Illinois, 295 miles

NCT07028489 on ClinicalTrials.gov

Strong evidence Strong. Discuss with your care team.

Ask your oncologist

- Have you had prior osimertinib? The trial mentions this therapy.
- Have you had prior osimertinib? The trial mentions this therapy.

▼

Met (quoted)

Requires an EGFR alteration: chart: EGFR exon 19 deletion.

Trial's own words

> *Evidence of a confirmed diagnosis of EGFR mutant (EGFR exon 19del or EGFR exon 21 L858R) locally advanced or metastatic NSCLC diagnosed on Jan 1st 2024 or later.*

▶

Not met (quoted)

▶

Discuss with your care team

Discuss these findings with your care team.

▶ Other trials to ask about (198)

▶ Show 70 unlikely trials

- Would any of these NCT IDs be worth discussing for me?
- Is there a criterion marked unknown that you already know from my chart?
- If a trial looks close, what would we need to confirm before a referral?

Trial records come from ClinicalTrials.gov, a service of the National Library of Medicine, National Institutes of Health. ClinicalTrials.gov last processed these records at 2026-11-10T10:00:05 (shown in the source timezone from the API). We keep only regional sites and add a local stage flag. Those are our modifications, not ClinicalTrials.gov fields.