


ENM 360: Introduction to Data-driven Modeling

Lecture #6: Statistical estimation



Paris Perdikaris
September 21, 2021



Correlation and linear dependence




The multivariate Gaussian



$$p(\mathbf{x} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-\frac{D}{2}} |\boldsymbol{\Sigma}|^{-\frac{1}{2}} \exp \left(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right)$$


The multivariate Gaussian



Notation	$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$
Parameters	$\boldsymbol{\mu} \in \mathbf{R}^k$ — location $\boldsymbol{\Sigma} \in \mathbf{R}^{k \times k}$ — covariance (positive semi-definite matrix)
Support	$\mathbf{x} \in \boldsymbol{\mu} + \text{span}(\boldsymbol{\Sigma}) \subseteq \mathbf{R}^k$
PDF	$\det(2\pi\boldsymbol{\Sigma})^{-\frac{1}{2}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}$, exists only when $\boldsymbol{\Sigma}$ is positive-definite
Mean	$\boldsymbol{\mu}$
Mode	$\boldsymbol{\mu}$
Variance	$\boldsymbol{\Sigma}$


Marginals and conditionals of a Gaussian

$$p(\mathbf{x}, \mathbf{y}) = \mathcal{N} \left(\begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}, \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{bmatrix} \right)$$



Marginal distribution

$$p(\mathbf{x}) = \int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} = \mathcal{N}(\mathbf{x} \mid \mu_x, \Sigma_{xx})$$




Conditional distribution

$$p(\mathbf{x} \mid \mathbf{y}) = \mathcal{N}(\mu_{x \mid y}, \Sigma_{x \mid y})$$

$$\mu_{x \mid y} = \mu_x + \Sigma_{xy} \Sigma_{yy}^{-1} (\mathbf{y} - \mu_y)$$


$$\Sigma_{x \mid y} = \Sigma_{xx} - \Sigma_{xy} \Sigma_{yy}^{-1} \Sigma_{yx}.$$



These are unique properties that make the Gaussian distribution very simple and attractive to compute with! It is essentially our main building block for computing under uncertainty.

Maximum likelihood estimation

$$\theta_{\text{MLE}} = \arg \max_{\theta \in \Theta} p(\mathcal{D}|\theta)$$



Bayesian estimation

$$p(\theta|\mathcal{D}) = \frac{p(\mathcal{D}|\theta)p(\theta)}{p(\mathcal{D})}$$

